

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,400

Open access books available

133,000

International authors and editors

165M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# Maintenance Of Calcium Homeostasis in Embryonic Stem Cell-Derived Cardiomyocytes

Iek Chi Lo<sup>1</sup>, Chun Kit Wong<sup>1</sup> and Suk Ying Tsang<sup>1,2,3,4\*</sup>

<sup>1</sup>School of Life Sciences, The Chinese University of Hong Kong

<sup>2</sup>State Key Laboratory of Agrobiotechnology, The Chinese University of Hong Kong

<sup>3</sup>Stem Cell and Regeneration Program, School of Biomedical Sciences,  
The Chinese University of Hong Kong

<sup>4</sup>Key Laboratory for Regenerative Medicine, Ministry of Education,  
The Chinese University of Hong Kong  
Hong Kong Special Administrative Region

## 1. Introduction

Myocardial infarction is one of the major causes of morbidity and mortality in many developed countries. A potential method for treatment of such disease is the cell replacement therapy which involves the transplantation of cardiomyocytes (CMs). However, CMs are of very limited supply. Embryonic stem cells (ESCs) isolated from the inner cell mass of blastocysts are capable of self-renewal and can differentiate into all cell lineages, including CMs (He et al., 2003; Kehat et al., 2002; Kehat et al., 2001; Moore et al., 2008; Mummery et al., 2003; Ng et al., 2010; Thomson et al., 1998; Xu et al., 2002; Xue et al., 2005). Therefore, ESCs can be an excellent source of CMs for regenerative medicine.

Calcium ( $\text{Ca}^{2+}$ ) is a universal signaling molecule that regulates a wide variety of cellular functions. In CMs, intracellular  $\text{Ca}^{2+}$  concentration ( $[\text{Ca}^{2+}]_i$ ) plays an important role in the contraction and relaxation of CMs. The  $[\text{Ca}^{2+}]_i$  is tightly regulated by many proteins, including ion channels, receptors, pumps, and exchangers that are located on the cell surface plasma membrane and on the sarcoplasmic reticulum (SR). The aim of this book chapter is to provide a thorough review on  $\text{Ca}^{2+}$  handling in ESC-derived CMs.

For each protein of interest, some basic information on the protein was firstly presented. Then, changes in the expression of the protein and their contribution to the  $\text{Ca}^{2+}$  homeostasis and  $\text{Ca}^{2+}$  transient as described in human (h) and/or mouse (m) ESC studies were presented. *In vivo* data in mouse embryo studies were also presented for comparison purposes.

The information reviewed in this chapter would be important not only for understanding the basic biology of early differentiating CMs, it would also be important for providing insights into the future uses of ESC-derived CMs for cell replacement therapies.

### 2.1 Regulation of intracellular calcium level by voltage-operated calcium channels

Voltage-operated  $\text{Ca}^{2+}$  channels are typically composed of five subunits, namely  $\alpha_1$ ,  $\alpha_2$ ,  $\beta$ ,  $\delta$  and  $\gamma$  (Catterall, 1995; Catterall, 2000; De Waard et al., 1996; Dolphin, 2006; Moreno Davila,

---

\* Corresponding author.

1999). The  $\alpha_1$  subunit forms the ion conduction pore of the channels while the other subunits modulate the functions of the channels. The  $\alpha_1$  subunit is itself a tetramer of proteins, each of which consists of six transmembrane segments (S1-S6). This subunit confers all of the major properties to voltage-operated  $\text{Ca}^{2+}$  channels in that it contains a voltage sensor at segment S4 and forms a  $\text{Ca}^{2+}$ -selective pore at segments S5-S6. The other subunits are regarded as to providing ancillary functions to the channels. The  $\alpha_2$  and  $\delta$  subunits are two products of the same gene; the protein is cleaved into two peptides following translation. Subsequent disulfide bond formation anchors the extracellular  $\alpha_2$  subunit to the membrane via the transmembrane  $\delta$  subunit. The  $\beta$  subunit is a cytoplasmic protein; it contains an  $\alpha_1$ -binding pocket, where specific amino acids in the linker region between segments S1 and S2 of the  $\alpha_1$  subunit bind (Richards et al., 2004). The  $\gamma$  subunit is a monomer, which consists of four transmembrane segments (S1-S4). A diverse range of functionally distinct voltage-operated  $\text{Ca}^{2+}$  channels are formed by the combinations of different isoforms that have been identified for each of these subunits. The  $\alpha_1$  subunit is encoded by a family of ten distinct genes (the  $\text{Cav}$  gene family), which can be divided into three sub-families;  $\text{Cav}1$  consists of  $\text{Cav}1.1-1.4$ ,  $\text{Cav}2$  consists of  $\text{Cav}2.1-2.3$  and  $\text{Cav}3$  consists of  $\text{Cav}3.1-3.3$  (Catterall et al., 2005). The  $\alpha_2/\delta$  and  $\beta$  subunits are each encoded by four genes;  $\alpha_2/\delta_{1-4}$  and  $\beta_{1-4}$  respectively, while the  $\gamma$  subunit is encoded by eight genes;  $\gamma_{1-8}$  (Arikkath and Campbell, 2003; Yang et al., 2011). Voltage-operated  $\text{Ca}^{2+}$  channels function by regulating the entry of extracellular  $\text{Ca}^{2+}$  into cells. Two types of voltage-operated  $\text{Ca}^{2+}$  channels exist in adult CMs (Catterall et al., 2005; Ono and Iijima, 2005). These are T-type  $\text{Ca}^{2+}$  channels, which open in response to more negative membrane potentials at about  $>-70\text{mV}$  for very short durations of 0.5 to 2 milliseconds (transient activation), and L-type  $\text{Ca}^{2+}$  channels, which open in response to less negative membrane potentials at about  $>-30\text{mV}$  for relatively longer periods of 0.5 to 10 milliseconds (long-lasting activation) (De Waard et al., 1996).

### 2.1.1 T-type calcium channels

T-type  $\text{Ca}^{2+}$  channels could be the simplest type of voltage-operated  $\text{Ca}^{2+}$  channels known thus far. Its structure was initially predicted *in silico* by homology to other  $\alpha_1$  subunits known at the time (Perez-Reyes, 2006). Since expression of the synthetic products was able to reproduce most of the electrophysiological properties observed in its native form, T-type  $\text{Ca}^{2+}$  channels were thought to consist of only one  $\alpha_1$  subunit; the other ancillary subunits ( $\alpha_2$ ,  $\beta$ ,  $\delta$  and  $\gamma$ ) were considered absent. Although later co-expression studies could describe a role for the other ancillary subunits in the regulating functions of T-type  $\text{Ca}^{2+}$  channels *in vitro*, presence of these subunits have not been validated *in vivo* (Dolphin et al., 1999; Dubel et al., 2004; Green et al., 2001; Hobom et al., 2000; Lacinova and Klugbauer, 2004). Therefore, further investigations will be required to ascertain the genuine structure of this type of voltage-operated  $\text{Ca}^{2+}$  channel.

The  $\alpha_1$  subunit of T-type  $\text{Ca}^{2+}$  channels is encoded by either one of the three  $\text{Cav}3$  genes, among which  $\text{Cav}3.1$  and  $\text{Cav}3.2$  are expressed in sinoatrial (SA) nodal cells of the heart (Bohn et al., 2000; Perez-Reyes, 2003). The SA node is the primary site, where spontaneous rhythmic action potentials are initiated. At the beginning of each action potential in an SA nodal cell, an influx of extracellular  $\text{Na}^+$  into the cytosol, known as the funny current ( $I_f$ ), first depolarizes the cell to about  $-50\text{mV}$ , at which T-type  $\text{Ca}^{2+}$  channels on the membrane open to allow an influx of extracellular  $\text{Ca}^{2+}$  into the cytosol. This produces a T-type  $\text{Ca}^{2+}$  current ( $I_{\text{Ca, T}}$ ), which further depolarizes the cell to about  $-40\text{mV}$ . L-type  $\text{Ca}^{2+}$  channels on the membrane then open at this membrane potential to allow a greater influx of extracellular

Ca<sup>2+</sup> into the cytosol and produce an L-type Ca<sup>2+</sup> current ( $I_{Ca, L}$ ) to a membrane potential until the threshold to produce an action potential is reached. Action potentials generated in the SA nodal cells are conducted to CMs within the working myocardium via gap junctions, ultimately resulting in regular contractions of the adult heart.

Ca<sub>v</sub>3.1 mRNA and protein were simultaneously detected as early as embryonic day 14 (E14) after the heart has fully matured *in vivo* within mouse embryos (Cribbs et al., 2001). In a separate experiment, the mRNA level of Ca<sub>v</sub>3.1 was moderately reduced from E18 when compared to its expression at adult stage (Yasui et al., 2005). Ni<sup>2+</sup> at a high concentration of 100 μM could be used to selectively block 50% of Ca<sub>v</sub>3.1 T-type channels. The application of Ni<sup>2+</sup> at this concentration was found to reduce  $I_{Ca, T}$  by about 40% in CMs derived from mouse embryos at E12.5, indicating that the Ca<sub>v</sub>3.1 T-type channels were functional by this time-point (Cribbs et al., 2001). In addition, homozygous null Ca<sub>v</sub>3.1<sup>-/-</sup> mice displayed bradycardia, i.e. a reduced rate of cardiac contraction (Mangoni et al., 2006). Expression of Ca<sub>v</sub>3.1 has also been studied *in vitro*. Its mRNA could be detected as early as day 5 and generally increased up until its last measurement at day 15 post-differentiation in ht7 mouse ESC- (mESC-) derived CMs (Mizuta et al., 2005). In R1 mESC-derived CMs, however, Ca<sub>v</sub>3.1 mRNA could only be detected at two copies per cell by day 12, with its expression reaching the highest level at day 23 and declining by day 34 post-differentiation; the expression pattern reflected the amplitudes of  $I_{Ca, T}$  measured at these time-points. (Zhang et al., 2003). A reduction of about 46% in Ca<sub>v</sub>3.1 expression from day 9.5 to day 23.5 post-differentiation was also reported in EMG7 mESC-derived CMs; this down-regulation was found to be associated with a decrease in the contraction rates of the CMs in that  $I_{Ca, T}$  was smaller in non-contracting myocytes at day 23.5 when compared to contracting CMs at day 9.5 and that the contraction rate at day 9.5 could be reduced via the application of 10 μM efonidipine, an L- and T-type blocker, to the CMs (Yanagi et al., 2007). These data suggested a role for Ca<sub>v</sub>3.1 in regulating the early contractions observed in the developing embryonic heart; by maintaining these contractions exclusively in developing pacemaker cells, other cells could then develop into non-contracting atrial or ventricular cells.

Ca<sub>v</sub>3.2 mRNA was detected as early as eight-week gestation in humans (Qu and Boutjdir, 2001). In mouse, the expression of Ca<sub>v</sub>3.2 was also measured at three time-points; its mRNA was detected as early as E9.5 and then its level was decreased by E18 until it was no longer detected at adult stage (Yasui et al., 2005). Again, Ca<sub>v</sub>3.2 has been studied *in vitro*. Its mRNA was detected as early as day 5, but peaked at day 6 and gradually declined and was still detectable up until the last measurement at day 15 post-differentiation in ht7 mESC-derived CMs (Mizuta et al., 2005). In line with this finding, Ca<sub>v</sub>3.2 mRNA level was also found to be down-regulated by about 24% from day 9.5 to day 23.5 post-differentiation in EMG7 mESC-derived CMs (Yanagi et al., 2007). Ni<sup>2+</sup> at a low concentration of <50 μM could be used to selectively block 50% of Ca<sub>v</sub>3.2 T-type channels. When Ni<sup>2+</sup> was applied at 40 μM to ht7 mESC-derived CMs at day 8 post-differentiation,  $I_{Ca, T}$  was found to be evidently decreased by about 60%, which signified the presence of functional Ca<sub>v</sub>3.2 T-type channels by this time-point. Homozygous null Ca<sub>v</sub>3.2<sup>-/-</sup> mice showed no sign of cardiac arrhythmia, but cardiac fibrosis with age-dependent severity (Chen et al., 2003). These data suggested a role for Ca<sub>v</sub>3.2 in regulating the growth and maturation of the developing embryonic heart.

### 2.1.2 L-type calcium channels

L-type Ca<sup>2+</sup> channels are distinguished from other voltage-operated Ca<sup>2+</sup> channels that open at high membrane potentials for long periods on the basis of their sensitivity to 1, 4-

dihydropyridines (DHPs) (Hess et al., 1984). DHP-binding sites are contained within the  $\alpha_1$  subunit of L-type  $\text{Ca}^{2+}$  channels, which are encoded by either one of the four genes of the  $\text{Ca}_v1$  sub-family. Among these,  $\text{Ca}_v1.1$  is expressed predominantly in skeletal myocytes,  $\text{Ca}_v1.2$  in adult CMs and  $\text{Ca}_v1.3$  exclusively in SA nodal and atrial cells of the heart (Catterall, 2000; Qu et al., 2005). Unlike T-type, L-type  $\text{Ca}^{2+}$  channels are typical voltage-operated  $\text{Ca}^{2+}$  channels, which are made up of five subunits. Together with one of each of the  $\alpha_2/\delta_{1-4}$ ,  $\beta_{1-3}$  and  $\gamma_{4, 6-8}$  ancillary subunits,  $\text{Ca}_v1.2$ - and  $\text{Ca}_v1.3$ -encoded  $\alpha_1$  proteins form the functional L-type  $\text{Ca}^{2+}$  channels that are found within the adult heart (Freise et al., 1999; Hosey et al., 1996; Yang et al., 2011).

$\text{Ca}_v1.2$  L-type  $\text{Ca}^{2+}$  channels are responsible for cardiac excitation-contraction coupling (E-C coupling), a mechanism by which an action potential is transformed into contraction in adult CMs. Upon the arrival of an action potential from SA nodal cells, L-type  $\text{Ca}^{2+}$  channels, which are clustered at transverse tubules (T-tubules), briefly open at the sarcolemma to allow an influx of extracellular  $\text{Ca}^{2+}$  into the cytosol of the CMs. This only accounts for about 30% of the increase in cytosolic  $\text{Ca}^{2+}$ , which is maintained and magnified through a process known as calcium-induced calcium release (CICR) (Bers, 2002). CICR is mediated via ryanodine receptor (RyR)  $\text{Ca}^{2+}$  release channels, which detect the  $I_{\text{Ca}, L}$  and open to release  $\text{Ca}^{2+}$  from its intracellular stores within the SR into the cytosol of the CMs. This rise enhances binding of  $\text{Ca}^{2+}$  to troponin within the CMs, ultimately bringing about contractions of the adult heart. Conversely,  $\text{Ca}_v1.3$  L-type  $\text{Ca}^{2+}$  channels are responsible for producing  $I_{\text{Ca}, L}$ , which constitutes one of the ionic currents to initiate action potentials at the SA node. Hence, these L-type  $\text{Ca}^{2+}$  channels contribute in maintaining the pacemaker activity of the adult heart (see 'T-type calcium channels' for details).

Consistent with its role in E-C coupling in the adult heart,  $\text{Ca}_v1.2$  mRNA was detected as early as eight-week gestation in humans, with its level culminating to its maximum at adult stage (Qu and Boutjdir, 2001).  $\text{Ca}_v1.2$  mRNA was also present *in vitro* at <40 days post-beating in H9.2 human ESC- (hESC-) derived CMs (Satin et al., 2008). In H7 hESC-derived CMs,  $\text{Ca}^{2+}$  transients that were detected at day 17 post-differentiation could be eliminated by the application of 10 $\mu\text{M}$  diltiazem, an L-type specific antagonist; this signified the functionality of L-type channels by this time-point (Zhu et al., 2009). Similar to humans, the expression of  $\text{Ca}_v1.2$  was also up-regulated from its initial detection at E9.5 when the earliest contractions were observed by about three-fold at E15.5 when the heart has fully matured in mouse embryos; its protein was detected by E12.5 in mouse embryos (Acosta et al., 2004; Xu et al., 2003). In mESCs, patch clamp and  $\text{Ca}^{2+}$  imaging experiments indicated the functional expression of the  $I_{\text{Ca}, L}$  starting from differentiation day 7, even before the appearance of spontaneous contractions (Kolossoff et al., 1998). The current density of  $I_{\text{Ca}}$  continued to increase to day 10, the day of beginning of spontaneous contractions in most differentiating embryoid bodies (EBs). Thereafter, similar current density was recorded on day 17, the last day of measurement (Kolossoff et al., 1998). Homozygous null  $\text{Ca}_v1.2^{-/-}$  mice died *in utero* by E14.5, probably due to an absence of CICR-regulated contractions, resulting in reduced oxygen supply to the embryos (Seisenberger et al., 2000). Despite the lethality observed, normal rhythmic contractions were sustained in the  $\text{Ca}_v1.2^{-/-}$  mice until E12.5. For this reason, CICR was initially thought to be non-essential for regulating the early cardiac contractions observed before E12.5 until later time-points. This conclusion was further supported by the fact that both the contractions and  $\text{Ca}^{2+}$  oscillations of D3 mESC-derived CMs were insensitive to the application of 50nM nisoldipine, a specific L-type channel blocker, at days 8-11 post-differentiation (Viatchenko-Karpinski et al., 1999). In addition,



these cells were insensitive to the application of high  $K^+$  when compared to D3 mESC-derived CMs at day 16 post-differentiation; exposure to high  $K^+$  at this stage would normally lead to immediate hyper-contraction and death of these cells. Albeit these findings, the notion that  $I_{Ca, L}$  was not required at earlier stages of development was overthrown when evidence for the existence of a compensatory mechanism emerged; the expression of  $Ca_v1.3$  was eminently up-regulated at E9.5 and E12.5 by four fold to produce a DHPR-insensitive  $I_{Ca, L}$  in  $Ca_v1.2^{-/-}$  mouse embryos when compared to wild-type (Xu et al., 2003). This has not only reinstated the importance of  $I_{Ca, L}$  in maintaining cardiac contractions, even at the embryonic stage of the heart, but also appointed a new role for  $Ca_v1.3$  to substitute for  $Ca_v1.2$  and preserve its role in CICR for E-C coupling in its absence. In contrast with  $Ca_v1.2$ ,  $Ca_v1.3$  mRNA was detected by E9.5, with its level elevated only by a modest amount at E15.5 in mouse embryos. Its protein was also detected by E12.5 in mouse embryos (Xu et al., 2003). Analogous to  $Ca_v3.1^{-/-}$  mice, homozygous null  $Ca_v1.3^{-/-}$  mice developed bradycardia, consistent with its primary role in regulating the pacemaker activity of the heart (Platzer et al., 2000; Zhang et al., 2002). In R1 mESC-derived CMs, the expression of L-type  $Ca^{2+}$  channels could be detected as early as day 7 post-differentiation; given that the application of  $1\mu M$  nifedipine, an L-type channel-specific antagonist, could eliminate  $Ca^{2+}$  transients that were normally observed at early (days 9-11) and intermediate (days 13-15), but only partially inhibit those at late (18-21 days post-differentiation) stages, this indicated that L-type channels were, in fact, playing a more dominant role at earlier developmental stages (Fu et al., 2006b).

## 2.2 Regulation of intracellular calcium level by ligand-operated calcium release channels

### 2.2.1 Ryanodine receptor (RyR) calcium release channels

RyR- $Ca^{2+}$  release channels of a molecular mass of greater than two Mega-Daltons probably form the largest ion channels known thus far (Lanner et al., 2010). RyR- $Ca^{2+}$  release channels are composed of homo-tetramers of RyR proteins. Owing to its enormous size, structural elucidation of these channels has been a major challenge. Nonetheless, RyR- $Ca^{2+}$  release channels have been predicted to be largely cytoplasmic, with an ion-conducting pore consisting of around four to twelve transmembrane segments.

Three mammalian isoforms of RyRs exist; RyR1 is predominantly expressed in skeletal myocytes, RyR2 in CMs and RyR3 in astrocytes. The isoforms differ in three particular regions, which have been named domains D1-3 (Ma et al., 2004). As mentioned earlier in this chapter, RyR- $Ca^{2+}$  release channels are induced to open in response to  $Ca^{2+}$  influx via L-type  $Ca^{2+}$  channels to facilitate E-C coupling in muscle cells. In skeletal myocytes, E-C coupling takes place through direct physical interaction between RyR1 and the D2 region, which is located between segments S2 and S3 of the  $Ca_v1.1$ -encoded  $\alpha_1$  subunit of L-type  $Ca^{2+}$  channels. In CMs, however, sequence divergence in the D2 domain between RyR1 and RyR2 means that RyR2 cannot physically interact with the  $Ca_v1.2$ -encoded  $\alpha_1$  subunit of L-type  $Ca^{2+}$  channels; E-C coupling can, therefore, only occur via CICR. RyR- $Ca^{2+}$  release channels are, thus, organized into large arrays at junctions between the SR and the sarcolemma, in close proximity to the  $Ca_v1.2$  L-type  $Ca^{2+}$  channels in adult CMs (Bers, 2004). Localized  $Ca^{2+}$  release events are referred to as  $Ca^{2+}$  sparks, which are collectively synchronized by  $I_{Ca, L}$  to produce large, whole cell  $Ca^{2+}$  transients in the CMs.

Release of  $Ca^{2+}$  from the SR stores greatly increases the amount of cytosolic  $Ca^{2+}$  available to bind troponins in adult CMs. Adult CMs are mostly occupied by bouquets of thick and thin

filaments. Thick filaments are composed of myosin II molecules, while thin filaments are made up of troponin, tropomyosin and actin molecules. Troponin is itself a globular complex, which consists of three subunits, namely troponin T (TnT), troponin C (TnC) and troponin I (TnI). The TnT subunit binds tropomyosin, which coils around strands of actin molecules in the thin filament. Binding of  $\text{Ca}^{2+}$  to the TnC subunit induces a conformational change in the TnI subunit, thereby removing the steric hindrance on actin from tropomyosin. Binding of myosin II to actin can then occur, inducing a conformational change in the complex. This pulls the actin-associated thin filament past the myosin II-associated thick filament, resulting in a contracting phenomenon in the CMs.

RyR2 protein was present as early as day 17 post-differentiation in H7 hESC-derived CMs (Zhu et al., 2009). In H9.2 hESC-derived CMs, the application of a puff of 10mM caffeine was able to induce  $\text{Ca}^{2+}$  release from the SR as early as day 2 post-beating; this signified the functionality of RyR-channels in these cells at this early time-point (Satin et al., 2008). In mouse embryos, RyR2 mRNA was detected as early as E8.5, the level of which continued to increase up until its last measurement at E16.5 (Rosemlit et al., 1999).  $\text{Ca}^{2+}$  release from the SR was inducible by the application of 10mM caffeine, indicating the functionality of RyR2 channels by E8.5 in mouse embryos. In a different study, RyR2 protein was also detected at E18 and the detection persisted until adult stage (Liu et al., 2002). Homozygous null RyR2<sup>-/-</sup> mice died at E10, displaying morphological abnormalities in the heart tube (Takeshima et al., 1998). RyR2 expression has also been studied *in vitro* in mESC-derived CMs. RyR2 mRNA and protein were detected as early as day 5 and day 9 post-differentiation in mESC-derived CMs respectively (Boheler et al., 2002; Fu et al., 2006a). Immunohistochemistry revealed a continuous increase in RyR immunofluorescence intensity in differentiation day 15-25 CMs compared with differentiation day 8-11 CMs, suggesting an increasing density of RyRs during cardiac differentiation (Sauer et al., 2001). The application of 10mM caffeine elicited  $\text{Ca}^{2+}$  transients, the amplitudes of which increased from day 8 to day 17 post-differentiation in mESC-derived CMs (Kapur and Banach, 2007). Hence, RyR2 channels were functional by day 8 post-differentiation in mESC-derived CMs. RyR2<sup>-/-</sup> knockout R1 mESC-derived CMs exhibited no difference in the amplitudes of  $\text{Ca}^{2+}$  transients but contractions at a reduced rate from early (days 9-11) to intermediate (days 13- 15) to late (days 18-21 post-differentiation) differentiation stages when compared with wild-type (Fu et al., 2006a; Yang et al., 2002). The applications of 10 $\mu$ M ryanodine and 10mM caffeine were able to inhibit and induce  $\text{Ca}^{2+}$  release from the SR respectively. Cardiac differentiation was not affected, as indicated by the number of contracting colonies present in the differentiating cultures. Both effects of ryanodine and caffeine were seen to increase with time post-differentiation, indicating that SR  $\text{Ca}^{2+}$  loads increased during differentiation (Fu et al., 2006a; Sauer et al., 2001). Altogether, these findings suggested a role for RyR to mediate SR  $\text{Ca}^{2+}$  release, thereby regulating the rate of the earliest contractions observed in the developing embryonic heart, and that this regulation increases with differentiation.

### 2.2.2 Inositol 1, 4, 5-trisphosphate Receptor (IP<sub>3</sub>R) calcium release channels

IP<sub>3</sub>R  $\text{Ca}^{2+}$  release channels represent a type of enigmatic intracellular  $\text{Ca}^{2+}$  channels in that both its structure and function(s) are not fully understood (Foskett et al., 2007; Taylor et al., 2004; Taylor and Tovey, 2010; Yule et al., 2010). IP<sub>3</sub>R- $\text{Ca}^{2+}$  release channels can be formed from either homo- or hetero-tetramers of IP<sub>3</sub>R proteins. Each subunit of IP<sub>3</sub>R proteins contains an IP<sub>3</sub>-binding domain at its N-terminus and six transmembrane segments (S1-6) at its C-terminus; the  $\text{Ca}^{2+}$ -selective pore is formed at segments S5-6. Three mammalian

isoforms of IP<sub>3</sub>R exist, namely IP<sub>3</sub>R1, IP<sub>3</sub>R2 and IP<sub>3</sub>R3. These isoforms differ in their binding affinities for IP<sub>3</sub>, with IP<sub>3</sub>R2 being the most and IP<sub>3</sub>R3 being the least sensitive to the ligand (Iwai et al., 2007; Tu et al., 2005). All three IP<sub>3</sub>Rs are expressed in human adult CMs, but their subcellular localizations have not been studied (Nakazawa et al., 2011; Uchida et al., 2010). In rat adult CMs, these are expressed at about 50-fold lower than those of RyRs; IP<sub>3</sub>R1s are localized around the nuclear envelope and SR, both of which are connected, while IP<sub>3</sub>R2s are dispersed throughout the cytosol in a punctate pattern (Bare et al., 2005; Li et al., 2005; Moschella and Marks, 1993). Thus far, the role of IP<sub>3</sub>R in the adult heart has not been defined, but an up-regulation in their expression has been associated with conditions of cardiac failure in that the mRNA levels of IP<sub>3</sub>R1 and IP<sub>3</sub>R2 were increased by 123% and 93% respectively in failing compared with normal heart tissues (Ai et al., 2005; Go et al., 1995).

Activation of IP<sub>3</sub>R-Ca<sup>2+</sup> release channels is regulated by both the concentration of cytosolic Ca<sup>2+</sup> ions and the binding of IP<sub>3</sub> to one or more of the IP<sub>3</sub>-binding domains (Iino, 1990; Taylor and Laude, 2002). Low concentrations of cytosolic Ca<sup>2+</sup> ions are known to activate pore opening, while high concentrations inhibit it. IP<sub>3</sub> is a second messenger that is generated via a G protein-coupled receptor-mediated signal transduction pathway. G protein-coupled receptors are transmembrane receptors that are able to sense external stimuli and relay these signals into the cell via their interactions with cytosolic G proteins. G proteins are heterotrimeric complexes, which consist of a GDP-bound G<sub>α</sub> subunit and a G<sub>βγ</sub> dimer. G protein-coupled receptors are classified into different types according to the isoform of the G<sub>α</sub> subunits contained within their G protein-interacting partners. Generation of IP<sub>3</sub> involves the stimulation of a G<sub>q</sub> type of G protein-coupled receptors. Upon stimulation with their agonists, a conformational change is induced in the G<sub>q</sub> protein-coupled receptors. This causes an exchange of GDP for GTP in the G<sub>α</sub> subunits, which then dissociate from their G<sub>q</sub> protein complexes. GTP-bound G<sub>α</sub> subunits of G<sub>q</sub> protein complexes are responsible for activating phospholipase C. It is this enzyme, which hydrolyses phosphatidylinositol 4, 5-bisphosphate (PIP<sub>2</sub>), a phospholipid component of the plasma membrane, to ultimately yield IP<sub>3</sub> and another second messenger, diacylglycerol (DAG). IP<sub>3</sub> then binds to and activates IP<sub>3</sub>R-Ca<sup>2+</sup> release channels.

IP<sub>3</sub>R mRNA was present at detectable levels in D3 mESCs at the undifferentiated state (Yanagida et al., 2004). Application of 5μM ATP generated a Ca<sup>2+</sup> transient in the undifferentiated D3 mESCs, which was inhibited by pre-treatment with 75μM 2-APB, an IP<sub>3</sub>R blocker. This verified the functionality of IP<sub>3</sub>R-channels in these cells. In H9.2 hESCs, immunostaining indicated that both IP<sub>3</sub>R1 and IP<sub>3</sub>R2 were expressed in hESC-derived CMs (Sedan et al., 2010; Sedan et al., 2008); and the expression of IP<sub>3</sub>R2 was shown to gradually decline with maturation as revealed by quantitative RT-PCR (Satin et al., 2008). The expression of IP<sub>3</sub>R was also tested during development *in vivo*; its mRNA was first measured and detected as early as E5.5 in mouse embryos; it was present at high levels by E8.5 and continued to increase until E16.5 when its level started to drop (Rosemlit et al., 1999). These were demonstrated to be functional at two time-points; application of 5μM of IP<sub>3</sub> was able to induce Ca<sup>2+</sup> release from the SR in CMs derived from mouse embryos at E5.5 and E8.5. Application of 5μM xestospongine C, an IP<sub>3</sub>R antagonist, to mouse embryos at E10 diminished its Ca<sup>2+</sup> spiking; washing the drug out allowed slow recovery of this spiking activity (Mery et al., 2005). Likewise, application of 5μM xestospongine C also abrogated the Ca<sup>2+</sup> spiking activity observed in CGR8 mESC-derived CMs at day 8-10 post-differentiation. In addition, during the whole course of cardiac differentiation of R1 mESCs, 2-APB



decreased both the amplitude and upstroke velocity of  $\text{Ca}^{2+}$  transients, with the inhibitory effect decreased as differentiation proceeded, suggesting that  $\text{IP}_3\text{R}$  contributes to the  $\text{Ca}^{2+}$  transient and its effect decreases with differentiation (Fu et al., 2006b). In an attempt to discriminate between the roles of RyR- and  $\text{IP}_3\text{R}$ - channels, 50 $\mu\text{M}$  ryanodine was first used to block RyR-channels before the application of 5-20 $\mu\text{M}$   $\text{IP}_3\text{-AM}$  to CMs derived from mouse embryos at E10 (Rapila et al., 2008). This was able to induce a  $\text{Ca}^{2+}$  leak from the SR of the CMs in a concentration-dependent manner, whereby the slopes of the  $\text{Ca}^{2+}$  transients were elevated, despite its frequency stayed unchanged. In the absence of ryanodine, however, application of 10 $\mu\text{M}$   $\text{IP}_3\text{-AM}$  led to an increase in the frequency of  $\text{Ca}^{2+}$  transients and, hence, increased contractions in the CMs at E10. These data suggested a role for  $\text{IP}_3\text{R}$  in regulating the rate of the earliest contractions observed in the developing embryonic heart, perhaps by providing a source of  $\text{Ca}^{2+}$  to bind RyRs so as to increase its open probability for greater  $\text{Ca}^{2+}$  release from the SR, as suggested by Rapila *et al.* (Rapila et al., 2008) In support of this finding, genetic knockdown of  $\text{IP}_3\text{R}$  in CGR8 mESC-derived pacemaker cells resulted in weak and infrequent contractions, although differentiation was not affected, whilst mice over-expressing the  $\text{IP}_3\text{R2}$  gene developed mild cardiac hypertrophy by three months of age (Mery et al., 2005; Nakayama et al., 2010).

### 2.3 Regulation of intracellular calcium level by exchanger and pump

As previously mentioned, upon arrival of cardiac action potential, activation of L-type  $\text{Ca}^{2+}$  channel followed by CICR increases  $[\text{Ca}^{2+}]_i$  for the contraction process. Subsequently, the excess  $\text{Ca}^{2+}$  has to be removed in order to initiate relaxation. In CMs, this  $\text{Ca}^{2+}$  removal process is mediated by different  $\text{Ca}^{2+}$  extrusion mechanisms via the action of sodium-calcium exchanger (NCX) and sarcoplasmic/endoplasmic reticulum  $\text{Ca}^{2+}$ -ATPase (SERCA).

#### 2.3.1 Sodium-calcium exchanger (NCX)

NCX contains 9 transmembrane segments with a large loop at the cytoplasmic side (Philipson and Nicoll, 2000). Mutation experiments demonstrated the importance of  $\alpha$ -helices in the transmembrane segments for the transport activity of NCX (Nicoll et al., 1996). The large intracellular loop is found to be not essential for the transport; however, it is important for catalyzing the ion translocation reaction and has important regulatory functions (Philipson and Nicoll, 2000). This loop consists of multiple regulatory sites, including the regulatory  $\text{Ca}^{2+}$  binding site and the exchanger inhibitory peptide (XIP) region. The activity of NCX is known to be regulated by at least 4 factors, including  $\text{Ca}^{2+}$  (DiPolo, 1979),  $\text{Na}^+$  (Hilgemann et al., 1992),  $\text{PIP}_2$  (Hilgemann, 1990), and phosphorylation (Iwamoto et al., 1996). Binding of  $\text{Ca}^{2+}$  to the  $\text{Ca}^{2+}$  binding site is required for activating the  $\text{Na}^+\text{-Ca}^{2+}$  exchange activity (Giladi et al., 2010; Ottolia et al., 2004; Wu et al., 2010). On the other hand, the XIP region is responsible for the  $\text{Na}^+$ -dependent inactivation and is involved in the elimination of the  $\text{Na}^+$ -dependent inactivation process by  $\text{PIP}_2$  (Matsuoka et al., 1997). The activity of NCX is also regulated by phosphorylation, with stronger phosphorylation leading to higher NCX activity (Reppel et al., 2007a). Requirements of direct phosphorylation for up-regulation of NCX function by PKA and PKC are still in debate. A recent study by Wanichawan *et al.* demonstrated that the PKA phosphorylation site in full-length NCX1 is inaccessible, suggesting that NCX1 is not a direct substrate of PKA (Wanichawan et al., 2011). On the other hand, another study showed that the activity of NCX1 is dependent on PKC, although direct phosphorylation by PKC is not required (Iwamoto et al., 1998). NCX exists in 3 different isoforms, namely NCX1, NCX2 and NCX3.

NCX1 is referred as the cardiac NCX isoform as it is highly expressed in CMs but only in a lesser extent in other tissues such as brain and kidney (Lee et al., 1994).

NCX is classified as a secondary active transporter, which uses the energy stored in the electrochemical gradient in  $\text{Na}^+$  to extrude  $\text{Ca}^{2+}$  out of the cells, while the electrochemical gradient of  $\text{Na}^+$  is maintained by the  $\text{Na}^+/\text{K}^+$ -ATPase. Under normal condition, NCX operates in the forward mode in which it constitutively brings 3  $\text{Na}^+$  into the cells and extrudes 1  $\text{Ca}^{2+}$  in each translocation cycle. The forward mode is stimulated in response to a rise in  $[\text{Ca}^{2+}]_i$ , and it serves to bring  $[\text{Ca}^{2+}]_i$  back to normal level. In CMs, the primary role of NCX is to extrude  $\text{Ca}^{2+}$  after excitation under normal physiological conditions (Philipson and Nicoll, 2000). Some studies also showed that NCX functions in shaping the cardiac action potential. Application of NCX blocker KB-R7943 leads to shortening of plateau phase of cardiac potential (Spencer and Sham, 2003); similarly, CMs from NCX knockout mice also has a shorter AP when compare to wild-type mice (Pott et al., 2005), while induced over-expression of NCX leads to a longer plateau (Wang et al., 2009). On the other hand, NCX also operates in the reverse mode in response to membrane depolarization in CMs. During the depolarization phase of action potential when the  $[\text{Ca}^{2+}]_i$  has not reached the peak of  $\text{Ca}^{2+}$  transients, the reverse mode is predominant (Sah et al., 2003).  $\text{Ca}^{2+}$  influx via the reverse mode of NCX has been suggested to act synergistically with  $\text{Ca}^{2+}$  influx via L-type  $\text{Ca}^{2+}$  channels to trigger  $\text{Ca}^{2+}$  release from the SR as  $I_{\text{Ca,L}}$  are small at depolarized membrane potential (Sah et al., 2003). In addition, NCX can positively regulate SR  $\text{Ca}^{2+}$  load via the reverse mode action (Hirota et al., 2007). Interestingly, under pathophysiological conditions such as cardiac failure, NCX also operates in the reverse mode to allow additional  $\text{Ca}^{2+}$  influx for contraction in order to compensate for the reduction in  $\text{Ca}^{2+}$  release from SR (Gaughan et al., 1999).

NCX is found to be essential for embryo development. Several studies showed that NCX1 is expressed restrictedly in the embryonic heart during early development. NCX knockout are embryonic lethal at ~9-11 days post coitum with immature heart development (Cho et al., 2000; Koushik et al., 2001; Reuter et al., 2003; Wakimoto et al., 2000). Molecular studies demonstrated that NCX mRNA expresses before the appearance of spontaneously beating mESC-CMs; the expression persists thereafter in the CMs (Fu et al., 2006b). It is suggested that expression of NCX at that early stage is essential for early EC-coupling as SR is not well-developed at that stage, NCX is hence essential for maintaining the proper  $\text{Ca}^{2+}$  homeostasis even in the very early stage cardiac development (Reppel et al., 2007a; Reppel et al., 2007b).

Two approaches were used to demonstrate the functional expression of NCX in ESC-CM. In Otsu *et al.* (Otsu et al., 2005), function of NCX was indirectly assessed by using high concentration of NCX blocker KB-R7943. Application of KB-R7943 induced sustained elevation of  $[\text{Ca}^{2+}]_i$ , and this elevation increased as differentiation of mESC-CMs proceeded (Otsu et al., 2005). Apart from using pharmacological blocker, direct measurement of NCX activity was performed by using patch-clamping.  $I_{\text{NCX}}$  was found to be increased as hESC-CMs developed from day 7+40 to day 7+90 (Fu et al., 2010). However, CMs derived from murine embryonic heart at late stage (E16.5) showed a significantly lower  $I_{\text{NCX}}$  density when compared to CMs at early stage (E10.5) (Reppel et al., 2007a), probably due to the high phosphorylation status of NCX in early stage. Consistently, upon differentiation, the proportion of  $\text{Ca}^{2+}$  extrusion by NCX declined from day 9 to day 17 in mESC-CMs (Kapur and Banach, 2007). The discrepancy between different studies on the absolute functional expression of NCX as development proceeds is unknown. Nonetheless, it is clear that NCX is important for  $\text{Ca}^{2+}$  extrusion in differentiating CMs and the decreased contribution by

NCX to  $\text{Ca}^{2+}$  extrusion as development proceeds may be explained by the gradual development of SERCA on the SR.

In the study by Fu *et al.*, the basal  $[\text{Ca}^{2+}]_i$  of mESC-CMs in both early and late developmental stages was increased after applying  $\text{Na}^+$ -free solution, suggesting that NCX is functional in maintaining  $\text{Ca}^{2+}$  homeostasis (Fu *et al.*, 2006b). Interestingly,  $\text{Na}^+$ -free solution completely blocked the  $\text{Ca}^{2+}$  transients in CMs from late developmental stage but not the CMs from early developmental stage, suggesting that NCX starts to regulate  $\text{Ca}^{2+}$  transients only in the late developmental stage (Fu *et al.*, 2006b). Similar results were obtained from hESC-CMs (Fu *et al.*, 2010). Basal  $[\text{Ca}^{2+}]_i$  was marginally increased in CMs at day 7+90 after applying  $\text{Na}^+$ -free solution, but the same was not observed in CMs at day 7+40. In addition, irregular  $\text{Ca}^{2+}$  transient pattern was observed in day 7+90 CMs treated with  $\text{Na}^+$ -free solution (Fu *et al.*, 2010). These suggested that the contribution of NCX to  $\text{Ca}^{2+}$  transients becomes more important as development proceeds in ESC-CMs.

### 2.3.2 Sarcoplasmic/endoplasmic reticulum $\text{Ca}^{2+}$ -ATPase (SERCA)

Details of the structure and function relationship of SERCA have been extensively reviewed (Periasamy *et al.*, 2008; Periasamy and Huke, 2001; Toyoshima, 2008; Toyoshima and Inesi, 2004; Wuytack *et al.*, 2002). SERCA is a single polypeptide with ~1000 amino acid residues located on the ER/SR membrane. It consists of 10 transmembrane (M) domains and 3 cytosolic domains, including actuator (A) domain, phosphorylation (P) domain, and nucleotide-binding (N) domain. A domain regulates the  $\text{Ca}^{2+}$  binding and release. N domain is connected to the P domain; it contains the adenosine binding site and forms the catalytic site. On the other hand, the  $\gamma$ -phosphate reacts with an amino acid residue in the P domain. SERCA utilizes the energy derived from ATP hydrolysis to pump  $\text{Ca}^{2+}$  against concentration gradient from the cytosol to the lumen of ER/SR. Two  $\text{Ca}^{2+}$  are transported by hydrolysis of one ATP in each catalytic cycle.

Regulation of SERCA is mainly achieved by the action of SR membrane proteins phospholamban and sarcolipin (Asahi *et al.*, 2003a; Edes and Kranias, 1987; MacLennan *et al.*, 2002; MacLennan and Kranias, 2003; Simmerman and Jones, 1998; Traaseth *et al.*, 2008). De-phosphorylated form of phospholamban interacts with SERCA and inhibits the pumping activities by decreasing the  $\text{Ca}^{2+}$  affinity of SERCA. Phospholamban exists in monomeric or pentameric form, while the monomeric form is inhibitory. Phosphorylation of phospholamban favors the formation of pentameric form, which in turns relieves the inhibitory effect on SERCA. Phosphorylation of phospholamban is regulated by cAMP-dependent protein kinase (Schwinger *et al.*, 1998) and  $\text{Ca}^{2+}$ /calmodulin-dependent kinase (Ji *et al.*, 2003). Sarcolipin is a shorter homolog of phospholamban (Hellstern *et al.*, 2001). Unlike phospholamban, sarcolipin has no obvious phosphorylation site (Odermatt *et al.*, 1997); therefore, the effect of sarcolipin on SERCA inhibition is mainly controlled by altering the expression level of sarcolipin (Odermatt *et al.*, 1998). Sarcolipin interacts directly with SERCA and inhibits its function by decreasing the  $\text{Ca}^{2+}$  affinity of SERCA. Sarcolipin can also exert its superinhibitory effect on SERCA by forming the tertiary complex phospholamban-sarcolipin-SERCA (Asahi *et al.*, 2002; Asahi *et al.*, 2003b). In addition, sarcolipin stabilizes the SERCA-phospholamban complex in the absence of phospholamban phosphorylation and also inhibits phospholamban phosphorylation (Asahi *et al.*, 2004).

SERCA plays a vital role in  $\text{Ca}^{2+}$  cycling between SR and cytosol, and this is important for EC-coupling. Over-expression of SERCA2a improved cardiac contractility by increasing SR  $\text{Ca}^{2+}$  loading and frequency of  $\text{Ca}^{2+}$  transients (Baker *et al.*, 1998; He *et al.*, 1997; Maier *et al.*,

2005; Prasad et al., 2004). On the other hand, homozygous SERCA2a knockout mice are embryonic lethal (Periasamy et al., 1999), while heterozygous knockout mice are alive and able to reproduce (Shull et al., 2003). Ji *et al.* has reported that the content of SR Ca<sup>2+</sup> stores and the amplitude of Ca<sup>2+</sup> transients were decreased by 40-60% and ~30-40%, respectively, in heterozygous CMs (Ji et al., 2000). Interestingly, heterozygous CMs showed a reduced phospholamban expression, an enhanced phospholamban phosphorylation, and an upregulated NCX expression. However, these changes in Ca<sup>2+</sup> handling proteins were not sufficient to compensate the effects on contractility by the loss of SERCA2a, indicating that SERCA2a is a critical regulator in controlling the E-C coupling of CMs (Ji et al., 2000). Apart from genetic manipulation of the expression level of SERCA, function of SERCA can also be assessed by using the pharmacological blocker thapsigargin. Acute application of thapsigargin caused the decrease in Ca<sup>2+</sup> transient amplitude, rate of decay of Ca<sup>2+</sup> transients, and duration of action potential in isolated ventricular myocytes (Kirby et al., 1992), again suggesting the involvement of SERCA in the E-C coupling of CMs.

In vertebrates, SERCA is encoded by three genes, including the SERCA1, SERCA2, and SERCA3. Alternative splicing of the transcripts from these genes produces more than 10 SERCA isoforms. In CMs, SERCA2a and SERCA2b are expressed, with SERCA2a being the predominant form (Periasamy and Kalyanasundaram, 2007).

SERCA2a mRNA is present before initial contraction of mESC-CMs, but has no obvious change in expression level during further differentiation (Fu et al., 2006b). From embryo studies, SERCA2 protein increased from E9.5 to E18 in mouse heart (Liu et al., 2002). In human, it was reported that SERCA2a protein level remained steady between 8 to 15th week gestation, and started to increase afterwards (Qu and Boutjdir, 2001). Role of SERCA in regulating Ca<sup>2+</sup> transients in ESC-CMs has been studied by several groups. Zhu *et al.* demonstrated that SERCA inhibitors, including thapsigargin and cyclopiazonic acid, reduced ~70% amplitude of Ca<sup>2+</sup> transients in hESC-CMs, but had no effect on the time of decay (Zhu et al., 2009). In case of mESC-CMs, thapsigargin reduced both the amplitude and decay of Ca<sup>2+</sup> transients, but exerted similar inhibitory effect on CMs from the 3 developmental stages (Fu et al., 2006a; Fu et al., 2006b). These findings are therefore consistent with the mRNA expression level of SERCA during mESC-CM differentiation. However, the contribution of Ca<sup>2+</sup> removal by SERCA is estimated to be more important as differentiation proceeds based on Ca<sup>2+</sup> imaging experiments (Kapur and Banach, 2007). Therefore, the role of other SR Ca<sup>2+</sup> handling proteins cannot be neglected. For example, expression level of calsequestrin increases as SR matures (Fu et al., 2006a). This increases the capacity of SR Ca<sup>2+</sup> load and may account for the requirement of greater contribution of Ca<sup>2+</sup> removal by SERCA in later developmental stage of mESC-CMs.

To summarize for the whole chapter, Figure 1 represents a summary of the relative contributions of different proteins responsible for regulating Ca<sup>2+</sup> transients in early differentiating ESC-CMs as development proceeds.

### 3. Conclusion

In summary, early differentiating ESC-CMs have already developed a scheme for regulating their [Ca<sup>2+</sup>]<sub>i</sub> for E-C coupling. The relative contributions of the proteins that regulate Ca<sup>2+</sup> transients alter upon the maturation of CMs. By comparing the regulation of [Ca<sup>2+</sup>]<sub>i</sub> in ESC-CMs and that in adult CMs, we can obtain important insights into the potential strategies for 'fine-tuning' ESC-CMs to better-suit different therapeutic and research purposes.






Fig. 1. Summary of the regulation of  $\text{Ca}^{2+}$  transients in CMs at early differentiation stage (EDS) and late differentiation stage (LDS). Number of a particular protein represents the relative changes in the expression level of that particular protein as differentiation proceeds. Thickness of the arrows represents the relative contribution of a particular path to the  $\text{Ca}^{2+}$  transients as differentiation proceeds.

#### 4. Acknowledgements

When writing up this book chapter, due to space limitations, we were unable to include all the important studies in the field related to calcium homeostasis in ESC-CMs; we apologize to those investigators whom we may have missed here. I. C. Lo and C. K. Wong contributed equally to this work. I. C. Lo and C. K. Wong were supported by postgraduate studentships from the Chinese University of Hong Kong (CUHK). This work was supported by the Competitive Earmarked Research Grant (474907) from the University Grants Committee of the Hong Kong Special Administrative Region and the Focused Investment Scheme (1901073) from the CUHK.

#### 5. References

- Acosta L, Haase H, Morano I, Moorman AF, Franco D. 2004. Regional expression of L-type calcium channel subunits during cardiac development. *Dev Dyn* 230(1):131-136.
- Ai X, Curran JW, Shannon TR, Bers DM, Pogwizd SM. 2005. Ca<sup>2+</sup>/calmodulin-dependent protein kinase modulates cardiac ryanodine receptor phosphorylation and sarcoplasmic reticulum Ca<sup>2+</sup> leak in heart failure. *Circ Res* 97(12):1314-1322.
- Arikkath J, Campbell KP. 2003. Auxiliary subunits: essential components of the voltage-gated calcium channel complex. *Curr Opin Neurobiol* 13(3):298-307.
- Asahi M, Kurzydowski K, Tada M, MacLennan DH. 2002. Sarcolipin inhibits polymerization of phospholamban to induce superinhibition of sarco(endo)plasmic reticulum Ca<sup>2+</sup>-ATPases (SERCAs). *J Biol Chem* 277(30):26725-26728.
- Asahi M, Nakayama H, Tada M, Otsu K. 2003a. Regulation of sarco(endo)plasmic reticulum Ca<sup>2+</sup> adenosine triphosphatase by phospholamban and sarcolipin: implication for cardiac hypertrophy and failure. *Trends Cardiovasc Med* 13(4):152-157.
- Asahi M, Otsu K, Nakayama H, Hikoso S, Takeda T, Gramolini AO, Trivieri MG, Oudit GY, Morita T, Kusakari Y, Hirano S, Hongo K, Hirotani S, Yamaguchi O, Peterson A, Backx PH, Kurihara S, Hori M, MacLennan DH. 2004. Cardiac-specific overexpression of sarcolipin inhibits sarco(endo)plasmic reticulum Ca<sup>2+</sup> ATPase (SERCA2a) activity and impairs cardiac function in mice. *Proc Natl Acad Sci U S A* 101(25):9199-9204.
- Asahi M, Sugita Y, Kurzydowski K, De Leon S, Tada M, Toyoshima C, MacLennan DH. 2003b. Sarcolipin regulates sarco(endo)plasmic reticulum Ca<sup>2+</sup>-ATPase (SERCA) by binding to transmembrane helices alone or in association with phospholamban. *Proc Natl Acad Sci U S A* 100(9):5040-5045.
- Baker DL, Hashimoto K, Grupp IL, Ji Y, Reed T, Loukianov E, Grupp G, Bhagwhat A, Hoit B, Walsh R, Marban E, Periasamy M. 1998. Targeted overexpression of the sarcoplasmic reticulum Ca<sup>2+</sup>-ATPase increases cardiac contractility in transgenic mouse hearts. *Circ Res* 83(12):1205-1214.
- Bare DJ, Kettlun CS, Liang M, Bers DM, Mignery GA. 2005. Cardiac type 2 inositol 1,4,5-trisphosphate receptor: interaction and modulation by calcium/calmodulin-dependent protein kinase II. *J Biol Chem* 280(16):15912-15920.
- Bers DM. 2002. Cardiac excitation-contraction coupling. *Nature* 415(6868):198-205.
- Bers DM. 2004. Macromolecular complexes regulating cardiac ryanodine receptor function. *J Mol Cell Cardiol* 37(2):417-429.

- Boheler KR, Czyz J, Tweedie D, Yang HT, Anisimov SV, Wobus AM. 2002. Differentiation of pluripotent embryonic stem cells into cardiomyocytes. *Circ Res* 91(3):189-201.
- Bohn G, Moosmang S, Conrad H, Ludwig A, Hofmann F, Klugbauer N. 2000. Expression of T- and L-type calcium channel mRNA in murine sinoatrial node. *FEBS Lett* 481(1):73-76.
- Catterall WA. 1995. Structure and function of voltage-gated ion channels. *Annu Rev Biochem* 64:493-531.
- Catterall WA. 2000. Structure and regulation of voltage-gated Ca<sup>2+</sup> channels. *Annu Rev Cell Dev Biol* 16:521-555.
- Catterall WA, Perez-Reyes E, Snutch TP, Striessnig J. 2005. International Union of Pharmacology. XLVIII. Nomenclature and structure-function relationships of voltage-gated calcium channels. *Pharmacol Rev* 57(4):411-425.
- Chen CC, Lamping KG, Nuno DW, Barresi R, Prouty SJ, Lavoie JL, Cribbs LL, England SK, Sigmund CD, Weiss RM, Williamson RA, Hill JA, Campbell KP. 2003. Abnormal coronary function in mice deficient in alpha1H T-type Ca<sup>2+</sup> channels. *Science* 302(5649):1416-1418.
- Cho CH, Kim SS, Jeong MJ, Lee CO, Shin HS. 2000. The Na<sup>+</sup>-Ca<sup>2+</sup> exchanger is essential for embryonic heart development in mice. *Mol Cells* 10(6):712-722.
- Cribbs LL, Martin BL, Schroder EA, Keller BB, Delisle BP, Satin J. 2001. Identification of the t-type calcium channel (Ca(v)3.1d) in developing mouse heart. *Circ Res* 88(4):403-407.
- De Waard M, Gurnett CA, Campbell KP. 1996. Structural and functional diversity of voltage-activated calcium channels. *Ion Channels* 4:41-87.
- DiPolo R. 1979. Calcium influx in internally dialyzed squid giant axons. *J Gen Physiol* 73(1):91-113.
- Dolphin AC. 2006. A short history of voltage-gated calcium channels. *Br J Pharmacol* 147 Suppl 1:S56-62.
- Dolphin AC, Wyatt CN, Richards J, Beattie RE, Craig P, Lee JH, Cribbs LL, Volsen SG, Perez-Reyes E. 1999. The effect of alpha2-delta and other accessory subunits on expression and properties of the calcium channel alpha1G. *J Physiol* 519 Pt 1:35-45.
- Dubel SJ, Altier C, Chaumont S, Lory P, Bourinet E, Nargeot J. 2004. Plasma membrane expression of T-type calcium channel alpha(1) subunits is modulated by high voltage-activated auxiliary subunits. *J Biol Chem* 279(28):29263-29269.
- Edes I, Kranias EG. 1987. Regulation of cardiac sarcoplasmic reticulum function by phospholamban. *Membr Biochem* 7(3):175-192.
- Foskett JK, White C, Cheung KH, Mak DO. 2007. Inositol trisphosphate receptor Ca<sup>2+</sup> release channels. *Physiol Rev* 87(2):593-658.
- Freise D, Himmerkus N, Schroth G, Trost C, Weissgerber P, Freichel M, Flockerzi V. 1999. Mutations of calcium channel beta subunit genes in mice. *Biol Chem* 380(7-8):897-902.
- Fu JD, Jiang P, Rushing S, Liu J, Chiamvimonvat N, Li RA. 2010. Na<sup>+</sup>/Ca<sup>2+</sup> exchanger is a determinant of excitation-contraction coupling in human embryonic stem cell-derived ventricular cardiomyocytes. *Stem Cells Dev* 19(6):773-782.
- Fu JD, Li J, Tweedie D, Yu HM, Chen L, Wang R, Riordon DR, Brugh SA, Wang SQ, Boheler KR, Yang HT. 2006a. Crucial role of the sarcoplasmic reticulum in the

- developmental regulation of Ca<sup>2+</sup> transients and contraction in cardiomyocytes derived from embryonic stem cells. *FASEB J* 20(1):181-183.
- Fu JD, Yu HM, Wang R, Liang J, Yang HT. 2006b. Developmental regulation of intracellular calcium transients during cardiomyocyte differentiation of mouse embryonic stem cells. *Acta Pharmacol Sin* 27(7):901-910.
- Gaughan JP, Furukawa S, Jeevanandam V, Hefner CA, Kubo H, Margulies KB, McGowan BS, Mattiello JA, DiPaola K, Piacentino V, 3rd, Li S, Houser SR. 1999. Sodium/calcium exchange contributes to contraction and relaxation in failed human ventricular myocytes. *Am J Physiol* 277(2 Pt 2):H714-724.
- Giladi M, Boyman L, Mikhasenko H, Hiller R, Khananshvil D. 2010. Essential role of the CBD1-CBD2 linker in slow dissociation of Ca<sup>2+</sup> from the regulatory two-domain tandem of NCX1. *J Biol Chem* 285(36):28117-28125.
- Go LO, Moschella MC, Watras J, Handa KK, Fyfe BS, Marks AR. 1995. Differential regulation of two types of intracellular calcium release channels during end-stage heart failure. *J Clin Invest* 95(2):888-894.
- Green PJ, Warre R, Hayes PD, McNaughton NC, Medhurst AD, Pangalos M, Duckworth DM, Randall AD. 2001. Kinetic modification of the alpha(1I) subunit-mediated T-type Ca(2+) channel by a human neuronal Ca(2+) channel gamma subunit. *J Physiol* 533(Pt 2):467-478.
- He H, Giordano FJ, Hilal-Dandan R, Choi DJ, Rockman HA, McDonough PM, Bluhm WF, Meyer M, Sayen MR, Swanson E, Dillmann WH. 1997. Overexpression of the rat sarcoplasmic reticulum Ca<sup>2+</sup> ATPase gene in the heart of transgenic mice accelerates calcium transients and cardiac relaxation. *J Clin Invest* 100(2):380-389.
- He JQ, Ma Y, Lee Y, Thomson JA, Kamp TJ. 2003. Human embryonic stem cells develop into multiple types of cardiac myocytes: action potential characterization. *Circ Res* 93(1):32-39.
- Hellstern S, Pegoraro S, Karim CB, Lustig A, Thomas DD, Moroder L, Engel J. 2001. Sarcolipin, the shorter homologue of phospholamban, forms oligomeric structures in detergent micelles and in liposomes. *J Biol Chem* 276(33):30845-30852.
- Hess P, Lansman JB, Tsien RW. 1984. Different modes of Ca channel gating behaviour favoured by dihydropyridine Ca agonists and antagonists. *Nature* 311(5986):538-544.
- Hilgemann DW. 1990. Regulation and deregulation of cardiac Na(+)-Ca<sup>2+</sup> exchange in giant excised sarcolemmal membrane patches. *Nature* 344(6263):242-245.
- Hilgemann DW, Matsuoka S, Nagel GA, Collins A. 1992. Steady-state and dynamic properties of cardiac sodium-calcium exchange. Sodium-dependent inactivation. *J Gen Physiol* 100(6):905-932.
- Hirota S, Pertens E, Janssen LJ. 2007. The reverse mode of the Na(+)/Ca(2+) exchanger provides a source of Ca(2+) for store refilling following agonist-induced Ca(2+) mobilization. *Am J Physiol Lung Cell Mol Physiol* 292(2):L438-447.
- Hobom M, Dai S, Marais E, Lacinova L, Hofmann F, Klugbauer N. 2000. Neuronal distribution and functional characterization of the calcium channel alpha2delta-2 subunit. *Eur J Neurosci* 12(4):1217-1226.
- Hosey MM, Chien AJ, Puri TS. 1996. Structure and regulation of L-type calcium channels a current assessment of the properties and roles of channel subunits. *Trends Cardiovasc Med* 6(8):265-273.



- Iino M. 1990. Biphasic  $\text{Ca}^{2+}$  dependence of inositol 1,4,5-trisphosphate-induced Ca release in smooth muscle cells of the guinea pig taenia caeci. *J Gen Physiol* 95(6):1103-1122.
- Iwai M, Michikawa T, Bosanac I, Ikura M, Mikoshiba K. 2007. Molecular basis of the isoform-specific ligand-binding affinity of inositol 1,4,5-trisphosphate receptors. *J Biol Chem* 282(17):12755-12764.
- Iwamoto T, Pan Y, Nakamura TY, Wakabayashi S, Shigekawa M. 1998. Protein kinase C-dependent regulation of  $\text{Na}^{+}/\text{Ca}^{2+}$  exchanger isoforms NCX1 and NCX3 does not require their direct phosphorylation. *Biochemistry* 37(49):17230-17238.
- Iwamoto T, Pan Y, Wakabayashi S, Imagawa T, Yamanaka HI, Shigekawa M. 1996. Phosphorylation-dependent regulation of cardiac  $\text{Na}^{+}/\text{Ca}^{2+}$  exchanger via protein kinase C. *J Biol Chem* 271(23):13609-13615.
- Ji Y, Lalli MJ, Babu GJ, Xu Y, Kirkpatrick DL, Liu LH, Chiamvimonvat N, Walsh RA, Shull GE, Periasamy M. 2000. Disruption of a single copy of the SERCA2 gene results in altered  $\text{Ca}^{2+}$  homeostasis and cardiomyocyte function. *J Biol Chem* 275(48):38073-38080.
- Ji Y, Li B, Reed TD, Lorenz JN, Kaetzel MA, Dedman JR. 2003. Targeted inhibition of  $\text{Ca}^{2+}$ /calmodulin-dependent protein kinase II in cardiac longitudinal sarcoplasmic reticulum results in decreased phospholamban phosphorylation at threonine 17. *J Biol Chem* 278(27):25063-25071.
- Kapur N, Banach K. 2007. Inositol-1,4,5-trisphosphate-mediated spontaneous activity in mouse embryonic stem cell-derived cardiomyocytes. *J Physiol* 581(Pt 3):1113-1127.
- Kehat I, Gepstein A, Spira A, Itskovitz-Eldor J, Gepstein L. 2002. High-resolution electrophysiological assessment of human embryonic stem cell-derived cardiomyocytes: a novel in vitro model for the study of conduction. *Circ Res* 91(8):659-661.
- Kehat I, Kenyagin-Karsenti D, Snir M, Segev H, Amit M, Gepstein A, Livne E, Binah O, Itskovitz-Eldor J, Gepstein L. 2001. Human embryonic stem cells can differentiate into myocytes with structural and functional properties of cardiomyocytes. *J Clin Invest* 108(3):407-414.
- Kirby MS, Sagara Y, Gaa S, Inesi G, Lederer WJ, Rogers TB. 1992. Thapsigargin inhibits contraction and  $\text{Ca}^{2+}$  transient in cardiac cells by specific inhibition of the sarcoplasmic reticulum  $\text{Ca}^{2+}$  pump. *J Biol Chem* 267(18):12545-12551.
- Kolossov E, Fleischmann BK, Liu Q, Bloch W, Viatchenko-Karpinski S, Manzke O, Ji GJ, Bohlen H, Addicks K, Hescheler J. 1998. Functional characteristics of ES cell-derived cardiac precursor cells identified by tissue-specific expression of the green fluorescent protein. *J Cell Biol* 143(7):2045-2056.
- Koushik SV, Wang J, Rogers R, Moskophidis D, Lambert NA, Creazzo TL, Conway SJ. 2001. Targeted inactivation of the sodium-calcium exchanger (Ncx1) results in the lack of a heartbeat and abnormal myofibrillar organization. *FASEB J* 15(7):1209-1211.
- Lacinova L, Klugbauer N. 2004. Modulation of gating currents of the  $\text{Ca}(v)3.1$  calcium channel by alpha 2 delta 2 and gamma 5 subunits. *Arch Biochem Biophys* 425(2):207-213.
- Lanner JT, Georgiou DK, Joshi AD, Hamilton SL. 2010. Ryanodine receptors: structure, expression, molecular details, and function in calcium release. *Cold Spring Harb Perspect Biol* 2(11):a003996.

- Lee SL, Yu AS, Lytton J. 1994. Tissue-specific expression of Na(+)-Ca<sup>2+</sup> exchanger isoforms. *J Biol Chem* 269(21):14849-14852.
- Li X, Zima AV, Sheikh F, Blatter LA, Chen J. 2005. Endothelin-1-induced arrhythmogenic Ca<sup>2+</sup> signaling is abolished in atrial myocytes of inositol-1,4,5-trisphosphate(IP<sub>3</sub>)-receptor type 2-deficient mice. *Circ Res* 96(12):1274-1281.
- Liu W, Yasui K, Opthof T, Ishiki R, Lee JK, Kamiya K, Yokota M, Kodama I. 2002. Developmental changes of Ca(2+) handling in mouse ventricular cells from early embryo to adulthood. *Life Sci* 71(11):1279-1292.
- Ma J, Hayek SM, Bhat MB. 2004. Membrane topology and membrane retention of the ryanodine receptor calcium release channel. *Cell Biochem Biophys* 40(2):207-224.
- MacLennan DH, Abu-Abed M, Kang C. 2002. Structure-function relationships in Ca(2+) cycling proteins. *J Mol Cell Cardiol* 34(8):897-918.
- MacLennan DH, Kranias EG. 2003. Phospholamban: a crucial regulator of cardiac contractility. *Nat Rev Mol Cell Biol* 4(7):566-577.
- Maier LS, Wahl-Schott C, Horn W, Weichert S, Pagel C, Wagner S, Dybkova N, Muller OJ, Nabauer M, Franz WM, Pieske B. 2005. Increased SR Ca<sup>2+</sup> cycling contributes to improved contractile performance in SERCA2a-overexpressing transgenic rats. *Cardiovasc Res* 67(4):636-646.
- Mangoni ME, Traboulsie A, Leoni AL, Couette B, Marger L, Le Quang K, Kupfer E, Cohen-Solal A, Vilar J, Shin HS, Escande D, Charpentier F, Nargeot J, Lory P. 2006. Bradycardia and slowing of the atrioventricular conduction in mice lacking Cav3.1/alpha1G T-type calcium channels. *Circ Res* 98(11):1422-1430.
- Matsuoka S, Nicoll DA, He Z, Philipson KD. 1997. Regulation of cardiac Na(+)-Ca<sup>2+</sup> exchanger by the endogenous XIP region. *J Gen Physiol* 109(2):273-286.
- Mery A, Aimond F, Menard C, Mikoshiba K, Michalak M, Puceat M. 2005. Initiation of embryonic cardiac pacemaker activity by inositol 1,4,5-trisphosphate-dependent calcium signaling. *Mol Biol Cell* 16(5):2414-2423.
- Mizuta E, Miake J, Yano S, Furuichi H, Manabe K, Sasaki N, Igawa O, Hoshikawa Y, Shigemasa C, Nanba E, Ninomiya H, Hidaka K, Morisaki T, Tajima F, Hisatome I. 2005. Subtype switching of T-type Ca<sup>2+</sup> channels from Cav3.2 to Cav3.1 during differentiation of embryonic stem cells to cardiac cell lineage. *Circ J* 69(10):1284-1289.
- Moore JC, Tsang SY, Rushing SN, Lin D, Tse HF, Chan CW, Li RA. 2008. Functional consequences of overexpressing the gap junction Cx43 in the cardiogenic potential of pluripotent human embryonic stem cells. *Biochem Biophys Res Commun* 377(1):46-51.
- Moreno Davila H. 1999. Molecular and functional diversity of voltage-gated calcium channels. *Ann N Y Acad Sci* 868:102-117.
- Moschella MC, Marks AR. 1993. Inositol 1,4,5-trisphosphate receptor expression in cardiac myocytes. *J Cell Biol* 120(5):1137-1146.
- Mummery C, Ward-van Oostwaard D, Doevendans P, Spijker R, van den Brink S, Hassink R, van der Heyden M, Opthof T, Pera M, de la Riviere AB, Passier R, Tertoolen L. 2003. Differentiation of human embryonic stem cells to cardiomyocytes: role of coculture with visceral endoderm-like cells. *Circulation* 107(21):2733-2740.

- Nakayama H, Bodi I, Maillet M, DeSantiago J, Domeier TL, Mikoshiba K, Lorenz JN, Blatter LA, Bers DM, Molkentin JD. 2010. The IP<sub>3</sub> receptor regulates cardiac hypertrophy in response to select stimuli. *Circ Res* 107(5):659-666.
- Nakazawa M, Uchida K, Aramaki M, Kodo K, Yamagishi C, Takahashi T, Mikoshiba K, Yamagishi H. 2011. Inositol 1,4,5-trisphosphate receptors are essential for the development of the second heart field. *J Mol Cell Cardiol*.
- Ng SY, Wong CK, Tsang SY. 2010. Differential gene expressions in atrial and ventricular myocytes: insights into the road of applying embryonic stem cell-derived cardiomyocytes for future therapies. *Am J Physiol Cell Physiol* 299(6):C1234-1249.
- Nicoll DA, Hryshko LV, Matsuoka S, Frank JS, Philipson KD. 1996. Mutation of amino acid residues in the putative transmembrane segments of the cardiac sarcolemmal Na<sup>+</sup>-Ca<sup>2+</sup> exchanger. *J Biol Chem* 271(23):13385-13391.
- Odermatt A, Becker S, Khanna VK, Kurzydowski K, Leisner E, Pette D, MacLennan DH. 1998. Sarcolipin regulates the activity of SERCA1, the fast-twitch skeletal muscle sarcoplasmic reticulum Ca<sup>2+</sup>-ATPase. *J Biol Chem* 273(20):12360-12369.
- Odermatt A, Taschner PE, Scherer SW, Beatty B, Khanna VK, Cornblath DR, Chaudhry V, Yee WC, Schrank B, Karpati G, Breuning MH, Knoers N, MacLennan DH. 1997. Characterization of the gene encoding human sarcolipin (SLN), a proteolipid associated with SERCA1: absence of structural mutations in five patients with Brody disease. *Genomics* 45(3):541-553.
- Ono K, Iijima T. 2005. Pathophysiological significance of T-type Ca<sup>2+</sup> channels: properties and functional roles of T-type Ca<sup>2+</sup> channels in cardiac pacemaking. *J Pharmacol Sci* 99(3):197-204.
- Otsu K, Kuruma A, Yanagida E, Shoji S, Inoue T, Hirayama Y, Uematsu H, Hara Y, Kawano S. 2005. Na<sup>+</sup>/K<sup>+</sup> ATPase and its functional coupling with Na<sup>+</sup>/Ca<sup>2+</sup> exchanger in mouse embryonic stem cells during differentiation into cardiomyocytes. *Cell Calcium* 37(2):137-151.
- Ottolia M, Philipson KD, John S. 2004. Conformational changes of the Ca(2+) regulatory site of the Na(+)-Ca(2+) exchanger detected by FRET. *Biophys J* 87(2):899-906.
- Perez-Reyes E. 2003. Molecular physiology of low-voltage-activated t-type calcium channels. *Physiol Rev* 83(1):117-161.
- Perez-Reyes E. 2006. Molecular characterization of T-type calcium channels. *Cell Calcium* 40(2):89-96.
- Periasamy M, Bhupathy P, Babu GJ. 2008. Regulation of sarcoplasmic reticulum Ca<sup>2+</sup> ATPase pump expression and its relevance to cardiac muscle physiology and pathology. *Cardiovasc Res* 77(2):265-273.
- Periasamy M, Huke S. 2001. SERCA pump level is a critical determinant of Ca(2+)homeostasis and cardiac contractility. *J Mol Cell Cardiol* 33(6):1053-1063.
- Periasamy M, Kalyanasundaram A. 2007. SERCA pump isoforms: their role in calcium transport and disease. *Muscle Nerve* 35(4):430-442.
- Periasamy M, Reed TD, Liu LH, Ji Y, Loukianov E, Paul RJ, Nieman ML, Riddle T, Duffy JJ, Doetschman T, Lorenz JN, Shull GE. 1999. Impaired cardiac performance in heterozygous mice with a null mutation in the sarco(endo)plasmic reticulum Ca<sup>2+</sup>-ATPase isoform 2 (SERCA2) gene. *J Biol Chem* 274(4):2556-2562.
- Philipson KD, Nicoll DA. 2000. Sodium-calcium exchange: a molecular perspective. *Annu Rev Physiol* 62:111-133.

- Platzer J, Engel J, Schrott-Fischer A, Stephan K, Bova S, Chen H, Zheng H, Striessnig J. 2000. Congenital deafness and sinoatrial node dysfunction in mice lacking class D L-type Ca<sup>2+</sup> channels. *Cell* 102(1):89-97.
- Pott C, Philipson KD, Goldhaber JJ. 2005. Excitation-contraction coupling in Na<sup>+</sup>-Ca<sup>2+</sup> exchanger knockout mice: reduced transsarcolemmal Ca<sup>2+</sup> flux. *Circ Res* 97(12):1288-1295.
- Prasad V, Okunade GW, Miller ML, Shull GE. 2004. Phenotypes of SERCA and PMCA knockout mice. *Biochem Biophys Res Commun* 322(4):1192-1203.
- Qu Y, Baroudi G, Yue Y, El-Sherif N, Boutjdir M. 2005. Localization and modulation of  $\alpha$ 1D (Cav1.3) L-type Ca channel by protein kinase A. *Am J Physiol Heart Circ Physiol* 288(5):H2123-2130.
- Qu Y, Boutjdir M. 2001. Gene expression of SERCA2a and L- and T-type Ca channels during human heart development. *Pediatr Res* 50(5):569-574.
- Rapila R, Korhonen T, Tavi P. 2008. Excitation-contraction coupling of the mouse embryonic cardiomyocyte. *J Gen Physiol* 132(4):397-405.
- Reppel M, Fleischmann BK, Reuter H, Sasse P, Schunkert H, Hescheler J. 2007a. Regulation of the Na<sup>+</sup>/Ca<sup>2+</sup> exchanger (NCX) in the murine embryonic heart. *Cardiovasc Res* 75(1):99-108.
- Reppel M, Sasse P, Malan D, Nguemo F, Reuter H, Bloch W, Hescheler J, Fleischmann BK. 2007b. Functional expression of the Na<sup>+</sup>/Ca<sup>2+</sup> exchanger in the embryonic mouse heart. *J Mol Cell Cardiol* 42(1):121-132.
- Reuter H, Henderson SA, Han T, Mottino GA, Frank JS, Ross RS, Goldhaber JJ, Philipson KD. 2003. Cardiac excitation-contraction coupling in the absence of Na<sup>(+)</sup> - Ca<sup>2+</sup> exchange. *Cell Calcium* 34(1):19-26.
- Richards MW, Butcher AJ, Dolphin AC. 2004. Ca<sup>2+</sup> channel beta-subunits: structural insights AID our understanding. *Trends Pharmacol Sci* 25(12):626-632.
- Rosemblit N, Moschella MC, Ondriasa E, Gutstein DE, Ondrias K, Marks AR. 1999. Intracellular calcium release channel expression during embryogenesis. *Dev Biol* 206(2):163-177.
- Sah R, Ramirez RJ, Oudit GY, Gidrewicz D, Trivieri MG, Zobel C, Backx PH. 2003. Regulation of cardiac excitation-contraction coupling by action potential repolarization: role of the transient outward potassium current (I<sub>(to)</sub>). *J Physiol* 546(Pt 1):5-18.
- Satin J, Itzhaki I, Rapoport S, Schroder EA, Izu L, Arbel G, Beyar R, Balke CW, Schiller J, Gepstein L. 2008. Calcium handling in human embryonic stem cell-derived cardiomyocytes. *Stem Cells* 26(8):1961-1972.
- Sauer H, Theben T, Hescheler J, Lindner M, Brandt MC, Wartenberg M. 2001. Characteristics of calcium sparks in cardiomyocytes derived from embryonic stem cells. *Am J Physiol Heart Circ Physiol* 281(1):H411-421.
- Schwinger RH, Bolck B, Munch G, Brixius K, Muller-Ehmsen J, Erdmann E. 1998. cAMP-dependent protein kinase A-stimulated sarcoplasmic reticulum function in heart failure. *Ann N Y Acad Sci* 853:240-250.
- Sedan O, Dolnikov K, Zeevi-Levin N, Fleishmann N, Spiegel I, Berdichevski S, Amit M, Itskovitz-Eldor J, Binah O. 2010. Human embryonic stem cell-derived cardiomyocytes can mobilize 1,4,5-inositol trisphosphate-operated [Ca<sup>2+</sup>]<sub>i</sub> stores:




- the functionality of angiotensin-II/endothelin-1 signaling pathways. *Ann N Y Acad Sci* 1188:68-77.
- Sedan O, Dolnikov K, Zeevi-Levin N, Leibovich N, Amit M, Itskovitz-Eldor J, Binah O. 2008. 1,4,5-Inositol trisphosphate-operated intracellular Ca(2+) stores and angiotensin-II/endothelin-1 signaling pathway are functional in human embryonic stem cell-derived cardiomyocytes. *Stem Cells* 26(12):3130-3138.
- Seisenberger C, Specht V, Welling A, Platzer J, Pfeifer A, Kuhbandner S, Striessnig J, Klugbauer N, Feil R, Hofmann F. 2000. Functional embryonic cardiomyocytes after disruption of the L-type alpha1C (Cav1.2) calcium channel gene in the mouse. *J Biol Chem* 275(50):39193-39199.
- Shull GE, Okunade G, Liu LH, Kozel P, Periasamy M, Lorenz JN, Prasad V. 2003. Physiological functions of plasma membrane and intracellular Ca<sup>2+</sup> pumps revealed by analysis of null mutants. *Ann N Y Acad Sci* 986:453-460.
- Simmerman HK, Jones LR. 1998. Phospholamban: protein structure, mechanism of action, and role in cardiac function. *Physiol Rev* 78(4):921-947.
- Spencer CI, Sham JS. 2003. Effects of Na<sup>+</sup>/Ca<sup>2+</sup> exchange induced by SR Ca<sup>2+</sup> release on action potentials and afterdepolarizations in guinea pig ventricular myocytes. *Am J Physiol Heart Circ Physiol* 285(6):H2552-2562.
- Takeshima H, Komazaki S, Hirose K, Nishi M, Noda T, Iino M. 1998. Embryonic lethality and abnormal cardiac myocytes in mice lacking ryanodine receptor type 2. *EMBO J* 17(12):3309-3316.
- Taylor CW, da Fonseca PC, Morris EP. 2004. IP(3) receptors: the search for structure. *Trends Biochem Sci* 29(4):210-219.
- Taylor CW, Laude AJ. 2002. IP<sub>3</sub> receptors and their regulation by calmodulin and cytosolic Ca<sup>2+</sup>. *Cell Calcium* 32(5-6):321-334.
- Taylor CW, Tovey SC. 2010. IP(3) receptors: toward understanding their activation. *Cold Spring Harb Perspect Biol* 2(12):a004010.
- Thomson JA, Itskovitz-Eldor J, Shapiro SS, Waknitz MA, Swiergiel JJ, Marshall VS, Jones JM. 1998. Embryonic stem cell lines derived from human blastocysts. *Science* 282(5391):1145-1147.
- Toyoshima C. 2008. Structural aspects of ion pumping by Ca<sup>2+</sup>-ATPase of sarcoplasmic reticulum. *Arch Biochem Biophys* 476(1):3-11.
- Toyoshima C, Inesi G. 2004. Structural basis of ion pumping by Ca<sup>2+</sup>-ATPase of the sarcoplasmic reticulum. *Annu Rev Biochem* 73:269-292.
- Traaseth NJ, Ha KN, Verardi R, Shi L, Buffy JJ, Masterson LR, Veglia G. 2008. Structural and dynamic basis of phospholamban and sarcolipin inhibition of Ca(2+)-ATPase. *Biochemistry* 47(1):3-13.
- Tu H, Wang Z, Nosyreva E, De Smedt H, Bezprozvanny I. 2005. Functional characterization of mammalian inositol 1,4,5-trisphosphate receptor isoforms. *Biophys J* 88(2):1046-1055.
- Uchida K, Aramaki M, Nakazawa M, Yamagishi C, Makino S, Fukuda K, Nakamura T, Takahashi T, Mikoshiba K, Yamagishi H. 2010. Gene knock-outs of inositol 1,4,5-trisphosphate receptors types 1 and 2 result in perturbation of cardiogenesis. *PLoS One* 5(9).
- Viatchenko-Karpinski S, Fleischmann BK, Liu Q, Sauer H, Gryshchenko O, Ji GJ, Hescheler J. 1999. Intracellular Ca<sup>2+</sup> oscillations drive spontaneous contractions in

- cardiomyocytes during early development. *Proc Natl Acad Sci U S A* 96(14):8259-8264.
- Wakimoto K, Kobayashi K, Kuro OM, Yao A, Iwamoto T, Yanaka N, Kita S, Nishida A, Azuma S, Toyoda Y, Omori K, Imahie H, Oka T, Kudoh S, Kohmoto O, Yazaki Y, Shigekawa M, Imai Y, Nabeshima Y, Komuro I. 2000. Targeted disruption of Na<sup>+</sup>/Ca<sup>2+</sup> exchanger gene leads to cardiomyocyte apoptosis and defects in heartbeat. *J Biol Chem* 275(47):36991-36998.
- Wang J, Chan TO, Zhang XQ, Gao E, Song J, Koch WJ, Feldman AM, Cheung JY. 2009. Induced overexpression of Na<sup>+</sup>/Ca<sup>2+</sup> exchanger transgene: altered myocyte contractility, [Ca<sup>2+</sup>]<sub>i</sub> transients, SR Ca<sup>2+</sup> contents, and action potential duration. *Am J Physiol Heart Circ Physiol* 297(2):H590-601.
- Wanichawan P, Louch WE, Hortemo KH, Austbo B, Lunde PK, Scott JD, Sejersted OM, Carlson CR. 2011. Full-length cardiac Na<sup>+</sup>/Ca<sup>2+</sup> exchanger 1 protein is not phosphorylated by protein kinase A. *Am J Physiol Cell Physiol* 300(5):C989-997.
- Wu M, Le HD, Wang M, Yurkov V, Omelchenko A, Hnatowich M, Nix J, Hryshko LV, Zheng L. 2010. Crystal structures of progressive Ca<sup>2+</sup> binding states of the Ca<sup>2+</sup> sensor Ca<sup>2+</sup> binding domain 1 (CBD1) from the CALX Na<sup>+</sup>/Ca<sup>2+</sup> exchanger reveal incremental conformational transitions. *J Biol Chem* 285(4):2554-2561.
- Wuytack F, Raeymaekers L, Missiaen L. 2002. Molecular physiology of the SERCA and SPCA pumps. *Cell Calcium* 32(5-6):279-305.
- Xu C, Police S, Rao N, Carpenter MK. 2002. Characterization and enrichment of cardiomyocytes derived from human embryonic stem cells. *Circ Res* 91(6):501-508.
- Xu M, Welling A, Papparisto S, Hofmann F, Klugbauer N. 2003. Enhanced expression of L-type Cav1.3 calcium channels in murine embryonic hearts from Cav1.2-deficient mice. *J Biol Chem* 278(42):40837-40841.
- Xue T, Cho HC, Akar FG, Tsang SY, Jones SP, Marban E, Tomaselli GF, Li RA. 2005. Functional integration of electrically active cardiac derivatives from genetically engineered human embryonic stem cells with quiescent recipient ventricular cardiomyocytes: insights into the development of cell-based pacemakers. *Circulation* 111(1):11-20.
- Yanagi K, Takano M, Narazaki G, Uosaki H, Hoshino T, Ishii T, Misaki T, Yamashita JK. 2007. Hyperpolarization-activated cyclic nucleotide-gated channels and T-type calcium channels confer automaticity of embryonic stem cell-derived cardiomyocytes. *Stem Cells* 25(11):2712-2719.
- Yanagida E, Shoji S, Hirayama Y, Yoshikawa F, Otsu K, Uematsu H, Hiraoka M, Furuichi T, Kawano S. 2004. Functional expression of Ca<sup>2+</sup> signaling pathways in mouse embryonic stem cells. *Cell Calcium* 36(2):135-146.
- Yang HT, Tweedie D, Wang S, Guia A, Vinogradova T, Bogdanov K, Allen PD, Stern MD, Lakatta EG, Boheler KR. 2002. The ryanodine receptor modulates the spontaneous beating rate of cardiomyocytes during development. *Proc Natl Acad Sci U S A* 99(14):9225-9230.
- Yang L, Katchman A, Morrow JP, Doshi D, Marx SO. 2011. Cardiac L-type calcium channel (Cav1.2) associates with gamma subunits. *FASEB J* 25(3):928-936.
- Yasui K, Niwa N, Takemura H, Opthof T, Muto T, Horiba M, Shimizu A, Lee JK, Honjo H, Kamiya K, Kodama I. 2005. Pathophysiological significance of T-type Ca<sup>2+</sup>

- channels: expression of T-type  $\text{Ca}^{2+}$  channels in fetal and diseased heart. *J Pharmacol Sci* 99(3):205-210.
- Yule DI, Betzenhauser MJ, Joseph SK. 2010. Linking structure to function: Recent lessons from inositol 1,4,5-trisphosphate receptor mutagenesis. *Cell Calcium* 47(6):469-479.
- Zhang YM, Shang L, Hartzell C, Narlow M, Cribbs L, Dudley SC, Jr. 2003. Characterization and regulation of T-type  $\text{Ca}^{2+}$  channels in embryonic stem cell-derived cardiomyocytes. *Am J Physiol Heart Circ Physiol* 285(6):H2770-2779.
- Zhang Z, Xu Y, Song H, Rodriguez J, Tuteja D, Namkung Y, Shin HS, Chiamvimonvat N. 2002. Functional Roles of  $\text{Ca}_v1.3$  ( $\alpha_1D$ ) calcium channel in sinoatrial nodes: insight gained using gene-targeted null mutant mice. *Circ Res* 90(9):981-987.
- Zhu WZ, Santana LF, Laflamme MA. 2009. Local control of excitation-contraction coupling in human embryonic stem cell-derived cardiomyocytes. *PLoS One* 4(4):e5407.

IntechOpen



## **Embryonic Stem Cells - Differentiation and Pluripotent Alternatives**

Edited by Prof. Michael S. Kallos

ISBN 978-953-307-632-4

Hard cover, 506 pages

**Publisher** InTech

**Published online** 12, October, 2011

**Published in print edition** October, 2011

The ultimate clinical implementation of embryonic stem cells will require methods and protocols to turn these unspecialized cells into the fully functioning cell types found in a wide variety of tissues and organs. In order to achieve this, it is necessary to clearly understand the signals and cues that direct embryonic stem cell differentiation. This book provides a snapshot of current research on the differentiation of embryonic stem cells to a wide variety of cell types, including neural, cardiac, endothelial, osteogenic, and hepatic cells. In addition, induced pluripotent stem cells and other pluripotent stem cell sources are described. The book will serve as a valuable resource for engineers, scientists, and clinicians as well as students in a wide range of disciplines.

### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Iek Chi Lo, Chun Kit Wong and Suk Ying Tsang (2011). Maintenance Of Calcium Homeostasis in Embryonic Stem Cell-Derived Cardiomyocytes, Embryonic Stem Cells - Differentiation and Pluripotent Alternatives, Prof. Michael S. Kallos (Ed.), ISBN: 978-953-307-632-4, InTech, Available from:

<http://www.intechopen.com/books/embryonic-stem-cells-differentiation-and-pluripotent-alternatives/maintenance-of-calcium-homeostasis-in-embryonic-stem-cell-derived-cardiomyocytes>

**INTECH**  
open science | open minds

### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166  
[www.intechopen.com](http://www.intechopen.com)

### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821



© 2011 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen