

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,400

Open access books available

133,000

International authors and editors

165M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Asthma and Health Related Quality of Life in Childhood and Adolescence

Esther Hafkamp-de Groen and Hein Raat
*Erasmus MC-University Medical Center Rotterdam
The Netherlands*

1. Introduction

Asthma is the most frequent chronic disorder in childhood. Asthma puts a serious burden on children's health related quality of life, despite the availability of effective and safe treatment (Dalheim-Englund et al., 2004; Global Initiative for Asthma, 2010; Masoli et al., 2004; Mohangoo et al., 2005). The overall goal of asthma management is to achieve optimal disease control and health related quality of life improvements (Bateman et al., 2007; Pedersen et al., 2011). The World Health Organization has defined the term health related quality of life as the individual's perception of their position of life in the context of the culture and value systems in which they live and in relation to their goals, expectations and concerns (World Health Organization, 1993). The own perception is important because it emphasises that these are the impairments that patients themselves consider important. As in most medical conditions, the correlation between asthma control and health related quality of life is modest. Therefore, the impact that asthma has on a patient's health related quality of life cannot be inferred from the conventional clinical measures of asthma (e.g. spirometry); it must be measured directly (Juniper et al., 1999a, 1999b).

During the past decade, the use of health related quality of life as an essential outcome measure of childhood asthma treatment and management has increased (Merikallio et al., 2005). This review summarises recent literature on: 1) health related quality of life instruments for childhood asthma, 2) the impact of childhood asthma on children's health related quality of life, 3) the impact of children's asthma on caregiver's health related quality of life and 5) factors associated with health related quality of life in childhood asthma.

2. Health related quality of life instruments and childhood asthma

Several feasible, reliable and validated pediatric health related quality of life questionnaires are standardised and available to measure health related quality of life in asthmatic children (Fiese et al., 2005; Raat et al., 2006). Both generic and asthma-specific questionnaires are used to measure health related quality of life in school aged children. Generic health related quality of life questionnaires intend to measure all dimensions of health-related quality of life (Raat et al., 2006). Frequently applied generic health related quality of life questionnaires are: the Child Health Questionnaire (CHQ) (Gorelick et al., 2003), the Pediatric Quality of Life Inventory (PedsQL) (Varni et al., 2005), the TNO-AZL (Preschool) Children's Quality of Life questionnaire (TAPQoL/TACQoL) (Bunge et al., 2005), the Infant-Toddler Quality of

Life (ITQOL) questionnaire (Spuijbroek et al., 2011) and the KIDSCREEN/DISABKIDS questionnaires (Petersen et al., 2005). Asthma-specific health related quality of life questionnaires focus on those dimensions that are likely to be affected by asthma disease or treatment. The most prominent asthma-specific health related quality of life questionnaires are the Pediatric Asthma Quality of Life Questionnaire (PAQLQ) (Juniper et al., 1996; Raat et al. 2005), the How Are You (HAY) (Le Coq et al., 2000) instrument and the Childhood Asthma Questionnaire (CAQ) (Christie et al., 1993).

If children are unable to report about their own experience reliably, parents are appropriate sources of information about health related quality of life (Petsios et al., 2011). One study suggests that fathers may be better proxy reporters than mothers (Petsios et al., 2011). The correlation between child and parent reported quality of life improves with increasing age of the child (Annett et al., 2003). Although the agreement between child self-report and parent proxy report on health related quality of life has been showed as satisfactory, according to Petsios *et al.* (2011), parents may overestimate health related quality of life of their children with asthma. This has to be taken into account when interpreting results from parent reported health related quality of life questionnaires, in comparison with child self-reports.

The PAQLQ is the most frequently used disease-specific health related quality of life instrument with regard to childhood asthma. Therefore, using this instrument has the benefit for researchers that results can more easily be compared with previous findings. However, using the existing health related quality of life instruments may have some limitations. A recent study has investigated whether asthma-specific health related quality of life questionnaires actually include all relevant aspects of asthma-specific health related quality of life for children with asthma (Annett et al., 2003). They have found disagreement between distinct health related quality of life questionnaires on components of asthma-specific health related quality of life: only some components of the asthma symptoms domain and of the activity limitations domain are part of all questionnaires. Furthermore, according to Van den Bemt *et al.* (2010), not all essential components of asthma-specific health related quality of life, according to childhood asthma, are part of existing asthma-specific health related quality of life questionnaires.

When classifying health related quality of life questionnaires into standardised and individualised health related quality of life instruments, another limitation is revealed. In standardised health related quality of life instruments the questions and range of answers are predetermined and the same for all patients. As opposed to standardised health related quality of life instruments, individualised health related quality of life instruments allow patients to define their quality of life in relation to their goals and expectations. Carr & Higginson (2001) conclude that standardised health related quality of life questionnaires have limited ability to capture the health related quality of life of individual asthma patients.

The most appropriate approach to measure health related quality of life in asthmatic children would be to use a combination of parental and self-reports of both generic and asthma-specific health related quality of life by validated questionnaires (Raat et al., 2006). Whether such health related quality of life measures are truly patient centred and to what extent they actually represent the quality of life of individual or groups of asthmatic children should always be taken into account when one interprets study results (Carr & Higginson, 2001).

3. Impact of asthma on children's health related quality of life

Asthma might have physical, emotional and psychosocial impact on children's lives (Grootenhuis et al., 2007; Juniper, 1997; Merikallio et al., 2005; Sawyer et al., 2004). Important components of health related quality of life are the effects on, and consequences of asthma on peer relationships (e.g., being bullied), the dependence on medication, shortness of breath, cough, limitations in activities and limitations due to the response on cigarette smoke exposure (Van den Bemt et al., 2010). Compared to preschool children without asthma symptoms, preschool children with asthma symptoms have significantly lower health related quality of life scores for lung problems, sleeping, appetite, communication and positive mood health related quality of life scales (Mohangoo et al., 2005).

Most studies have focused on severity of symptoms to examine the impact of asthma symptoms on children's health related quality life; the results are conflicting (Everhart & Fiese, 2009). For example, disease severity is not consistently associated with children's health related quality of life in some studies (Erickson et al., 2002; Vila et al., 2003), whereas others report that children with moderate or severe asthma have a worse level of functioning in several domains of their health related quality of life compared to children with mild asthma (Annett et al., 2001; Merikallio et al., 2005; Mohangoo et al., 2007, 2011; Sawyer et al., 2000) suggesting there may be a 'dose-response' relationship between the frequency and intensity of children's asthma symptoms and their health related quality of life. Mohangoo *et al.* (2007, 2011) evaluated health related quality of life in infants and adolescents with asthma-like symptoms, such as attacks of wheezing and shortness of breath (Mohangoo et al., 2007, 2011). Asthma-like symptoms during the first year of life are associated with impaired health related quality of life at the age of 12 months. Also, the presence of at least four wheezing attacks during the past year was associated with impaired adolescents' health related quality of life. Frequent wheezing attacks mostly affect adolescents' general health, bodily pain, self esteem and mental health (Mohangoo et al., 2007). Previous studies have also found that wheezing attacks more often have a physical impact than a psychosocial impact (Merikallio et al., 2005).

As described earlier, one of the main goals of asthma management is to achieve good asthma control. Asthma control has been defined as the minimisation of night time and daytime symptoms, activity limitation, rescue bronchodilator use and airway narrowing (Global Initiative for Asthma, 2010). Poorly controlled asthma symptoms impair health related quality of life in children (Guilbert et al., 2011). An important issue is whether proper asthma management improves quality of life in asthma patients, and whether poor health related quality of life makes disease management harder. Studies have found that poor health related quality of life is predictive of subsequent asthma-related emergency department visits, which implicates poor asthma control (Magid et al, 2004). Pont *et al.* (2004) show that proper asthma management improves health related quality of life.

In short, children experience asthma as an interruption in daily life that influences them physically, emotionally and socially.

4. Impact of children's asthma on caregiver's health related quality of life

With childhood asthma, the family and particularly the primary caregiver may face a considerable burden. While there are several questionnaires for assessing parental/caregiver's health related quality of life not directly related to asthma (Osman &

Silverman, 1996), there is only one instrument to examine the specific impact of childhood asthma on parental/caregiver functioning: The Pediatric Asthma Caregiver's Quality of Life Questionnaire (PACQLQ) (Juniper et al., 1996).

Whereas some studies find no association between caregiver's health related quality of life and children's asthma symptoms (Annett et al., 2003), duration of asthma illness and asthma pre-treatment severity (Vila et al., 2003), other studies report that caregiver's and child's health related quality of life are significantly associated with each other (Dean et al., 2009, 2010; Garro, 2011; Halterman et al., 2004). Halterman *et al.* (2004) find that higher symptom levels with regard to childhood asthma are associated with lower parental health related quality of life. Further, when children's symptoms improve, parents show higher health related quality of life.

It should be considered how childhood asthma affects caregiver's health related quality of life. Caregivers of asthmatic children appear to be more compromised in their resistance to stress, mood, emotional stability, amount of spare time and leisure activities (Garro, 2011). Caregivers of children with uncontrolled asthma report significantly higher absenteeism than their controlled counterparts (Dean et al., 2009, 2010).

Both caregiver's health related quality of life, caregiver's perception of the child's asthma symptoms, and the child's health related quality of life may be important in diagnosis and control of established asthma in childhood (Skoner, 2002). While giving attention to the caregiver's health related quality of life, it should be taken into account that the profile of health related quality of life impairment is different in asthmatic children and in their parents (Farnik et al., 2010). Where activity limitation seems to be the most impaired domain in children, asthma symptom perception and emotional health appear to be the most affected health related quality of life domains in parents.

In addition to evaluation of the asthmatic child, the integral assessment of asthma requires the evaluation of caregiver's health related quality of life. Giving attention to caregiver's health related quality of life is needed in clinical practice in order to avoid possible interferences of the caregiver's distress in the optimization of child's asthma treatment outcomes (Majani et al., 2005).

5. Factors associated with health related quality of life in asthmatic children

As we described earlier, the frequency and severity of asthma attacks and effects of asthma management or treatment are associated with children's health related quality of life. Researchers have also investigated other variables in association to health related quality of life in childhood asthma (Annett et al., 2003; Erickson et al., 2002; Mrazek, 1992; Sawyer et al., 2000, 2001). Hospital admissions, absences from school, limitations of sport and other activities, sleeping problems (and fatigue) are associated with health related quality of life in asthmatic children (Mrazek, 1992). Erickson *et al.* (2002) show that both asthma morbidity and health related quality of life are related to socioeconomic status. Also, household income is most consistently associated with the health related quality of life of asthmatic children and their caregivers. Sawyer *et al.* (2001) report the impact of family functioning on health related quality of life in children with asthma. They have found that the degree to which children are upset by their asthma is related to general functioning of their families, and their symptom levels are associated with several dimensions of family functioning (Sawyer et al., 2000, 2001). Children living in families with more clearly defined roles, greater interest and concern for the well-being of each other and clearer rules have been

found to be less bothered by their asthma symptoms (Sawyer et al., 2000). A study by Annett *et al.* (2003) didn't find an association between health related quality of life of asthmatic children and family functioning, measured by the degree of cohesion among family members.

Results suggest that several factors may impact health related quality of life of asthmatic children. Important predictors of the health related quality of life of asthmatic children are socioeconomic status and family functioning. These findings implicate the need of specific attention to health related quality of life in asthmatic children from families with low socioeconomic status and poor family functioning.

6. Conclusion

Health care workers should be aware of the impact of asthma on children's life, their families and the factors associated with the health related quality of life of these children. Routine use of an health related quality of life questionnaire to evaluate health related quality of life in children with asthma symptoms and their caregivers should be recommended in health care. Specific application, for example, can be found in preventive child health care and in primary health care to prevent impairment of health related quality of life due to asthma symptoms and to realise adequate management of asthma symptoms. Attention should be given to health related quality of life in asthmatic children from families with low socioeconomic status and poor family functioning. Generally, a combination of parental and self-reports of both general and asthma-specific patient centred health related quality of life questionnaires should be applied. Further research should focus on which factors are responsible for the greatest burden on asthmatic children's health related quality of life and their caregivers' health related quality of life and how such risk factors should be prevented and managed.


7. References

- Annett, RD.; Bender, BG.; Lapidus, J.; DuHamel, TR. & Lincoln, A. (2001). Predicting children's quality of life in an asthma clinical trial: What did children's reports tell us? *J Pediatr*, Vol.139, No.6, pp. 854-861, ISSN 0022-3476
- Annett, RD.; Bender, BG.; DuHamel, TR. & Lapidus, J. (2003). Factors influencing parent reports on quality of life for children with asthma. *J Asthma*, Vol.40, No.5, pp. 577-587, ISSN 0277-0903
- Bateman, E.D.; Bousquet, J.; Keech, ML.; Busse, WW.; Clark, TJ. & Pedersen, SE. (2007). The correlation between asthma control and health status: the GOAL study. *Eur Respir J*, Vol.29, No.1, pp. 56-63, ISSN 0903-1936
- Bunge, EM.; Essink-Bot, ML.; Kobussen, MP.; Van Suijlekom-Smit, LW.; Moll, HA. & Raat, H. (2005). Reliability and validity of health status measurement by the TAPQOL. *Arch Dis Child*, Vol.90, No.4, pp. 351-358, ISSN 1468-2044
- Carr, AJ. & Higginson, IJ. (2001). Are quality of life measures patient centred? *BMJ*, Vol.322, No.7298, pp. 1357-1360, ISSN 0959-8138
- Christie, MJ.; French, D.; Sowden, A. & West, A. (1993). Development of child-centred disease-specific questionnaires for living with asthma. *Psychosom Med*, Vol.55, No.6, pp. 541-548, ISSN 0033-3174

- Dalheim-Englund, AC.; Rydström, I.; Rasmussen, BH.; Moller, C. & Sandman, PO. (2004). Having a child with asthma-quality of life for Swedish parents. *J Clin Nurs*, Vol.13, No.3, pp. 386-395, ISSN 0962-1067
- Dean, BB.; Calimlim, BC.; Kindermann, SL.; Khandker, RK. & Tinkelman, D. (2009). The impact of uncontrolled asthma on absenteeism and health-related quality of life. *J Asthma*, Vol.46, No.9, pp. 861-866, ISSN 1532-4303
- Dean, BB.; Calimlim, BC.; Sacco, P.; Aguilar, D.; Maykut, R. & Tinkelman, D. (2010). Uncontrolled asthma: assessing quality of life and productivity of children and their caregivers using a cross-sectional Internet-based survey. *Health Qual Life Outcomes*, Vol.8, pp. 96, ISSN 1477-7525
- Erickson, SR.; Munzenberger, PJ. & Plante, MJ.; Kirking, DM.; Hurwitz, ME. & Vanuya, RZ. (2002). Influence of sociodemographics on the health-related quality of life of pediatric patients with asthma and their caregivers. *J Asthma*, Vol.39, No.2, pp. 107-117, ISSN 0277-0903
- Everhart, RS. & Fiese, BH. (2009). Asthma severity and child quality of life in pediatric asthma: A systematic review. *Patient Educ Couns*, Vol.75, No.2, pp. 162-168, ISSN 0738-3991
- Farnik, M.; Pierzchała, W.; Brozek, G.; Zejda, JE. & Skrzypek, M. (2010). Quality of life protocol in the early asthma diagnosis in children. *Pediatr Pulmonol*, Vol.45, No.11, pp. 1095-1102, ISSN 1099-0496
- Fiese, BH.; Wamboldt, FS. & Anbar, RD. (2005). Family asthma management routines: connections to medical adherence and quality of life. *J Pediatrics*, Vol.146, No.2, pp. 171-176, ISSN 0022-3476
- Garro A. (2011). Health-related quality of life (HRQOL) in Latino families experiencing pediatric asthma. *J Child Health Care*, (Epub ahead of print), ISSN 1741-2889
- Global Initiative for Asthma (GINA) Executive Committee. (2010). Global strategy for asthma management and prevention, In: Global Initiative for Asthma (GINA), Available from: www.ginasthma.org
- Gorelick, MH.; Scribano, PV.; Stevens, MW. & Schultz, TR. (2003). Construct validity and responsiveness of the Child Health Questionnaire in children with acute asthma. *Ann Allergy Asthma Immunol*, Vol.90, No.6, pp. 622-628, ISSN 1081-1206
- Grootenhuis, MA.; Koopman, HM.; Verrips, EG.; Vogels, AG. & Last, BF. (2007). Health-related quality of life problems of children aged 8-11 years with a chronic disease. *Dev Neurorehabil*, Vol.10, No.1, pp. 27-33, ISSN 1751-8423
- Guilbert, TW.; Garris, C.; Jhingran, P.; Bonafede, M.; Tomaszewski, KJ.; Bonus, T.; Hahn, RM. & Schatz, M. (2011). Asthma that is not well-controlled is associated with increased healthcare utilization and decreased quality of life. *J Asthma*, Vol.48, No.2, pp. 126-132, ISSN 1532-4303
- Halterman, JS.; Yoos, HL.; Conn, KM.; Callahan, PM.; Montes, G.; Neely, TL. & Szilagyi, PG. (2004). The impact of childhood asthma on parental quality of life. *J Asthma*, Vol.41, No.6, pp. 645-653, ISSN 0277-0903
- Juniper, EF.; Guyatt, GH.; Feeny, DH.; Ferrie, PJ.; Griffith, LE. & Townsend, M. (1996). Measuring quality of life in children with asthma. *Qual Life Research*, Vol.5, No.1, pp. 35-46, ISSN 0962-9343
- Juniper, EF.; Guyatt, GH.; Feeny, DH.; Ferrie, PJ.; Griffith, LE. & Townsend, M. (1996). Measuring quality of life in the parents of children with asthma. *Qual Life Research*, Vol.5, No.1, pp. 27-34, ISSN 0962-9343

- Juniper, EF. (1997). How important is quality of life in pediatric asthma? *Pediatr Pulmonology*, Suppl.15, pp. 17-21, ISSN 1054-187X
- Juniper, EF.; Jenkins, C.; Price, MJ., Thwaites RMA. & James MH. (1999). Quality of life of asthma patients treated with salmeterol/fluticasone propionate combination product and budesonide (abstract). *Eur Respir J*, Vol.14, Suppl.30, pp. 370s
- Juniper, EF.; Svensson, K.; O'Byrne, PM.; Barnes, PJ.; Bauer, CA.; Lofdahl, CG.; Postma, DS.; Pauwels, RA.; Tattersfield, AE. & Ullman, A. (1999). Asthma quality of life during 1 year of treatment with budesonide with or without formoterol. *Eur Respir J*, Vol.14, No.5, pp. 1038-104, ISSN 0903-1936
- Le Coq, EM.; Colland, VT.; Boeke, AJ.; Boeke, P.; Bezemer, DP. & Van Eijk, JT. (2000). Reproducibility, construct validity, and responsiveness of the "How Are You?" (HAY), a self-report quality of life questionnaire for children with asthma. *J Asthma*, Vol.37, No.1, pp. 43-58, ISSN 0277-0903
- Magid, DJ.; Houry, D.; Ellis, J.; Lyons, E. & Rumsfeld, JS. (2004). Health-related quality of life predicts emergency department utilization for patients with asthma. *Ann Emerg Med*, Vol.43, No.5, pp. 551-557, ISSN 1097-6760
- Majani, G.; Baiardini, I.; Giardini, A.; Pasquali, M.; Tiozzo, M.; Tosca, M.; Cosentino, C.; La Grutta, S.; Marseglia, GL. &
- Canonica, GW. (2005). Impact of children's respiratory allergies on caregivers. *Monaldi Arch Chest Dis*, Vol.63, No.4, pp. 199-203, ISSN 1122-0643
- Masoli, M.; Fabian, D.; Holt, S. & Beasley, R. (2004). Global Initiative for Asthma (GINA) Program. The global burden of asthma: executive summary of the GINA Dissemination Committee report. *Allergy*, Vol.59, No.5, pp. 469-478, ISSN 0105-4538
- Merikallio, VJ.; Mustalahti, K., Remes, ST., Valovirta, EJ. & Kaila, M. (2005). Comparison of quality of life between asthmatic and healthy school children. *Pediatr Allergy Immunol*, Vol.16, No.4, pp. 332-340, ISSN 0905-6157
- Mohangoo, AD.; Essink-Bot, ML.; Juniper, EF.; Moll, HA.; De Koning, HJ. & Raat, H. (2005). Health-related quality of life in preschool children with wheezing and dyspnea: Preliminary results from a random general population sample. *Qual Life Research*, Vol. 14, No.8, pp. 1931-1936, ISSN 0962-9343
- Mohangoo, AD.; De Koning, HJ.; Mangunkusumo, RT. & Raat, H. (2007). Health-Related Quality of Life in Adolescents with Wheezing Attacks. *J Adolesc Health*, Vol.41, No.5, pp. 464-471, ISSN 1879-1972
- Mohangoo, AD.; De Koning, HJ.; De Jongste, JC.; Landgraf, JM.; Van der Wouden, JC.; Jaddoe, VW.; Hofman, A.; Moll, HA.; Mackenbach, JP. & Raat, H. (2011). Asthma-like symptoms in the first year of life and health-related quality of life at age 12 months: the Generation R study. *Qual Life Research*, In Press (Accepted: 2011-06-14), ISSN 1573-2649
- Mrazek D. (1992). Psychiatric complications of pediatric asthma. *Ann Allergy*, Vol.69, No.4, pp. 285-293, ISSN 0003-4738
- Osman, L. & Silverman, M. (1996). Measuring quality of life for young children with asthma and their families. *Eur Respir J*, Suppl.21, pp. 35s-41s, ISSN 0904-1850
- Pedersen, SE.; Hurd, SS.; Lemanske, RF. (Jr); Becker, A.; Zar, HJ.; Sly, PD.; Soto-Quiroz, M.; Wong, G. & Bateman, ED. (2011). Global strategy for the diagnosis and management of asthma in children 5 years and younger. *Pediatr Pulmonol*, Vol.46, No.1, pp. 1-17, ISSN 1099-0496
- Petersen, C.; Schmidt, S.; Power, M. & Bullinger, M. (2005). Development and pilot-testing of a health-related quality of life chronic generic module for children and adolescents

- with chronic health conditions: a European perspective. *Qual Life Research*, Vol.14, No.4, 1065-1077, ISSN 0962-9343
- Petsios, K.; Priftis, KN.; Tsoumakas, C.; Hatziaiorou, E.; Tsanakas, JN.; Galanis, P.; Antonogeorgos, G. & Matziou, V. (2011). Level of parent-asthmatic child agreement on health-related quality of life. *J Asthma*, Vol.48, No.3, pp. 286-297, ISSN 1532-4303
- Pont, LG.; Van der Molen, T.; Denig, P.; Van der Werf, GT. & Haaijer-Ruskamp, FM. (2004). Relationship between guideline treatment and health-related quality of life in asthma. *Eur Respir J*, Vol.23, No.5, pp. 718-722, ISSN 0903-1936
- Raat, H.; Bueving, H.J.; De Jongste, J.C.; Grol, MH.; Juniper, EF. & Van der Wouden, JC. (2005). Responsiveness, longitudinal- and cross-sectional construct validity of the Pediatric Asthma Quality of Life Questionnaire (PAQLQ) in Dutch children with asthma. *Qual Life Research*, Vol.14, No.1, pp. 265-272, ISSN 0962-9343
- Raat, H.; Mohangoo, A.D. & Grootenhuis, M.A. (2006). Pediatric health-related quality of life questionnaires in clinical trials. *Curr Opin Allergy Clin Immunol*, Vol.6, No.3, pp. 180-185, ISSN 1528-4050
- Sawyer, MG.; Spurrier, N.; Whaites, L.; Kennedy, D.; Martin, AJ. & Baghurst, P. (2000). The relationship between asthma severity, family functioning, and the health related quality of life of children with asthma. *Qual Life Research*, Vol.9, No.10, pp. 1105-1115, ISSN 0962-9343
- Sawyer, MG.; Spurrier, N.; Kennedy, D. & Martin, J. (2001). The relationship between the quality of life of children with asthma and family functioning. *J Asthma*, Vol.38, No.3, pp. 279-284, ISSN
- Sawyer, MG.; Reynolds, KE.; Couper, JJ.; French, DJ.; Kennedy, D.; Martin, J.; Staugas, R.; Ziaian, T. & Baghurst, PA. (2004). Health-related quality of life of children and adolescents with chronic illness - a two year prospective study. *Qual Life Research*, Vol.13, No.7, pp. 1309-1319, ISSN 0962-9343
- Skoner D. (2002). Outcome measures in childhood asthma. *Pediatrics*, Vol.109, Suppl.2, pp. 393-398, ISSN 1098-4275
- Spuijbroek, AT.; Oostenbrink, R.; Landgraf, JM.; Rietveld, E.; De Goede-Bolder, A.; Van Beeck, EF.; Van Baar, M.; Raat, H. & Moll, HA. (2011). Health-related quality of life in preschool children in five health conditions. *Qual Life Research*, Vol.20, No.5, pp. 779-786, ISSN 1573-2649
- Van den Bemt, L.; Kooijman, S.; Linssen, V.; Lucassen, P.; Muris, J.; Slabbers, G. & Schermer, T. (2010). How does asthma influence the daily life of children? Results of focus group interviews. *Health Qual Life Outcomes*, Vol.8, pp. 5, ISSN 1477-7525
- Varni, JW.; Burwinkle, TM.; Sherman, SA.; Hanna, K.; Berrin, SJ.; Malcarne, VL. & Chambers, HG. (2005). Health-related quality of life of children and adolescents with cerebral palsy: hearing the voices of the children. *Dev Med Child Neurol*, Vol.47, No.9, pp. 592-597, ISSN 0012-1622
- Vila, G.; Hayder, R.; Bertrand, C.; Falissard, B.; De Blic, J.; Mouren-Simeoni, MC. & Scheinmann, P. (2003). Psychopathology and quality of life for adolescents with asthma and their parents. *Psychosomatics*, Vol.44, No.4, pp. 319-328, ISSN 0033-3182
- World Health Organization (WHO), Division of Mental Health. (1993). Measurement of Quality of Life in Children. Available from:
www.who.int/mental_health/media/en/663.pdf



Allergic Diseases - Highlights in the Clinic, Mechanisms and Treatment

Edited by Prof. Celso Pereira

ISBN 978-953-51-0227-4

Hard cover, 554 pages

Publisher InTech

Published online 14, March, 2012

Published in print edition March, 2012

The present Edition "Allergic diseases - highlights in the clinic, mechanisms and treatment" aims to present some recent aspects related to one of the most prevalent daily clinical expression disease. The effort of a group of outstanding experts from many countries reflects a set of scientific studies very promising for a better clinical care and also to the treatment and control of the allergy. This book provides a valuable reference text in several topics of the clinical allergy and basic issues related to the immune system response. The inflammatory reaction understanding in allergic disease is clearly evidenced, as well as new strategies for further researches.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Esther Hafkamp-de Groen and Hein Raat (2012). Asthma and Health Related Quality of Life in Childhood and Adolescence, Allergic Diseases - Highlights in the Clinic, Mechanisms and Treatment, Prof. Celso Pereira (Ed.), ISBN: 978-953-51-0227-4, InTech, Available from: <http://www.intechopen.com/books/allergic-diseases-highlights-in-the-clinic-mechanisms-and-treatment/health-related-quality-of-life-in-children-with-asthma>

INTECH

open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen