

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,300

Open access books available

129,000

International authors and editors

155M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Role of Creatine Kinase – Hexokinase Complex in the Migration of Adenine Nucleotides in Mitochondrial Dysfunction

Elena Erlykina and Tatiana Sergeeva
*Nizhny Novgorod State Medical Academy
Russian Federation*

1. Introduction

Creatine phosphokinase (CK) (ATP: creatine phosphotransferase, EC 2.7.3.2.) is found in a variety of cells with high and fluctuating energy requirements. It catalyses the reversible transfer of the high-energy-N-phosphoryl group from phosphocreatine to ADP. Creatine kinase connects sites of energy production with sites of energy consumption (Dolder et al., 2001; Focant et al., 1970; Grossmann et al., 1985; Lipskaya et al., 1989; Walzel et al., 2002; Wyss, 2000).

There are known to be three cytosolic and two mitochondrial isoforms of CK. The more basic mitochondrial creatine kinase MiCK_b is accumulated in mitochondria of cardiac muscle and skeletal muscle. The more acidic mitochondrial creatine kinase MiCK_a was found in the brain (Eppenberger-Eberhardt et al., 1991; Fridman, Roberts, 1994).

Creatine kinase can exist in two interconvertible forms: dimer and octamer (Eriksson et al., 1998; Shen et al., 2002). Creatine kinase binds to the outer leaflet of the entire inner mitochondrial membrane and is specifically enriched in the so-called contact sites where inner and outer membranes are in close proximity (Boero et al., 2003; Chen et al., 1994; Lin et al., al., 1996; Wang et al., 2005).

A change in the octamer/dimer ratio may influence on the association behavior of mitochondrial creatine kinase in general and thus modulate mitochondrial energy flux (Brdiczka, 2003; Dolder et al., 2001; Schnyder et al., 1995).

Mitochondrial creatine kinase forms the functional microcompartment together with the mitochondrial porin (voltage-dependent anion channel) in the outer membrane and as well as the transmembrane protein adenine nucleotide translocase in the inner membrane (Fritz-Wolf et al., 1996; Kaldis, Wallimann, 1994; Schnyder et al., 1988).

Hexokinase (HK) (ATP:D-hexokinase-6-phosphotransferase, EC 2.7.11) is the enzyme with variable cellular localization (Mulichak et al., 1998; Xie & Wilson, 1990).

The type I isoenzyme of mammalian hexokinase is ubiquitously expressed in mammalian tissues but is found particularly at high levels in the brain where it plays an important role in regulating the rate of cerebral glucose metabolism (Schwab & Wilson, 1989; Wilson, 1985).

The major portion of the hexokinase activity in the brain is associated with mitochondria. About 85% of hexokinase is bound to the outer mitochondrial membrane, forming the specific complex with porin (Magnani et al., 1982; Redker et al., 1972; Wilson, 1995). This

physical proximity provides the basis for functional interaction between glucose phosphorylation by hexokinase and mitochondrial ATP production by oxidative phosphorylation with resulting coordination of the glycolytic and oxidative phases of glucose metabolism (Aleshin, 1998; Rosano et al., 1999). The outer mitochondrial membrane protein - porin, which forms the transmembrane channel, is responsible for specific interaction with hexokinase (Aflalo & Azoulay, 1998; Linden et al., 1982; Schlattner et al., 2001; Vyssokikh & Brdiczka, 2003).

The preferential mitochondrial localization of hexokinase in rat brain provides a predominant access to ATP, generated in mitochondria. The ADP produced by hexokinase activity is known to control both membrane potential and reactive oxygen species generation (Rose & Warms, 1967; Smith and Wilson, 1991; Viitanen et al., 1984; Wilson, 1980).

Thus both enzymes - creatine kinase and hexokinase - play an important role in dynamic compartmentation of adenine nucleotides.

Mitochondrial creatine kinase is a key enzyme of oxidative cellular energy metabolism in the brain (Bessman, 1981; Guo et al., 2003; Hemmer et al., 1994; Levin et al., 1990; Takagi et al., 2001; Wallimann et al., 1992; Wallimann et al., 1998). Hexokinase is an enzyme involved in the first step of glycolysis. Mitochondrial creatine kinase - hexokinase complex takes part in transport of adenine nucleotides from mitochondria to cytoplasm. Functioning of this complex depends on interaction of enzymes with the mitochondrial membrane and the oligomeric state of mitochondrial creatine kinase.

Mitochondrial dysfunction is one of the main reasons of the pathological changes in cerebral ischemia (Clostre, 2001; Delivoria-Papadopoulos et al., 2007; Fiscum, 2000; Kuznetsov, Margreiter, 2009; Mattson, Liu, 2002; Sas et al., 2007; Siesjo, 1999).

Stroke is a leading cause of disability and death in many countries. Understanding the molecular mechanisms of ischemic injury helps to find the novel therapeutic strategies for stroke. 80% of human strokes are ischemic in origin (Levine et al., 1992; Sappey-Marini et al., 2002; Ueda et al., 2000).

Thus experimental models of cerebral ischemia have been developed in an attempt to closely mimic the changes that occur during and after human ischemic stroke. Changes in the amount and activity of enzyme proteins are critical factors in regulating intracellular metabolism under ischemic conditions (Cherubini et al., 2000; Dos Santos et al., 2004; Maulik et al., 1999; Rauchova et al., 2002).

According to modern data, membrane-associated enzyme in contrast to soluble enzyme has other catalytic properties (Beutner et al., 1998; da-Silva et al., 2004; Dolder et al., 2001; Kellershohn & Ricard, 1994; Linden et al., 1982; Lyubarev, 1997; Ovadi & Srere, 2000). The reverse adsorption on the mitochondrial membrane is controlled by ions and metabolites thus broadening the regulatory possibility of the cells under hypoxic conditions.

2. Materials and methods

Animals

Experiments were performed on male outbred albino rats weighing 150-180 g.

Cerebral ischemia was produced by bilateral ligation of the common carotid arteries. The animals were anesthetized with nembutal (30 mg/kg intraperitoneally). The brain tissue was examined 30 minutes (acute ischemia), 1.5, 4, 18 hours after surgical impairment of cerebral hemodynamics.

The animals were divided into two groups due to their physiological state after acute ischemia (severe and moderate). The severity of ischemia was estimated according to the behavior of rats, the respiratory rate and survival one. In acute severe ischemia the rats after ligation of the common carotid arteries were in severe state: they were passive, in lateral recumbent position, with agonal breathing (20-30 times per minute with respiratory arrest). In acute moderate ischemia the state of animals was satisfactory: they were active, moved in a cage, the respiratory rate was 50-70 times per minute. Due to the increase in the severity of general physiological state of rats in case of long-term ischemia we could not divide the animals into two groups; they were included into one group.

Preparation of brain tissue

The mitochondrial fraction of the brain was isolated by differential centrifugation (Fonyo, Somogy, 1960; Dizhe et al., 2003). The brain tissue was homogenized at 4 °C in a medium containing 0.32 M sucrose, 10 mM tris-HCl, 1 mM EDTA, pH 7.4. The total tissue homogenate was centrifuged at 2000 g for 10 minutes. The resulting supernatant was collected and centrifuged further at 12 500g for 15 minutes. The pellet containing mitochondria was resuspended in 0.32 M sucrose and centrifuged at 16 500g for 15 minutes. The fraction enriched mitochondria was collected and washed by 0.32 M sucrose.

The mitochondria were then swollen by incubation in distilled water (at a protein concentration of 1 mg) for 30 minutes, followed by centrifugation at 20 000g for 30 minutes. The resulting supernatant was collected for further analysis. The pellet containing mitochondrial membranes was resuspended in 0.32 M sucrose with 0.25 M dithiothreitol, pH 7.4.

Enzyme assay

Creatine kinase activity was measured by the pH-stat method using ADP and creatine phosphate as substrates (Kuby, Noltman, 1962). The velocity of the creatine kinase reaction is estimated by the change in pH. The reaction mixture (3 ml) contained (final concentration): 0.25 M sucrose, 2.5 mM tris-HCl, 12 mM MgCl₂, 10 mM KCl, 0.25 mM dithiothreitol, 5 mM creatine phosphate, 2 mM ADP. The reaction was started by addition of 100 µg protein. Then the mixture was titrated by addition of 10 µl 0.1 N HCl.

Creatine kinase activity is expressed as 1 unit corresponds to 1 µg-equ H⁺/ min per 1 mg of protein.

Hexokinase activity was measured spectrophotometrically (Felgner, Wilson 1976). The reaction mixture (3 ml) contained (final concentration): 50 mM tris-HCl, pH 8.0, 2 mM glucose, 2 mM ATP, 5 mM MgCl₂, 0.25 mM NADP, 0.4 IU/min glucose-6-phosphate dehydrogenase. The reaction was started by addition of 100 µg protein.

Hexokinase activity is expressed as follows: 1 unit corresponds to 1 nmol of NADP transformed/ min per 1 mg of protein.

Solubilization of creatine kinase

Mitochondria were resuspended in the proper (0.1M KCl; K-Na phosphate buffer 0.1-1.75 M, 0.5% (v/v) Triton X-100; 0.1% deoxycholate Na) solubilizing solution and incubated for 30 minutes. The samples were centrifuged at 4°C and 20 000g, 60 minutes. Percentage of solubilization was determined as the difference of the activity before and after solubilization of the enzyme.

Solubilization of hexokinase

Mitochondria were resuspended at a protein concentration of 0.5-1 mg/ml in 0.1M tris-HCl pH 6.6, 0.1 M KCl or 0.5% (v/v) Triton X-100. After incubation for 30 minutes on ice, with

occasional mixing, the samples were centrifuged at 4°C and 20 000g for 30 minutes. Percentage of solubilization was determined as the difference of the activity before and after solubilization of the enzyme.

According to Wilson (2003) there are 2 types of binding sites for hexokinase on brain mitochondria. Hexokinase is readily solubilized from Type A sites by glucose-6-phosphate while hexokinase bound to Type B sites remains bound even in the presence of glucose-6-phosphate.

Mitochondria were resuspended in 2 mM glucose-6-phosphate; tris-HCl buffer, pH 8 and incubated for 30 minutes at the room temperature, and centrifuged 100 000g for 15 minutes. Aliquots of supernatant contain hexokinase Type A.

The sediment of mitochondria which contained hexokinase type B was resuspended again in 0,32 M sucrose, 0.5% (v/v) Triton X-100, 0.1M tris-HCl, pH 8. After incubation for 5 minutes on ice the samples were centrifuged at 20 000g for 10 minutes. The sediment was resuspended again in 0.32M sucrose, 0.1M KCl, 1% (v/v) Triton X-100, 0.1M tris-HCl, pH 8. After incubation for 20 minutes on ice the samples were centrifuged at 20 000g for 10 minutes. The aliquots of supernatant contained hexokinase Type B.

Dissociation of creatine kinase

Mitochondrial creatine kinase was dissociated by incubation of the total mitochondrial fraction and mitochondrial membrane pellet with substrates for the transition-state analogue complex (MgCl₂, ADP, KNO₃, and creatine) at 4 °C for 2 hours (Lipskaya et al., 1989).

The free radical oxidation intensity assay

The intensity of the free radical oxidation (FRO) was estimated by the method of H₂O₂, Fe²⁺-induced chemiluminescence on a BChL-07 biochemiluminometer. This method is based on the catalytic decomposition of hydrogen peroxide by ions of metal with variable valency (bivalent iron) (the Phenton reaction). The reaction mixture contained: 0,05 mM Fe₂SO₄, a phosphate buffer and a mitochondrial fraction. The reaction was started by addition of 2 % solution of hydrogen peroxide. Proceeding process of free radical oxidation was registered within 30 seconds. It is the time of the greatest information about its intensity. The ideal curve of the process is presented in figure 1.

The following parameters are the most informative for the estimation of the chemiluminescence intensity: the total luminescence yield (S, enables to estimate a balance between lipid peroxidation and antioxidants), maximum flash amplitude (I_{max}, shows a potential ability of the biological sample to free radical oxidation), and K index characterizing antioxidant potential were used as integral parameters of chemiluminescence (Kuzmina et al., 2009).

The protein concentration assay

Protein concentration was measured by the method of Bredford (Bredford & Spector, 1978).

Statistical analysis

The data are expressed as mean and standard error of the mean (SEM). The results were analyzed by means of Primer of Biostatistics 4.03 (Glantz, 2005). The significance of differences between the samples was evaluated by Student's test. The level of significance was set at $p < 0.05$.




Fig. 1. The kinetic curve of the development of an induced chemiluminescence signal.

3. Results

Two forms of creatine kinase and hexokinase – membrane and soluble – were found in the brain.

The activity of membrane-associated creatine kinase equals to half of the enzyme activity in mitochondria. In contrast hexokinase activity is concentrated on the outer surface of the mitochondrial membrane (Table 1).

Form of the enzyme	Creatine kinase, U/ mg*min.	Hexokinase, U/ mg*min.
Membrane-associated	1.40±0.07 n=23	11.56±0.19 n=12
Soluble	1.78±0.10 n=19	1.53±0.04 n=12

Table 1. The distribution of creatine kinase and hexokinase activity between membrane-associated and soluble forms of the enzymes

Catalytic and kinetic properties of mitochondrial creatine kinase and hexokinase were shown to depend on the interaction with the membrane.

Different solubilizing agents (electrolyte, detergent and the endogenous metabolite glucose-6-phosphate) were used to analyze the character of interaction of hexokinase with the mitochondrial membrane.

All these agents solubilized only a third of the hexokinase activity, and only the sequence of action of electrolyte, detergent and glucose-6-phosphate removed the enzyme from the mitochondrial membrane (Fig. 2). Thus it shows the lability of protein-protein interaction and the possibility of its regulation under the certain pathological conditions.




Fig. 2. Solubilization (%) of hexokinase from mitochondrial membranes of intact rats.

Creatine kinase can not dissociate from the mitochondrial membrane even in the presence of the simultaneous action of electrolyte and detergent (Fig. 3).

So the brain creatine kinase exists in different molecular forms: the first - soluble, which is located in the intermembrane space, the second is associated, which is loosely bound to the inner mitochondrial membrane and under the certain solubilizing agents can remove into the intermembrane space, the third form (about 18%) is tightly bound with the membrane.

In mitochondria from intact animals, mitochondrial creatine kinase presents as a mixture of two oligomeric forms (dimer and octamer; 65 and 35%, respectively). We consider that the tightly bound creatine kinase to exist mainly in the contact sites in the octamer form.




Fig. 3. Solubilization (%) of creatine kinase from mitochondrial membranes of intact rats.

Rebinding of both phosphokinases with the membrane changes their catalytic properties. Binding of hexokinase with the membrane increases the velocity of the reaction in 3 fold, but the kinetic behavior is not changed (Fig. 4).




Fig. 4. Dependence of V_0 on concentration of MgATP of membrane-associated hexokinase of intact rats.

Creatine kinase has different types of kinetic behavior (Fig. 5-7). We consider that the membrane associated form of the enzyme binds by ionic interaction with the membrane and the character of the curve reveals the classical kinetic behavior (Fig. 6). The tightly bound




Fig. 5. Kinetic of creatine kinase reaction in total mitochondrial fraction of intact rats.

form of membrane enzyme has the abnormal kinetic behavior due to ionic and hydrophobic interaction (Fig. 7). These data describe the role of specific microenvironment in the modification of the enzyme properties.




Fig. 6. Kinetic of creatine kinase reaction on mitochondrial membrane of intact rats.




Fig. 7. Kinetic of creatine kinase reaction on mitochondrial membrane after solubilization by phosphate buffer of intact rats.

Thus the catalytic properties depend on the binding with the membrane and this process is controlled by the endogenous metabolites and the functional state of mitochondria.

All forms of hypoxia and ischemia are accompanied by activation of free radical oxidation (Ayer, Zhang, 2008; da-Silva et al., 2004; Kuznetsov, Margreiter, 2009; Meyer et al., 2006;

Wang et al., 2005). As a result of this activation the properties of the mitochondrial membrane-associated enzymes are changed in acute ischemia.

Severe ischemia reduced the binding of the investigated enzymes with the membrane (Fig. 8, 9). The activity of the enzymes decreased 2 fold for creatine kinase and 3 fold for hexokinase. Glucose-6-phosphate and the products of membrane degradation inhibited hexokinase activity in cerebral ischemia (Ishibashi, 1999; Wilson et al., 2000).




Fig. 8. Membrane associated hexokinase activity on the membrane before and after solubilization by 0.1M KCl of intact rats and in 30 minutes ischemia.




Fig. 9. Creatine kinase activity on the membrane of intact rats and in 30 minutes ischemia before and after solubilization by phosphate buffer.

The bar diagrams display average activities with error bars representing the standard deviation. * $p < 0.05$ versus intact animals. ** $p < 0.05$ versus initial hexokinase activity on the membrane within the same group.

The bar diagrams display average activities with error bars representing the standard deviation. $p < 0.05$ versus intact animals. ** $p < 0.05$ versus initial creatine kinase activity on the membrane within the same group.

In the second group (moderate ischemia) the activity of hexokinase was increased by 29% and by 92% for creatine kinase in comparison with intact animals. After the solubilization of the hexokinase by 0.1 M KCl the enzyme lost 38% of the initial activity. The effect of solubilization for membrane-bound creatine kinase was 69% instead of 35% for the intact rats.

The study of behavior of creatine kinase revealed the modification of its properties in acute ischemia. They differed significantly from those of the intact rats. It is connected with the realization of interconvertible transformation of oligomeric subunits of creatine kinase. This changing in kinetic behavior provides the higher sensitivity of the enzyme to the changes in substrate concentration (Fig. 10-13).




Fig. 10. Dependence of V_0 on concentration of phosphocreatine in total mitochondrial fraction in moderate ischemia




Fig. 11. Dependence of V_0 on concentration of phosphocreatine in fraction of mitochondrial membranes in moderate ischemia




Fig. 12. Dependence of V_0 on concentration of phosphocreatine in total mitochondrial fraction in severe ischemia.

The study of the creatine kinase reaction in the group of animals after severe ischemia showed the abnormal kinetic behavior of the enzyme, the appearance of the intermediate plateau at the low concentration (0.3-0.4 mM) of creatine phosphate, the V_0 decreased 1.4-2 fold in comparison with intact rats.




Fig. 13. Dependence of V_0 on concentration of phosphocreatine in fraction of mitochondrial membranes in severe ischemia.

The reversibility of these alterations has been shown during the increasing of duration of ischemia.

Increasing of the duration of cerebral ischemia to 4 and 18 hours was accompanied by changes in activity distribution for hexokinase. The activity of hexokinase progressively increased. The level of activity of the enzyme under these conditions was higher than in acute ischemia. However, the level of hexokinase activity in animals during long-term ischemia remained lower than in intact specimens (Fig. 14).

The bar diagrams display average activities with error bars representing the standard deviation. * $p < 0.05$ versus intact animals. ** $p < 0.05$ versus severe ischemia (30 minutes).

An increase in duration of cerebral ischemia influenced on the adsorption properties of hexokinase. Solubilization of hexokinase by 0.1 KCl was accompanied by decrease in the activity of the enzyme by 78% in acute cerebral ischemia. The percentage of solubilizing enzyme was 37% in 1.5 hours ischemia. It was by 11% higher than in intact specimens. The percentage of solubilizing enzyme was 30% and 27% in 4 hours and 18 hours ischemia, respectively. It did not differ from that in intact animals.

Therefore increase in the duration of cerebral ischemia was followed by an increase in the resistance of membrane structures. These changes were manifested in reduction of hexokinase desorption from the mitochondrial membrane.




Fig. 14. The activity of membrane-bound hexokinase before and after solubilization by 0.1M KCl in cerebral ischemia

The cerebral ischemia causes the imbalance between reactive oxygen species production and the level of antioxidant defense, which leads to oxidative stress. Neuronal membranes contain a considerable amount of unsaturated lipids. The low level of activity of antioxidant enzymes and formation of free radicals in neurochemical reactions provide conditions for lipid oxidation and induce enzyme modification. To evaluate the state of membranes, the intensity of free radical oxidation and antioxidant properties of the brain tissue were estimated in various periods of ischemia (Table 2).

Experimental groups	Imax, mV	S, imp. *30 sec.	K=1/S
Intact animals	1.02±0.03 n=8	10.64±0.34 n=8	0.094
Ischemia, 30 minutes	1.89±0.07* n=7	17.22±0.98* n=7	0.058
Ischemia, 1.5 hours	1.52±0.04* n=7	15.18±0.91* n=7	0.066
Ischemia, 4 hours	1.47±0.05* n=8	14.55±0.03* n=8	0.068
Ischemia, 18 hours	1.08±0.11 n=8	12.63±0.79* n=8	0.079

* $p < 0.05$ versus intact animals

Table 2. The intensity of free radical oxidation and the activity of antioxidant system in the brain.

Various characteristics of chemiluminescence (maximum flash amplitude and total yield of slow flash) in the mitochondrial fraction were elevated during various periods of cerebral ischemia. These changes reflect activation of free radical processes in the brain. Total yield of slow flash was 1.6-fold and 1.4-fold higher than in the intact animals in 30 min and 1.5 hours ischemia.

Parameters of free radical oxidation (I_{max} and S) remained practically unchanged by 4 hours ischemia. The increase in the duration of cerebral ischemia to 18 hours was accompanied by a decrease in the intensity of free radical oxidation. Parameter I_{max} did not differ from the corresponding parameter in intact animals.

Acute ischemia was not only followed by damage of the cell membrane structures and activation of free radical oxidation, but also induced the antioxidant system (Dziennis et al., 2008; Lai et al., 2003; Perez-Pinzon et al., 2005; Suzuki et al., 1997).

The K index serves as a criterion for the antioxidant potential of the cell. The level of the antioxidant activity of the brain tissue was elevated after ischemia for 1.5 and 4 hours. The conclusion was derived from the decrease in this index. By 18 hours ischemia, the K index did not differ from that in intact animals.

These data indicate that the prooxidant/antioxidant ratio returns to normal with increasing in the duration of cerebral circulatory disorder. The observed changes are probably related to activation of defense protein synthesis, which increases the resistance of membrane structures to the adverse effect of ischemia.

The kinetic curve for hexokinase was shown to have hyperbola form in various periods of ischemia except for 1.5 hours ischemia (Fig. 15). MgATP did not inhibit hexokinase when increasing the duration of ischemia. By 18 hours, the K_m (0.13 mM) is 2-fold lower than in intact animals (0.26 mM) and is 5-fold lower than in 30 minutes ischemia (0.7 mM).




Fig. 15. Kinetic of the hexokinase reaction.

The major problem in the involvement of cell structures in the regulation of enzyme activity is the dependence of enzyme properties on the association of this enzyme with the membrane under conditions of functional changes in the organism.

Mitochondrial creatine kinase is associated with mitochondrial membranes due to the forces of electrostatic and hydrophobic interaction. Cerebral ischemia was followed by changes in the activity of associated and tightly-bound mitochondrial creatine kinase (Fig. 16).

The activity of associated mitochondrial creatine kinase increased in comparison with intact animals in 1.5 hours ischemia. However the activity of tightly-bound creatine kinase did not change. The activity of tightly-bound mitochondrial creatine kinase increased and the ratio between two forms of mitochondrial creatine kinase restored in 4 hours ischemia. By 18 hours, the percentage of tightly-bound form of the enzyme reached 90%.




Fig. 16. The activity of membrane-associated creatine kinase before and after solubilization by phosphate buffer in cerebral ischemia.

The bar diagrams display average activities with error bars representing the standard deviation. * $p < 0.05$ versus intact animals. ** $p < 0.05$ versus severe ischemia (30 minutes).

Cerebral ischemia was shown to change the kinetic properties of mitochondrial creatine kinase (Fig. 17). Mitochondrial creatine kinase showed abnormal kinetic with the appearance of intermediate plateau. By 18 hours, the kinetic curve acquired a hyperbola form.

In mitochondria, mitochondrial creatine kinase is presented by two oligomeric forms (dimer and octamer). They are characterized by dynamic equilibrium (Lipskaya et al., 1989). The transition-state analogue complex of mitochondrial creatine kinase was induced to evaluate the ratio between oligomeric forms of this enzyme under conditions of cerebral circulatory disorders.

Cerebral ischemia changes the dimer/octamer ratio. This ratio is shifted toward the formation of dimers after 30-min ischemia (79%). Phospholipids serve as the structural elements of membranes that are bound to mitochondrial creatine kinase. Membrane binding properties of mitochondrial creatine kinase depend strongly on the protein dimer/octamer ratio and degree of lipid oxidation. Activation of free radical oxidation during acute ischemia is probably followed by partial dissociation of octamers to dimers. Increasing of the duration of ischemia to 18 hours was followed by an increase in the octamer ratio (53%).




Fig. 17. Kinetic of the membrane-bound creatine kinase reaction

Published data suggest that octameric mitochondrial creatine kinase contributes to the appearance and strengthening of contact sites, which increases the efficiency of energy formation in brain mitochondria, consolidates the membrane structure, and determines the resistance of membranes to the adverse effect of hypoxia (Gross, Wallimann, 1995; Koufen et al., 1999; Lenz et al., 2007; Meyer et al., 2006). The existence of two oligomeric forms of this enzyme probably maintains the near-equilibrium state of reaction in a wide range of physiological conditions.

4. Conclusion

The results indicate that catalytic properties of mitochondrial creatine kinase and hexokinase depend on the functional interaction with mitochondrial membranes.

Acute ischemia impairs enzyme interaction with the mitochondrial membrane. Increasing in the duration of ischemia is not only followed by injury and dysfunction, but also activates the defense systems in the nervous tissue. It is manifested in the decrease in the intensity of free radical oxidation, increase in the percentage of tightly-bound mitochondrial creatine kinase, changes in kinetic properties of the enzyme and change in the dimer/octamer ratio toward the formation of octamer for the mitochondrial creatine kinase. These changes stabilize the mitochondrial creatine kinase complex. In contrast, increase in the duration of ischemia is accompanied by the decrease in the hexokinase activity on the membrane in spite the fact that it becomes higher than in acute 30 min ischemia, but the percentage of solubilizing enzyme does not differ from that in intact animals.

Therefore, the long-term ischemia leads to stabilization of the functional interaction between hexokinase and creatine kinase complex with the mitochondrial membranes at a new level,

providing the adequate energy supply of the nervous cells due to the new adaptive conditions.

5. References

- Aflalo, C. & Azoulay, H. (1998). Association of rat brain hexokinase to yeast mitochondria: effect of environmental factors and the source of porin. *J. Bioenerg. Biomembr.*, Vol. 30, pp. 245-255.
- Aleshin, A.E.; Fromm, H.J. & Honzatko, R.B. (1998). Multiple crystal form of hexokinase I: new insights regarding conformational dynamics, subunit interactions, and membrane association. *FEBS Lett*, Vol. 434, pp. 42-46.
- Bessman, S.P. & Carpenter, C.L. (1985). The creatine-creatine phosphate energy shuttle. *Annu. Rev. Biochem.*, Vol. 54, pp. 831-862.
- Beutner, G.; Ruck, A., Riede, B. & Brdiczka, D. (1998). Complexes between porin, hexokinase, mitochondrial creatine kinase and adenylate translocator display properties of the permeability transition pore. Implication for regulation of permeability transition by the kinases. *Biochem. Biophys. Acta*, Vol. 1368, pp. 7-18.
- Boero, J.; Qin, W., Cheg, J., Woolsey, T.A., Strauss, A.W. & Rhuchua, Z. (2003). Restricted neuronal expression of ubiquitous mitochondrial creatine kinase; changing patterns in development and with increased activity. *Mol. Cell Biochem*, Vol. 244, No 102, pp. 69-76.
- Brdiczka, D. (1994). Function of the outer mitochondrial compartment in regulation of energy metabolism. *Biochem. Biophys. Acta*, Vol. 1187, No 2, pp. 264-269.
- Bredford, T. & Spector, T. (1978). Refinement of the coomassie blue method of protein quantitation. A simple and linear spectrophotometric assay for less than or equal to 0.5 to 50 microgram of protein. *Anal. Biochem*, Vol. 86, No 1, pp. 142-146.
- Chen, L.; Roberts, R. & Friedman, D.L. (1995). Expression of brain-type creatine kinase and ubiquitous mitochondrial creatine kinase in the fetal rat brain: evidence for a nuclear energy shuttle. *J. Comp. Neurol.*, Vol. 363, No 3, pp. 389-401.
- Cherubini, A.; Ruggiero, C., Polidori, M.C. & Mecocci P. (2005). Potential markers of oxidative stress in stroke. *Free Radic. Biol. Med.*, Vol. 39, No 7, pp. 841-852.
- Clostre, F. (2001). Mitochondria: recent pathophysiological discoveries and new therapeutic perspectives. *Ann. Pharm. Fr.*, Vol. 59, No 1, pp. 3-21.
- da-Silva, W.S.; Gomez-Puyou, A., de Gomez-Puyou, M.T., Moreno-Sanchez, R., De Felice, F.G., de Meis, L., Oliveira, M.F. & Galina, A. (2004). Mitochondrial bound hexokinase activity as a preventive antioxidant defense: steady-state ADP formation as a regulatory mechanism of membrane potential and reactive oxygen species generation in mitochondria. *J. Biol. Chem.*, Vol. 279, No 38, pp. 39846-39855.
- Delivoria-Papadopoulos, M.; Gorn M., Ashraf, Q.M. & Mishra, O.P. (2007). ATP and cytochrome c-dependent activation of caspase-9 during hypoxia in the cerebral cortex of newborn piglets. *Neurosci. Lett.*, Vol. 429, No 2-3, pp. 115-119.
- Dizhe, G. P.; Eshchenko, N.D., Dizhe, A.A. & Krasovskaya, I.E. (2003). *Introduction to the Technique of Biochemical Assay* [in Russian], St. Petersburg, p. 86.
- Dolder, M., Wendt, S. & Wallimann, T. (2001). Mitochondrial creatine kinase in contact sites: interaction with porine and adenine nucleotide translocase, role in permeability

- transition and sensitivity to oxidative damage. *Biol. Signals Recept.*, Vol. 10, No 1-2, pp. 93-111.
- Dos Santos, P.; Laclau, M.N., Boudina, S. & Garlid, K.D. (2004). Alteration of the bioenergetics systems of the cell in acute chronic myocardial ischemia. *Mol. Cell Biochem.*, Vol. 256-257, No 1-2, pp. 157-166.
- Eppenberger-Eberhardt, M.; Riesinger, I., Messerli, M. et al. (1991). Adult rat cardiomyocytes cultured in creatine-deficient medium display large mitochondria with paracrystalline inclusions, enriched for creatine kinase. *J. Cell Biol.*, Vol. 113, pp. 289-302.
- Eriksson, O.; Fontaine, E. & Bernandi, P. (1998). Chemical modification of arginines by 2,3-butanedione and phenylglyoxal causes closure of the mitochondrial permeability transition pore. *Biol. Chem.*, Vol. 273, pp. 12669-12674.
- Felgner, P.L. & Wilson, J.E. (1976) Purification of nonbindable and membrane bindable mitochondrial hexokinase from rat brain. *Biochem. Biophys. Res. Commun.*, Vol. 68, No 2, pp. 592-597.
- Fiscum, G. (2000) Mitochondrial participation in ischemic and traumatic neural cell death. *J. Neurotrauma*, Vol. 17, No 10, pp. 843-855.
- Focant, T. (1970). Isolement et propriétés de la créatine-kinase de muscle lisse de boeuf. *FEBS Lett.*, Vol. 10, pp. 57-61.
- Fonyo, A. & Somogyi, J. (1960). The phosphorylation of adenosinediphosphate and glucose in isolated brain mitochondria at different osmotic concentrations. *Acta physiol. Acad. Sci., Hung.*, Vol. 8, No 3, pp. 191.
- Fridman, D.L. & Roberts, R. (1994). Compartmentation of brain type creatine kinase and ubiquitous mitochondrial creatine kinase in neurons; evidence for a creatine phosphate shuttle in adult rat brain. *J. Comp. Neurol.*, Vol. 343, No 3, pp. 500-511.
- Fritz-Wolf, K. ; Schnyder, T. , Wallimann, T. & Kabsch, W (1996). Structure of mitochondrial creatine kinase. *J. Nature*, Vol. 381, No 6580, pp. 341-345.
- Glantz, S.A. (2005). *Primer of Biostatistics*, 6th ed., McGraw-Hill Medical Publishing, p. 500.
- Gross, M. & Wallimann, T. (1995). Dimer-dimer interactions in octameric mitochondrial creatine kinase. *Biochemistry*, Vol. 34, No 20, pp. 6660-6667.
- Grossman, S. & Mixon, D. (1985). Further Characterization of the reassembly of creatine kinase and effect of substrate. *Arch. Biochem. Biophys.*, Vol. 236, pp. 797.
- Guo, Z.; Wang, Z. & Wang, X. (2003). Studies on the stability of creatine kinase isoenzymes. *Biochem Cell Biol.*, Vol.81, No 1, pp. 9-16.
- Hemmer, W.; Zanolla, E., Furter-Graves, E.M., Eppenberger, H.M. & Wallimann, T. (1994). Creatine kinase isozymes in chicken cerebellum: specific localization of brain-type creatine kinase in Bergmann glial cells and muscle-type creatine kinase in Purkinje neurons. *Eur. J. Neurosci*, Vol. 6, No 4, pp. 538-549.
- Ishibashi, S. (1999). Cooperation of membrane proteins and cytosolic proteins in metabolic regulation – involvement of binding of hexokinase to mitochondria in regulation of glucose metabolism and association and complex formation between membrane proteins and cytosolic proteins in regulation of active oxygen production. *Yakugaku Zasshi*, Vol. 119, No 1, pp. 16-35.

- Kaldis, P.; Wallimann T. (1995). Functional differences between dimeric and octameric mitochondrial creatine kinase. *Biochem. J.*, Vol. 308, pp. 623-627.
- Kellershohn, N. & Ricard, J. (1994). Coordination of catalytic activities within enzyme complex. *Eur. J. Biochem.*, Vol. 220, No 2, pp. 955-961.
- Koufen, P. et al. (1999). Free radical-induced inactivation of creatine kinase: influence on the octameric and dimeric states of the mitochondrial enzyme (Mi_b-CK). *Biochem. J.*, Vol. 344, pp. 413-417.
- Kuby, S.A.; Noda, L. & Lardy, H.A. (1954). Adenosinetriphosphate-creatine transphosphorylase. I. Isolation of the crystalline enzyme from rabbit muscle. *J. Biol. Chem.*, Vol. 209, No 1, pp. 191-201.
- Kuzmina, E.I. et al. (2009). The evaluation of antioxidant action of medications on modeling systems in vitro. *Revista de Ozonoterapia*, Vol. 3, suppl., No 1, pp. 67-68.
- Kuznetsov, A.V.; Margreiter, R. (2009). Heterogeneity of mitochondria and mitochondrial function within cells as another level of mitochondrial complexity. *Int. J. Mol. Sci.*, Vol. 10, No 4, pp. 1911-1929.
- Lai, J.C.; White, B.K., Buerstatte, C.R. & Haddad G.G., Novotny, E.J. Jr. & Behar, K.L. (2003). Chronic hypoxia in development selectively alters the activities of key enzymes of glucose oxidative metabolism in brain regions. *Neurochem. Res.*, Vol. 28, No 6, pp. 933-940.
- Lenz, H.; Schmidt, M., Welge, V., Kueper, T., Schlattner, U., Wallimann, T., Elsässer, H.P., Wittern, K.P., Wenck, H., Staeb, F. & Blatt, T. (2007). Inhibition of cytosolic and mitochondrial creatine kinase by siRNA in HaCaT – and HeLaS3-cells affects cell viability and mitochondrial morphology. *Mol. Cell. Biochem.*, Vol. 306, pp. 153-162.
- Levin, R.M.; Longhurst, P.A., Levin, S.S., Haugaard, N. & Wein, A.J. (1990). Creatine kinase activity of urinary bladder and skeletal muscle from control and streptozotocin-diabetic rats. *Mol. Cell Biochem.*, Vol. 97, No 2, pp. 153-159.
- Levine, S.R. et al. Human focal cerebral ischemia: evaluation of brain pH and energy metabolism with P-31 NMR spectroscopy. *Radiology*, Vol. 185, No 2, pp. 537-544.
- Lin, L.; Perryman, M.B., Friedman, D., Roberts, R. & Ma, T.S. (1994). Determination of the catalytic site of creatine kinase by site-directed mutagenesis. *Biochim Biophys Acta*, Vol. 1206, No 1, pp. 97-104.
- Linden, M.; Gellerfors, P. & Nelson, B.D. (1982). Pore protein and the hexokinase-binding protein from the outer membrane of rat liver mitochondria are identical. *FEBS Lett*, Vol. 141, pp. 189-192.
- Lipskaya, T.Yu.; Trofimova, M.E. & Moiseeva, N.S. (1989). Kinetic properties of the octameric and dimeric forms of mitochondrial creatine kinase and physiological role of the enzyme. *Biochem. Int.*, Vol. 19, No 3, pp. 603-613.
- Lyubarev, A.E. (1977). Origin of biochemical organization. *Biosystems*, Vol. 42, No 2-3, pp. 103-110.
- Magnani, M.; Serafini, G., Stocche, V., Bossù, M. & Dachà, M. Solubilization, purification and properties of rabbit brain hexokinase. *Arch. Biochem. and Biophys.*, Vol. 216, No 2, pp. 449-454.
- Mattson, M.P.; Liu, D. (2002). Energetics and oxidative stress in synaptic plasticity and neurodegenerative disorders. *Neuromolecular Med*, Vol. 2, No 2, pp. 215-231.


- Maulik, N.; Goswami, S., Galang, N. & Das, D.K. (1999). Differential regulation of Bcl-2, AP-1 and NF-kappaB on cardiomyocyte apoptosis during myocardial ischemic stress adaptation. *FEBS Lett.*, Vol. 443, No 3, pp. 331-336.
- Meyer, L.E.; Machado, L.B., Santiago, A.P., da-Silva, W.S., De Felice, F.G., Holub, O., Oliveira, M.F. & Galina, A. (2006). Mitochondrial creatine kinase activity prevents reactive oxygen species generation. Antioxidant role of mitochondrial kinase-dependent ADP re-cycling activity. *The journal of biological chemistry*, Vol. 281, No 49, pp. 37361-37371.
- Mulichak, A.M.; Wilson, J.E., Padmanabhan, K. & Garavito, R.M. (1998). The structure of mammalian hexokinase I. *Nat. Struct. Biol.*, Vol. 5, pp. 555-560.
- Ovadi, J. & Srere, P.A. (2000). Macromolecular compartmentation and channeling. *Intl. Rev. Cytol.*, Vol. 192, pp. 255-280.
- Perez-Pinzon, M.A.; Dave, K.R. & Raval, A.P. (2005). Role of reactive oxygen species and protein kinase C in ischemic tolerance in the brain. *Antioxid. Redox Signal*, Vol. 7, No 9-10, pp. 1150-1157.
- Rauchova, H.; Kaudelova, J., Drahota, Z. & Mourek J. (2002). Hypoxia-induced lipid peroxidation in rat brain and protective effect of carnitine and phosphocreatine. *Neurochem. Res.*, Vol. 27, No 9, pp. 899-904.
- Redker, V.D. & Kenkare, V.W. (1972). Bovine brain mitochondrial hexokinase. Solubilization, purification and role of sulphhydryl residues. *J. Biol. Chem.*, Vol. 247, pp. 1576-1584.
- Rosano, C. E.; Sabini, M. Rizzi, M., Deriu, D., Murshudov, G., Bianchi, M., Serafini, G., Magnani, M. & Bolognesi, M. (1999). Binding of non-catalytic ATP to human hexokinase I highlights the structural components for enzyme-membrane association control. *Structure*, Vol. 7, No 11, pp. 1427-1437.
- Rose, I. & Warms, J.V.B. (1967). Mitochondrial hexokinase. Release. Rebinding, and location. *J. Biol. Chem.*, Vol. 242, pp. 1635-1645.
- Sappey-Mariniere, D.; Hubesch, B., Matson, G.B. & Weiner, M.W. (2002). Decreased phosphorus metabolite concentrations and alkalosis in chronic cerebral infarction. *Radiology*, Vol. 182, pp. 29-34.
- Sas, K.; Robotka, H., Toldi, J. & Vécsei, L. (2007). Mitochondria, metabolic disturbances, oxidative stress and the kynurenine system, with focus on neurodegenerative disorders. *J. Neurol. Sci*, Vol. 257, No 1-2, pp. 221-239.
- Schlattner, U.; Dolder, M., Wallimann, N. & Tokarska-Schlattner, M. (2001). Mitochondrial creatine kinase and mitochondrial outer membrane porin show a direct interaction that is modulated by calcium. *J. Biol. Chem.*, Vol. 276, No 51, pp. 48027-48030.
- Schnyder, T.; Engel, A., Lustig, A. & Wallimann, T. (1988). Native mitochondrial creatine kinase forms octameric structures. II. Characterization of dimers and octamers by ultracentrifugation, direct mass measurements by scanning transmission electron microscopy, and image analysis of single mitochondrial creatine kinase octamers. *J. Biol. Chem*, Vol. 263, No 32, pp. 16954-16962.
- Schnyder, T.; Tittmann, P., Winkler, H., Gross, H. & Wallimann, T. (1995). Localization of reactive cysteine residues by maleidoyl undecagold in the mitochondrial creatine kinase octamer. *J. Struct. Biol*, Vol. 114, No 3, pp. 209-217.

- Schwab, D.A & Wilson, J.E. (1989). Complete amino acid sequence of rat brain hexokinase, deduced from the cloned cDNA, and proposed structure of a mammalian hexokinase. *Biochem.*, Vol. 86, pp. 2563-2567.
- Shen, W.; Willis, D., Zhang, Y., Schlattner, U., Wallimann, T. & Molloy, G.R. (1978). Expression of creatine kinase isozyme genes during postnatal development of rat brain cerebellum: Evidence for transcriptional regulation. *Biochem. J.*, Vol. 367, No 2, pp. 369-380.
- Siesjo, B.K. Brain energy metabolism / B.K. Siesjo. - Ed. J.Wiley and Sons, 1978. - 607p.
- Smith, A.D. & Wilson, J.E. (1991). Disposition of mitochondrially bound hexokinase at the membrane surface, deduced from reactivity with monoclonal anti bodies recognizing epitopes of defined location. *Arch. Biochem. Biophys.*, Vol. 287, pp. 359-366.
- Suzuki, Y.J.; Forman, H.J. & Sevanian, A. Oxidants as stimulators of signal transduction. *Free Radic. Biol. Med.*, Vol. 22, No 1-2, pp. 269-285.
- Takagi, Y.; Yasuhara, T. & Gomi K. (2001). Creatine kinase and its isozymes. *Rinsho Byori*, Suppl. 116, pp. 52-61.
- Ueda, M.; Muramatsu, H., Kamiya, T., Muramatsu, A., Mori, T., Terashi, A. & Katayama, Y. (2000). Pyruvate dehydrogenase activity and energy metabolite level following bilateral common carotid artery occlusion in rat brain. *Life Sci*, Vol. 67, No 7, pp. 821-826.
- Viitanen, P.V.; Geiger, P.J., Erickson-Viitanen, S. & Bessman, S.P. (1984). Evidence for functional hexokinase compartmentation in rat skeletal muscle mitochondria. *J. Biol. Chem.*, Vol. 259, No 15, pp. 9679-9686.
- Vyssokikh, M. & Brdiczka, D. (2003). The Function of complexes between the outer mitochondrial membrane pore (VDAC) and the adenine nucleotide translocase in regulation of energy metabolism and apoptosis. *Acta Biochimica Polonica*, Vol. 50, No 2, pp. 389-404.
- Wallimann, T. et al. (1992). Intracellular compartmentation, structure and function of creatine kinase isoenzymes in tissues with high and fluctuating energy demands: the "phosphocreatine circuit" for cellular energy homeostasis. *Biochem. J.*, Vol. 281, pp. 21-40.
- Wallimann, T. et al. (1998). Some new aspects of creatine kinase (CK): compartmentation, structure, function and regulation for cellular and mitochondrial bioenergetics and physiology. *Biofactors*, Vol. 8, No 3-4, pp. 229-234.
- Walzel, B.; Speer, O., Zanolla, E., et al. (2002). Novel mitochondrial creatine transport activity: implications for intracellular creatine compartments and bioenergetics. *J. Biol. Chem.*, Vol. 277, pp. 37503-37511.
- Wang, P.F.; Flynn, A.J., McLeish, M.J. & Kenyon, G.L. (2005). Loop movement and catalysis in creatine kinase. *IUBMB Life*, Vol. 57, No 4-5, pp. 355-362.
- Wilson, J.E. (1980). Brain hexokinase, the prototype ubiquitous enzyme. *Curr. Top. Cell. Regul.*, Vol. 16., pp. 2-54.
- Wilson, J.E. (1985). Regulation of mammalian hexokinase activity Regul. *Carbohydr. Metab.*, No 1, pp. 45-85.
- Wilson, J.E. (1995). Hexokinases. *Rev. Physiol. Biochem. Pharmacol.*, Vol. 126, pp. 65-198.

- Wilson, J.E. (2003). Isozymes of mammalian hexokinase: structure, subcellular localization and metabolic function. *The J. Exp. Biol.*, Vol. 206, pp. 2049-2057.
- Wyss, M. & Kaddurah-Daouk, R. (2000). Creatine and creatinine metabolism. *Physiol. Rev.*, Vol. 80, pp. 1107-1213.
- Xie, G. & Wilson, J.E. (1990). Tetrameric structure of mitochondrially bound rat brain hexokinase: a crosslinking study. *Arch. Biochem. Biophys.*, Vol. 276, No 1, pp. 285-293.

IntechOpen

IntechOpen



Advances in the Preclinical Study of Ischemic Stroke

Edited by Dr. Maurizio Balestrino

ISBN 978-953-51-0290-8

Hard cover, 530 pages

Publisher InTech

Published online 16, March, 2012

Published in print edition March, 2012

This book reports innovations in the preclinical study of stroke, including - novel tools and findings in animal models of stroke, - novel biochemical mechanisms through which ischemic damage may be both generated and limited, - novel pathways to neuroprotection. Although hypothermia has been so far the sole "neuroprotection" treatment that has survived the translation from preclinical to clinical studies, progress in both preclinical studies and in the design of clinical trials will hopefully provide more and better treatments for ischemic stroke. This book aims at providing the preclinical scientist with innovative knowledge and tools to investigate novel mechanisms of, and treatments for, ischemic brain damage.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Elena Erlykina and Tatiana Sergeeva (2012). Role of Creatine Kinase – Hexokinase Complex in the Migration of Adenine Nucleotides in Mitochondrial Dysfunction, *Advances in the Preclinical Study of Ischemic Stroke*, Dr. Maurizio Balestrino (Ed.), ISBN: 978-953-51-0290-8, InTech, Available from:
<http://www.intechopen.com/books/advances-in-the-preclinical-study-of-ischemic-stroke/role-of-creatine-kinase-hexokinase-complex-in-the-migration-of-adenine-nucleotides-under-mitochondri>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen