

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,800

Open access books available

142,000

International authors and editors

180M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Slit/Robo Signaling: Inhibition of Directional Leukocyte Migration

Ilya M. Mukovozov and Lisa A. Robinson
*The University of Toronto,
The Hospital for Sick Children,
Canada*

1. Introduction

Localized inflammation and the associated influx of leukocytes is a hallmark of the pathogenesis of many diseases. The ability to target the recruitment of leukocytes holds vast therapeutic potential in inflammatory diseases where there is excessive cell recruitment due to an overactive immune response, or the improper resolution of the initial response resulting in chronic leukocyte infiltration.

1.1 The neutrophil

Polymorphonuclear leukocytes, or neutrophils, are a critical component of the innate immune system, participating in host defence against bacterial and fungal infections. Not surprisingly, neutropenias can lead to severe infections and sepsis. During an inflammatory response, neutrophils are recruited to the sites of infection and/or injury by chemoattractants, including the chemokine family of proteins. Once in the tissue, neutrophils fight infections by ingesting microorganisms and producing reactive oxygen intermediates (ROI) as well as other antimicrobial substances, such as defensins (Ganz, T., 2003). Neutrophils can also produce and/or exacerbate inflammatory disease states as a result of the potent systems that have evolved in these cells for microbial killing. Inappropriate or excessive activation of these systems results in tissue damage (Fujishima et al., 1995). To better understand the role of the neutrophil in the fine balance between host defence and tissue injury, the mechanisms underlying neutrophil recruitment will be discussed.

1.2 Neutrophils and tissue injury

Neutrophils have been implicated in the pathogenesis of several inflammatory conditions, including: ischemia reperfusion injury (following coronary artery occlusion) (Frangogiannis et al., 2002), idiopathic pulmonary fibrosis (Haslam et al., 1980), arthritis (Weissmann et al., 1984), asthma (Lemanske et al., 1983), vasculitis (Fauci et al., 1978), glomerulonephritis (Holdsworth et al., 1984) and acute respiratory distress syndrome (ARDS) (Wieland et al., 1999). Neutrophil-mediated tissue injury results from the release of neutrophil antimicrobial factors such as ROI and proteases, and other mediators that amplify cell recruitment into the extracellular milieu (Frangogiannis et al., 2001). This can occur in two ways: 1) activation of neutrophils leads to fusion of antimicrobial granules to the plasma

membrane and subsequent release of granule contents, and 2) attempts to ingest large particles result in a large open vacuole, and subsequent granule fusion and release of granule contents into the extracellular space (Weissmann et al., 1971).

ROI are strong oxidizing and reducing agents that damage the integrity of cell membranes by lipid peroxidation (Li et al., 2002). ROI also promote arachidonic acid synthesis by activating phospholipase A2. Arachidonic acid is an important precursor of eicosanoids and prostaglandins, including thromboxane A₂ and leukotriene B₄ (Toyokuni et al., 1999). Increased production of these pro-inflammatory molecules enhances recruitment of leukocytes. ROI also induce activation of transcription factors such as nuclear factor κ B (NF- κ B) and activator protein1 (AP1) (Toyokuni et al., 1999), leading to increased expression of adhesion molecules, including P-selectin, and chemokines (such as IL-8) thereby facilitating leukocyte arrest and recruitment from the circulation (Eltzschig et al., 2004).

Activated neutrophils also secrete matrix metalloproteases (MMPs), including collagenase and gelatinase. These enzymes are structurally specialized to digest basement membranes and interstitial structural proteins to facilitate neutrophil extravasation and subsequent migration through the interstitium (Kang et al., 2001). MMPs degrade several major structural components of the extracellular matrix (ECM), including collagen, fibronectin, proteoglycans, laminin and gelatin. MMPs are antagonized by tissue inhibitors of metalloproteases (TIMPs; Own et al., 1999). It has been shown that the imbalance between TIMPs and neutrophil-derived MMPs is a key feature of inflammatory conditions, including ARDS and asthma (Cederqvist et al., 2001). Neutrophil derived elastase is another bactericidal protease that is also associated with tissue damage. Like the MMPs, elastase displays proteolytic activity against structural components of the ECM. Elevated levels of neutrophil derived elastase and collagenase have been detected in patients with chronic inflammatory conditions, such as rheumatoid arthritis (Garcia et al., 1987). Increased neutrophil-derived protease activity has also been linked to cartilage destruction (Mohr et al., 1981). In ARDS, elastase activity has been associated with degradation of surfactant proteins in the lung (Hirche et al., 2004; Rubio et al., 2004). These proteins increase bacterial opsonization and clearance of apoptotic neutrophils (Vandivier et al., 2002). Therefore, increased elastase activity could indirectly increase susceptibility to infection and delay resolution of inflammation in the lung.

Commonly prescribed anti-inflammatory drugs, such as aspirin and glucocorticoids, have shown some success in reducing neutrophil-mediated tissue damage. However, these drugs generally attenuate activation of transcription factors such as NF- κ B, thereby non-specifically reducing expression of cytokines and leukocyte adhesion molecules (Panes et al., 1999). One alternative method to prevent neutrophil-mediated tissue injury is to selectively block neutrophil recruitment to inflammatory foci. However, the redundancy in chemoattractant pathways means that interruption of a particular chemoattractant pathway may result in another pathway assuming its function. In principle, localized general chemoattractant blockade could be a useful strategy. Unique strategies to achieve this may be gained from studying central nervous system (CNS) development, in which positive and negative guidance cues for neuronal migration and axonal pathfinding have been defined.

1.3 Leukocyte trafficking and the adhesion cascade

The purpose of the inflammatory response is to selectively recruit the appropriate subsets of leukocytes to a site of inflammation. Inflammatory cytokines, such as interleukin 1 (IL-1)

and tumour necrosis factor α (TNF- α), and soluble chemoattractants, are released within the local inflammatory environment. This results in local vasodilation, increased volume of blood perfusing the inflamed area, and a simultaneous decrease in the flow velocity within the vessel, facilitating extravasation of circulating leukocytes. Leukocytes are recruited to sites of inflammation in a series of coordinated interactions with endothelial cells lining the vascular wall. The classical leukocyte adhesion cascade involves these main steps: i) leukocyte capture and rolling, ii) activation and arrest, and iii) transendothelial migration (Fig. 1). Failure in any one of these steps can result in severe immunodeficiencies (Beutler, B., 2004). However, there exists a substantial therapeutic potential for the localized blockade of leukocyte adhesion and diapedesis.




Fig. 1. Endothelial-leukocyte interactions leading to transmigration across the vascular wall. (i) Capture and (ii)rolling: The initial tethering of leukocytes to the endothelial cells lining the vessel wall is mediated by the selectins. These structural interactions enable the leukocyte to roll along the venular wall and to 'sample' the endothelial surface for activating factors (iii) Arrest: These interactions lead to leukocyte integrin activation. Firm adhesion of the leukocyte is mediated through binding of integrins to members of the immunoglobulin superfamily expressed in stimulated endothelial cells.(iv) Diapedesis: Following firm adhesion, the cell changes shape in response to local chemoattractant gradients and transmigrates across the endothelial barrier.

Selectins are a family of adhesion molecules that are structurally specialized for the initial capture of circulating leukocytes. Rolling is mediated by E-selectin and P-selectin, expressed by endothelial cells, and by L-selectin expressed on the majority of circulating neutrophils, monocytes, eosinophils, and T and B lymphocytes (Kansas, G., 1996). The broad expression pattern of L-selectin allows for nonspecific recruitment of all leukocyte lineages. P-selectin is constitutively found in Weibel-Palade bodies of endothelial cells, and mobilized to the cell surface within minutes following activation by inflammatory mediators (Frangogiannis et al., 2002). All of the selectins interact with P-selectin glycoprotein ligand 1 (PSGL1), although other glycoprotein ligands exist, such as CD34 and MadCAM-1 (McEver et al., 1997; Puri et al., 1995). Following initial leukocyte capture, the binding of leukocyte L-selectin to PSGL1 facilitates secondary leukocyte capture, where adherent leukocytes assist

in the recruitment of additional cells (Eriksson et al., 2001). Interactions of selectins with their ligands allow leukocytes to roll on inflamed endothelium under the rapid flow of the bloodstream (Alon et al., 1995). In fact, shear stress is required to support L-selectin and P-selectin dependent adhesion, and rolling cells detach when flow is stopped (Finger et al., 1996; Lawrence et al., 1997). This selectin-mediated slow rolling allows the leukocyte to 'sample' the repertoire of chemokines and other activation signals presented on the luminal surface of endothelial cells.

In addition to selectins, various integrins participate in rolling. Integrins bind members of the immunoglobulin superfamily, including vascular cell-adhesion molecule 1 (VCAM-1) and intercellular adhesion molecule 1 (ICAM-1). Neutrophils roll on immobilized VCAM-1 by engaging the leukocyte integrin receptor, very late antigen 4 (VLA-4; $\alpha_4\beta_1$ -integrin). β_2 -integrins also support rolling (Sigal et al., 2000). Resting mouse neutrophils roll on surfaces coated with E-selectin ligand and ICAM-1. Ligation of endothelial E-selectin induces a structural conformational change in leukocyte lymphocyte function-associated antigen 1 (LFA-1; $\alpha_L\beta_2$ -integrin) allowing it to bind to its endothelial ligand, ICAM-1 (Salas et al., 2004). In addition, it has recently been demonstrated that the mechanochemical design of LFA-1 allows shear stress to induce and maintain a state of high ligand-binding affinity (Astrof et al., 2006). Rolling *in vivo* requires E-selectin (Kunkel et al., 1996), engagement of the β_2 -integrins (Jung et al., 1998), LFA-1 and macrophage antigen-1 (MAC1; Dunne et al., 2002).

Although leukocytes (particularly neutrophils) roll under normal conditions, during inflammation leukocytes undergo integrin-dependent arrest. Arrest of leukocytes on endothelial cells is rapidly triggered by the binding of chemokines and other chemoattractants (Campbell et al., 1998). These chemoattractants are secreted by activated endothelial cells and platelets. In fact, platelets can deposit chemokines, such as CC-chemokine ligand 5 (CCL5), CXC-chemokine ligand 4 (CXCL4), and CXCL5 onto the inflamed endothelial lumen to trigger leukocyte arrest (von Hundelshausen et al., 2001; Huo et al., 2003).

Following firm arrest, leukocytes migrate, by a process called diapedesis, across the endothelial cell barrier, its associated basement membrane, and the pericyte sheath. Leukocyte diapedesis and chemotaxis is triggered by chemokines (such as IL-8) presented to rolling leukocytes on the luminal surface of endothelial cells. Leukocytes can cross the endothelium between adjacent endothelial cells (paracellular route) or directly through an endothelial cell (transcellular route). Transcellular migration generally occurs in 'thin' parts of the endothelium where there is less distance for the leukocyte to migrate (Ley et al., 2007). In addition, caveolae containing ICAM-1 link together to form vesiculo-vacuolar organelles (VVOs), providing shortcuts for transcellular leukocyte diapedesis (Dvorak et al., 2001). This creates a channel inside the cell through which leukocytes can migrate. During paracellular migration, ligation of endothelial-cell adhesion molecules results in reduced interendothelial contacts, facilitating the migration of leukocytes through endothelial cell junctions (Ley et al., 2007). Transendothelial migration requires an increase in intracellular endothelial calcium, which promotes opening of endothelial cell junctions via the activation of myosin light chain kinase and endothelial cell contraction. The route of leukocyte migration is determined by both the surface density of ICAM-1 and the shape of endothelial cells (Yang et al., 2005). Both a high density of ICAM-1 and endothelial cells with a polygonal morphology promote transcellular migration (Yang et al., 2005). Many endothelial junctional molecules, such as platelet/endothelial-cell adhesion molecule 1 (PECAM-1), ICAM -1,

ICAM-2, junctional adhesion molecule A (JAM-A), (JAM-B), (JAM-C), endothelial cell-selective adhesion molecule (ESAM), and CD99, play a role in leukocyte transmigration. Although the leukocyte adhesion cascade has been divided into several steps, these are not temporally exclusive, but instead synergistically promote leukocyte arrest and diapedesis. Leukocyte diapedesis was described almost 200 years ago, but its molecular mechanisms are only now beginning to be more fully understood (Imhof et al., 2004). In the past decade, new insights have been gained into the signaling events that underlie integrin activation, post-adhesion strengthening of leukocyte attachment and the structural significance of molecules involved in diapedesis (Muller, W., 2003).

1.4 Chemotaxis

Following extravasation, leukocytes migrate through the interstitial ECM, following a chemoattractant gradient, to reach the site of inflammation. Chemotaxis, directed cell migration towards external chemical gradients, occurs in many eukaryotic cells including: free-living organisms, leukocytes (during inflammation), endothelial cells (angiogenesis), spermatocytes (fertilization) and neurons (neurogenesis) (Singer et al., 1986). Upon exposure to a chemoattractant the cell orients itself in the direction of locomotion along the chemoattractant gradient. Polarization results from preferential pseudopod extension towards areas of higher chemoattractant concentration (Zigmond, S., 1974). Efficient chemotaxis requires coordination between pseudopod formation at the leading edge of the cell, and uropod retraction at the trailing edge. During chemotaxis, neutrophils extend short surface protrusions called filopodia, or microspikes, which are membrane extensions of approximately 0.1-0.2 μm in diameter and up to 20 μm in length. These structures act as cellular tentacles and are supported by a core bundle of actin microfilaments (Mattila et al., 2008). In neutrophils, filopodia support thin sheets of membrane-enclosed cytoplasm, called lamellipodia. Lamellipodia contain actin filaments and a meshwork of myosin II-associated microfilaments. In neutrophils, the actin network within the lamellipodia, together with other structural and regulatory proteins, comprises the molecular motor which drives cell locomotion (Jones et al., 1998). This locomotory apparatus works against cell-to-substratum adhesions called focal contacts or focal adhesions. Focal adhesions are molecular structures that utilize integrins to link the myosin II-containing bundles of cytoplasmic microfilaments (called stress fibers) to proteins in the extracellular matrix (ECM) (Critchley et al., 1999). In neutrophils, integrin-mediated contacts to the ECM take two forms: focal complexes and podosomes. Focal complexes are structurally similar to focal adhesions but lack stress fibers (Allen et al., 1997), while podosomes are distinct circular structures that are only observed in cells of the myeloid lineage (DeFife et al., 1999; Correia et al., 1999; Linder et al., 2003). In this way, cytoskeletal rearrangement permits leukocytes to migrate toward chemoattractant gradients.

1.5 Chemoattractants

Many types of chemoattractant recruit leukocytes to inflammatory foci. These include bacterial components, leukotrienes, complement factors and chemokines. C5a, the first chemoattractant identified, is a cleaved product derived from complement component C5 (Shin et al., 1968). Bacterial products such as fMLP (N-formyl-methionyl-leucyl-phenylalanine) and other N-formylpeptides also act as chemoattractants that non-specifically recruit leukocyte subsets to inflammatory foci. An important family of

chemoattractants involved in leukocyte recruitment to inflammatory foci is a family of chemoattractant cytokines called chemokines. Chemokines constitute a large family of small peptides that are structurally similar and that bind to a family of seven transmembrane G-protein coupled receptors (Rossi et al., 2000). The specific expression, regulation, and receptor binding patterns of each chemokine determine their functional diversity. Most chemokines are structurally conserved to bind to glycosaminoglycans (GAGs) on the luminal surface of endothelial cells. This binding is required for leukocyte recruitment *in vivo*. Indeed, chemokines with mutations in their GAG binding domains can induce *in vitro* chemotaxis, but are unable to recruit leukocytes to the peritoneal cavity *in vivo* (Johnson et al., 2005).

The binding of chemoattractants to their receptors activates leukocyte integrins instantaneously by inside-out signalling mechanisms (Shamri et al., 2005). They rapidly regulate integrin avidity by increasing both integrin affinity (by a conformational change that results in increased ligand binding energy and a decreased ligand dissociation rate), and valency (the density of integrins per area of plasma membrane involved in adhesion, determined by expression levels and lateral mobility) (Laudanna et al., 2002; Constantin et al., 2000). Through these signaling mechanisms, chemokines work as powerful activators of integrin-mediated adhesion and leukocyte recruitment.

1.6 Intracellular signaling of chemoattractant receptors

Several neutrophil chemoattractants, particularly chemokines, interact with specific receptors on the plasma membrane, transducing signals by coupling to heterotrimeric G proteins. Heterotrimeric G proteins are composed of an α , β , and γ subunit. The α subunit is the GDP/GTP binding element. When bound to GDP, the α subunit interacts with the β and γ subunits to form an inactive heterotrimer complex. Chemoattractant binding induces a conformational change in the receptor, exchanging GDP for GTP on the α subunit. The α subunit then dissociates from the receptor, releasing the $G\beta\gamma$ complex. The free $G\alpha$ and $G\beta\gamma$ subunits are then available to bind and activate target enzymes such as phosphatidylinositol 3-kinase (PI3K), phospholipase C (PLC), or adenylyl cyclase (Fig. 2). These enzymes generate secondary intracellular messengers that initiate a cascade of signaling events that ultimately culminate in cytoskeletal rearrangement and leukocyte migration.

Ligation of chemoattractant receptors leads to the activation of four major signaling pathways (Fig. 2): PLC, PI3K, mitogen-activated protein kinases (MAPKs) and Rho guanosine triphosphatases (GTPases). Once the $G\alpha$ subunit dissociates, the $G\beta\gamma$ complex activates PLC, which cleaves phosphatidylinositol (4,5)-bisphosphate (PI(4,5)P₂) to generate inositol (1,4,5)-triphosphate (IP₃) and diacylglycerol (DAG). Generation of IP₃ leads to the mobilization of intracellular calcium stores from the endoplasmic reticulum, and together with DAG, activates protein kinase C (PKC) (Li et al., 2000). The activation and recruitment of PKC to the plasma membrane promotes changes in the actin cytoskeleton that facilitate and/or drive cell spreading and migration (Fig. 2).

A convincing role for PI3K in chemoattractant receptor signaling and chemotaxis has been established (Li et al., 2000; Sasaki et al., 2000; Hirsch et al., 2000; Servant et al., 2000; Jin et al., 2000). Although there are at least four Class I PI3K isoforms in mammalian cells (Vanhaesebroeck et al., 1999), only a single Class IB variant has been shown to interact with

chemoattractant receptors in leukocytes. The outcome of Class I PI3K activation is phosphorylation of membrane PI(4,5)P₂ by activated PI3K, generating PI(3,4,5)P₃ at the plasmalemma. The Gβγ complex also activates PI3Kγ, activating Src-family kinases and generating PI(3,4,5)P₃ from membrane PI(4,5)P₂ (Krugmann et al., 1999), resulting in the recruitment of Ras GTPases and subsequent activation of MAPK pathways (Fig. 2) (Kintscher et al., 2000). Although MAPK signaling pathways are involved in chemotaxis and adhesion, the most important biochemical events for cell polarization are the production of PIP₃ and activation of Rho GTPases at the leading edge of the cell.




Fig. 2. Intracellular signaling cascade upon ligation of chemoattractant receptors. Chemoattractant binding to GPCRs induces a conformational change that results in the dissociation of Gα subunits from the Gβγ complex. This leads to rapid outside-in signaling resulting in the activation of four major signaling pathways that contribute to the generation of cell polarity and chemotaxis: Rho GTPases, PI3K, PLC, and MAPKs.

The PI3K dependent production of PIP₃ at the cell membrane allows for the recruitment of the Rho-family GTPases, Rac and Cdc42, to the cell membrane. The localization of PIP₃, Rac and Cdc42 then stimulate polymerization of actin, a process necessary for the formation of

filopodia and lamellipodia at the front of the cell. At the back of the cell, Rho-kinase phosphorylation results in inactivation of myosin light chain phosphatase, leading to increased myosin light-chain kinase (MLK) dependent activation of myosin (Nguyen et al., 1999). These biochemical conditions favour the formation of actomyosin bundles, contraction, de-adhesion from the substratum and tail retraction (Ridley, A., 2001; Bokoch, G., 2005). Interestingly, signals at the leading edge inhibit signals at the trailing edge, allowing for the maintenance of cell polarity (Fenteany et al., 2004). To prevent the accumulation of PIP₃ at the trailing edge, PTEN dephosphorylates PI(3,4,5)P₃ to PI(4,5)P₂. The lack of PIP₃ in the back of the cell prevents activation and recruitment of Rho GTPases and subsequent actin polymerization, allowing the formation of actomyosin bundles and tail retraction (Worthylake et al., 2001). Actin polymerization at the leading edge coupled with tail retraction in the back allows for directed leukocyte chemotaxis.

1.7 Rho-family GTPases: Rac, Cdc42, and Rho

Small GTPases of the Rho family are a part of the Ras superfamily of small GTP-binding proteins. They are pivotal regulators of many signaling networks that are activated by a diverse variety of receptor types. To date, over 20 mammalian Rho GTPases have been characterized, and these can be grouped into 6 different classes: Rac (Rac1, Rac2, Rac3, RhoG), Rho (RhoA, RhoB, RhoC), Cdc42 (Cdc42Hs, G25K, TC10), Rnd (Rnd3/RhoE, Rnd1/Rho6, Rnd2/Rho7), RhoD, and RhoH/translocation three four (TTF) (Aspenström, P., 1999; Kjoller et al., 1999). When activated, Rho GTPases regulate many important processes in all eukaryotic cells, including actin cytoskeleton dynamics, transcription, cell cycle progression, and membrane trafficking. The activity of Rho GTPases is regulated by outside-in signals from a variety of receptor types, including GPCR, tyrosine kinase receptors, cytokine receptors and adhesion receptors. Rho-family GTPases play a critical role in regulating leukocyte chemotaxis, adhesion and phagocytosis.

1.7.1 Rho GTPases: Structure and regulation

All Rho GTPases contain two main structural domains, the C-terminal 'CAAX' motif and a catalytic GTP domain. The 'CAAX' motif undergoes post-translational processing, involving carboxy-terminal proteolysis of the AAX residues followed by carboxyl-methylation. The modified C-terminal domain can then attach to membrane lipids and facilitates membrane association and subcellular localization of Rho GTPases (Gutierrez et al., 1989; Casey et al., 1989; Fujiyama et al., 1990). The catalytic domain contains two regions, switch I and switch II. These domains correspond to different structural conformations in the GTP-bound and GDP-bound forms. Rho GTPases function as molecular switches by cycling between GDP-bound and GTP-bound forms. When bound to GDP, Rho GTPases are inactive. Binding of ligands to cell surface receptors, results in exchange of GDP for GTP, switching the protein to an active state. The active form interacts with downstream effector molecules. The intrinsic GTPase activity of Rho GTPases completes this cycle by hydrolyzing GTP, returning the GTPase to its inactive GDP-bound state.

Three classes of molecules interact with Rho GTPases and regulate their activation state: guanine nucleotide exchange factors (GEFs), GTPase-activating proteins (GAPs), and

guanine nucleotide dissociation inhibitors (GDIs). GEFs catalyze the exchange of GDP for GTP, leading to the activation of Rho GTPases. To date, over 69 mammalian GEFs for Rho GTPases have been identified (Rossman et al., 2005). They are characterized by the presence of a Dbl homology domain (DH), which interacts with both the switch I and switch II regions and catalyses the exchange of GDP for GTP. In addition, many of these DH-domain containing proteins, such as Vav, contain a Pleckstrin homology (PH) domain which allows GEFs to bind phosphoinositides, such as PIP₃. This localizes GEFs to the plasma membrane where they can bind other Rho-family GTPase-interacting proteins. GAPs enhance the intrinsic GTPase activity of Rho GTPases, and thus suppress their activity. Although GTPases possess intrinsic GTPase activity, the actual rate of GTP hydrolysis is relatively slow. Therefore, the interaction with a GAP is required for efficient GTP hydrolysis, as this accelerates the cleavage step by several orders of magnitude (Vetter et al., 2001). To date, more than 70 eukaryotic RhoGAPs have been discovered, of which 35 are found in humans (Tcherkezian et al., 2007). There exists a large diversity in the primary sequences of the various GAPs. However, each one contains a Rho GAP domain with a conserved tertiary structure composed of α helices and a catalytically critical 'arginine finger' which stabilizes the formation of the transition state during GTP hydrolysis (Nassar et al., 1998). In addition, the Rho GAP domain interacts with both the switch I and switch II regions on the GTPase domain (Gamblin et al., 1998), allowing GAPs to facilitate the intrinsic hydrolysis of GTP, resulting in the inactivation of Rho GTPases.

Finally, GDIs associate with Rho GTPases in their inactive GDP-bound state and inhibit their activation by GEFs. GDIs also bind to GTP-bound GTPases, and suppress their activity (Oloffson, B., 1999). There is evidence that GDIs can bind to isoprenyl moieties on the C-terminus of GTPases in order to sequester them in the cytosol (Keep et al., 1997). The role of GDIs in partitioning GTPases between the membrane and cytosol may be physiologically more important than the inhibition of their activation, as this may provide a storage pool of Rho GTPases that is readily utilized upon cell activation. Overall, GDIs prevent the activation of Rho GTPases, prevent their interaction with membranes, and inhibit downstream signaling networks.

1.7.2 Rho GTPases and the actin cytoskeleton

The movement of eukaryotic cells relies on the coordinated extension of actin-rich lamellipodia in the leading edge and retraction of the uropod at the rear of the cell. The extension of lamellae in the leading edge involves rapid turnover of actin filaments (Symons et al., 1991; Wang, Y., 1985). More stable actin-myosin cables can be found in more established protrusions and in the middle and rear of the cell (DeBiasio et al., 1988). Recycling of the plasma membrane and integrin-mediated adhesion to the substratum and/or ECM are also important for cell motility (Bretscher, M., 1996; Martenson et al., 1993; Yamada et al., 1995; Mitra et al., 2005). Coordinated mobilization of the actin cytoskeleton is regulated by deployment of actin-binding proteins by activated Rho-family GTPases. Rho-family GTPases control cell motility and morphological changes in response to extracellular chemoattractants. Activation of Rho in fibroblasts results in the assembly of stress fibers and focal adhesions (Ridley et al., 1992). The activation of Rac causes extension of lamellipodia and assembly of small focal complexes (Nobes et al., 1995; Ridley et al., 1992). In contrast,

activation of the Cdc42 Rho-family GTPase leads to the formation of filopodial extensions (Nobes et al., 1995).

As discussed above, the influx of neutrophils and other leukocytes to inflammatory foci relies on activation of Rho-family GTPases and dynamic actin turnover. In principle, one method to prevent neutrophil-mediated tissue injury would involve blocking neutrophil recruitment. However, the redundancy in chemoattractant pathways means that interruption of a particular chemoattractant may result in another assuming its function. Thus, a localized general chemoattractant blockade could be a useful strategy. Unique strategies to target neutrophil recruitment may be gained from studying central nervous system (CNS) development, in which structurally distinct positive and negative guidance cues for migration and axonal pathfinding have been defined.

2. Slit2: A guidance cue for cell migration

During the development of the CNS, neurons must migrate and project axons over long distances. Most axons emanating from the CNS must cross the midline and then project longitudinally towards their synaptic targets. The molecular mechanisms that guide this pathfinding include contact attraction, chemoattraction, contact repulsion and chemorepulsion. Guidance cues selectively promote or repress migration of neurons and axonal projection. For example, netrins are diffusible chemotropic factors that attract commissural axons to the midline (Kennedy et al., 1994). The Slit family of secreted proteins, together with their cell-surface receptor Roundabout (Robo), repel neurons during CNS development. Once commissural axons have crossed the midline, midline glial cells express Slit to prevent axons from re-crossing the midline. Mutant *Drosophila* lacking Slit proteins exhibit midline defects, such as collapse of the regular scaffold of commissural and longitudinal axon tracts in the embryonic CNS (Rothberg et al., 1988; Rothberg et al., 1990). A similar defect is observed in mutant *Drosophila* lacking Robo, where projecting axon tracts cross the midline repeatedly (Kidd et al., 1998).


2.1 Slit and robo: Structure

The Slit family of proteins contain an N-terminal signal peptide, four leucine-rich repeats (LRRs), nine epidermal growth factor (EGF) repeats and a C-terminal cysteine knot (Fig. 3) (Rothberg et al., 1988; Rothberg et al., 1990; Rothberg et al., 1992). The EGF repeats and LRR allow Slit proteins to interact with ECM components, such as glypican-1, enabling them to act as localized, non-diffusible, signaling molecules (Ronca et al., 2001). Furthermore, Slit2 can be proteolytically cleaved after the fifth EGF repeat to form N-terminal (Slit2-N) and C-terminal (Slit2-C) fragments (Brose et al., 1999; Wang et al., 1999). Slit2-N includes the first 1118 amino acids and contains the four LRRs and the first five EGF repeats, while Slit2-C contains the remaining residues (Brose et al., 1999). Importantly, only the second LRR of human Slit2 is required to bind with the first Ig domain of Robo and initiate downstream signaling (Morlot et al., 2007). Therefore, both full length Slit2 and Slit2-N bind Robo receptors to repel migrating cells and projecting axons (Nguyen Ba-Charvet et al., 2001). Although the cleavage of Slit2 does not eliminate its activity, it may play a role in its diffusion since Slit-N appears to be more tightly associated with the cell membrane. In rat neural tissue both Slit2-N and Slit2-C were shown to bind heparan sulfate proteoglycan

glypican-1 (Liang et al., 1999), although Slit2-C bound with higher affinity, suggesting a possible regulatory mechanism for its diffusion.

Robo, a member of the immunoglobulin superfamily, is a single-pass type-1 receptor for the Slit proteins. The extracellular region of human Robo-1 contains five immunoglobulin (Ig) repeats and three fibronectin type III domains. The cytoplasmic region of Robo-1 contains four conserved cytoplasmic signaling motifs, CC0, CC1, CC2 and CC3 (Kidd et al., 1998; Zallen et al., 1998). Only the first Ig domain of Robo is required to bind to the second LRR domain in Slit2 and Slit2-N (Battye et al., 2001; Chen et al., 2001; Nguyen Ba-Charvet et al., 2001). The cytoplasmic CC motifs of Robo are required for response to Slit (Bashaw et al., 2000).

Slit2



Robo-1




Fig. 3. Primary Protein Structure of Mammalian Slit2 and Robo-1 Proteins. Mammalian Slit2 contains four leucine rich repeats (LRRs), nine epidermal growth factor (EGF) repeats, a laminin G (G) domain, and a cysteine rich C terminus. The Robo-1 receptor contains five immunoglobulin (Ig) repeats, three fibronectin (FN) type III, a transmembrane Domain (TM) and four conserved cytoplasmic (CC) signaling motifs.

The detection of an amino-terminal fragment of Robo-1 (Robo-1-NTF) in the conditioned medium of cancer cell lines and in the serum of patients with hepatocellular carcinoma suggests that Robo-1 may undergo proteolytic cleavage (Ito et al., 2006). The cleavage site was recently shown to be between Glu852 and Glu853, only 10 residues away from the plane of the plasma membrane (Seki et al., 2010). Following cleavage of transmembrane Robo-1 by MMPs, a soluble Robo-1-NTF is generated. The remaining carboxy-terminal fragment (Robo-1-CTF1) is subsequently cleaved by γ -secretase to form Robo-1-CTF2 (Fig. 4). Robo-1-CTF2 translocates to the nucleus, although its function is unknown (Seki et al., 2010).




Fig. 4. Successive cleavage of the Robo-1 receptor. Full-length Robo-1 is first cleaved by MMPs to form Robo-1-NTF and Robo-1-CTF1. The second cleavage, mediated by γ -secretase, releases Robo-1-CTF2 which translocates to the nucleus. The function of Robo-1-CTF2 at this location is unknown.

2.2 Slit and robo: Expression

Expression of the Slit genes has been demonstrated in many organisms, including *Drosophila* (Battye et al., 1999), *Caenorhabditis elegans* (Hao et al., 2001), *Xenopus* (Chen et al., 2000), *Gallus gallus domesticus* (Holmes et al., 2001; Vargesson et al., 2001), mice (Holmes et al., 1998; Piper et al., 2000), rats (Marillat et al., 2002) and humans (Itoh et al., 1998). In mammals there are three members of the Slit family. Although Slit1 is predominantly expressed in the developing CNS (Yuan et al., 1999), Slit2 and Slit3 are expressed outside the CNS, particularly in lung, kidney, and heart (Wu et al., 2001). Importantly, Slit expression persists in the adult organism, suggesting a role for Slit proteins beyond embryogenesis.

Expression of Robo has been demonstrated in *Drosophila* (Kidd et al., 1998), mice (Yuan et al., 1999) and humans (Kidd et al., 1998). There are four isoforms of Robo in mammals. Robo-1 is most highly expressed in tissues outside the CNS, including human leukocytes (Wu et al., 2001). Robo-2 is expressed during vertebrate limb development (Vargesson et al., 2001). Robo-3 is expressed following cerebellar and spinal cord lesions (Wehrle et al., 2005). Robo-4 is expressed in the adult organism by primary human endothelial cells, including

umbilical vein endothelial cells and microvascular endothelial cells (Suchting et al., 2005). Interestingly, the tissue expression of Slit and Robo is relatively complementary, suggesting a synergistic relationship (Yuan et al., 1999).

2.3 Slit and robo: Function

Recent studies demonstrate a role for Slit and Robo as repellents outside the CNS. For example, in mesoderm migration in *Drosophila*, myocyte precursors migrate away from the midline towards peripheral target sites where they fuse to form muscle fibers. In Slit and Robo mutants, these cells do not migrate away from the midline and instead fuse across it (Rothberg et al., 1990). Interestingly, this defect can be reversed by expressing Slit protein in midline cells (Kramer et al., 2001). Slit and Robo signaling also plays a role in nephrogenesis. During renal development, formation of a ureteric bud requires secretion of glial cell derived neurotrophic factor (GDNF) by nearby mesenchymal cells. Slit2 and Robo-2 knockout mice display abnormal patterns of GDNF secretion and develop multiple ureteric buds and multiple urinary collecting systems (Ray, L., 2004). Furthermore, polymorphisms in the human *Robo2* gene are associated with familial vesicoureteral reflux (Bertoli-Avella et al., 2008), a condition involving improper insertion of ureters into the bladder resulting in retrograde flow of urine from the bladder to the kidney. Slit2 also acts as a repellent in the mature organism. A recent study demonstrated that Slit2 inhibits vascular smooth muscle cell migration toward a gradient of platelet-derived growth factor (PDGF) (Liu et al., 2006). This inhibition occurred by suppression of activation of the small GTPase, Rac1. Slit2 has been shown to prevent cancer cell metastasis. The chemokine receptor, CXCR4, is expressed by some human breast cancer cells, allowing them to migrate towards gradients of the CXCR4 ligand, stromal cell-derived factor-1 (SDF-1 α), and promoting their metastasis to the lung. Slit2 inhibited chemotaxis, adhesion and chemoinvasion of these breast cancer cells (Prasad et al., 2004). Several other studies have demonstrated a role for Slit2 as a tumor suppressor. Slit2 was shown to inhibit colony formation in lung, colorectal and breast cancer cell lines (Dallol et al., 2002). *Slit2* has also been shown to be epigenetically silenced in more aggressive forms of these and other cancers (Dallol et al., 2003; Dallol et al., 2003; Dickinson et al., 2004). Collectively, these studies demonstrate a repellent role for Slit and Robo in the adult organism and in cancer biology.

The role of Robo-4 signaling in endothelial cells is controversial. Kaur et al. (2006) showed that Robo-4 signaling mediates attractive guidance mechanisms by activating Cdc42 and Rac1 in endothelial cells and inducing actin-mediated cell protrusions, including filopodia and lamellipodia. In fact, Robo-4-induced phenotypic effects in endothelial cells are rescued by dominant negative constructs of Cdc42. Thus, Robo-4 may mediate attractive signaling via activation of Rho-family GTPases, Cdc42 and Rac1. However, in 2008, Jones et al. showed that Slit inhibits endothelial cell migration and angiogenesis. In fact, Robo-4 signaling was shown to stabilize endothelial cell barriers (Jones et al., 2009). Thus, the precise role of Slit/Robo signaling in endothelial cells is yet to be determined.

2.4 Slit2/Robo-1 intracellular signal transduction

Studies of neuronal tissue have demonstrated that Robo-1 signals through two pathways that lead to remodeling of the cytoskeleton: Enabled (Ena) protein and Rho GTPases. Both of

these pathways require the CC motifs in the cytoplasmic domain of Robo. Ena and its mammalian homologue (Mena) are members of a family of proteins that link signal transduction to localized remodeling of the actin cytoskeleton by binding to profilin, an actin binding protein which regulates actin polymerization (Lanier et al., 1999; Wills et al., 1999). Ena is a substrate for Abelson kinase (Gertler et al., 1989). Ena and Abelson both bind to Robo. Ena binds to the CC1 motifs while Abelson binds to the CC3 motif (Fig. 5) (Bashaw et al., 2000). Impairing Ena binding reduces Robo function, while mutations in Abelson result in Robo hyperactivity (Bashaw et al., 2000).




Fig. 5. Intracellular signaling downstream of the Robo-1 receptor. Enabled protein bind to Robo-1 and may contribute to Slit-mediated repulsion. Abelson kinase phosphorylates intracellular domains of Robo and antagonizes Robo function. Ligation of Robo-1 by Slit2 results in the recruitment of srGAPs to the plasma membrane. srGAPs convert active GTP-bound forms of Cdc42 and Rac to their inactive, GDP-bound counterparts, thereby inhibiting the dynamic actin polymerization required for chemotaxis and preventing cell migration.

Slit/Robo also mediate cell repulsion through modulation of Rho GTPase activity. A family of GTPase activating proteins, Slit Robo GTPase activating proteins (srGAPs), were shown to bind Robo (Fig. 5) (Wong et al., 2001). The SH3 domain of srGAP binds the CC3 motif of Robo, while the GAP domain has activity for the Rho GTPases, Rac, Cdc42 and Rho (Wong et al., 2001). Ligation of Robo by Slit induces the recruitment of srGAP, thereby inactivating Rho-family GTPases and inhibiting actin remodeling and cell motility (Wong et al., 2001).

2.5 Slit/Robo in cell trafficking

Both neuronal and leukocyte chemotaxis require recognition of guidance cues, polarization of the cell, and mobilization of the actin cytoskeleton. In addition to repelling developing axons, Slit2 also inhibits chemotaxis of other cell types including vascular smooth muscle cells (Liu et al., 2006). However, the first study to demonstrate that Slit2 inhibits leukocyte chemotaxis, in 2001, utilized transwell migration assays to show that Slit2 inhibits chemotaxis of rat lymph node cells and neutrophil-like HL-60 cells towards MCP-1 and fMLP respectively (Wu et al., 2001). Subsequently, Kanellis et al. (2004) demonstrated that Slit2 inhibits chemotaxis of rat-derived macrophages towards MCP-1 and fMLP. Another study showed that Slit2 inhibited migration of dendritic cells (DCs) (Guan et al., 2003). In 2007, Prasad et al. demonstrated that Slit2 inhibits chemotaxis and transendothelial migration of primary CD4⁺ T lymphocytes toward SDF-1. Recently, Slit2 was shown to promote chemotaxis of eosinophils towards the chemokine, eotaxin, and to exacerbate allergic airway inflammation (Ye et al., 2010). Thus, Slit2 can negatively or positively regulate directional migration of individual leukocyte subsets.

3. Slit2/Robo-1 signaling inhibits neutrophil migration

Using immunoblotting, we previously demonstrated Robo-1 protein in human and mouse neutrophils (Tole et al., 2009). Immunofluorescence microscopy and flow cytometry revealed that Robo-1 was on the surface of cells.

We used Transwell migration assays to study the effects of Slit2 on chemotaxis of primary human neutrophils. In the presence of Slit2, fMLP-induced migration of neutrophils was inhibited in a dose-dependent manner. In fact, we observed that Slit2 is a potent inhibitor of neutrophil migration toward diverse types of chemotactic cues, including IL-8 and C5a (Tole et al., 2009).

Neutrophil exposure to chemoattractants results in the activation of the Rho GTPases, Rac and Cdc42 and the subsequent reorganization of actin filaments. (Sun et al., 2004; Srinivasan et al., 2003). Since the predominant isoform of Rac in human neutrophils is Rac2, not Rac1, the activation of Rac2 was studied. Following stimulation with fMLP, levels of activated Cdc42 and Rac2 in the presence of Slit2 were less than half of those observed in untreated control cells. We found that Slit2 inhibits neutrophil chemotaxis and actin polymerization by preventing cell polarization and disrupting generation and recruitment of activated Rac2 and Cdc42.

We examined the effects of Slit2 on the activation of kinase signaling pathways associated with neutrophil chemotaxis, namely, PI3K, Akt, Erk, and p38 MAPK. Stimulation of neutrophils with fMLP resulted in levels of activated Akt that were comparable in the presence or absence of Slit2, indicating that Slit2 does not impair the ability of neutrophils to

generate membrane PIP_3 . Similarly, Slit2 treatment had no effect on fMLP-induced phosphorylation of Erk and p38 MAPK. Thus, Slit2 inhibits neutrophil chemotaxis by specifically preventing activation of Cdc42 and Rac2 but not activation of Akt, Erk, or p38 MAPK (Tole et al., 2009).

We studied the effect of Slit2 on neutrophil recruitment *in vivo* using a mouse model of chemical irritant peritonitis (Glogauer et al., 2003). The administration of Slit2 prior to induction of peritonitis with sodium periodate, resulted in a significant decrease in neutrophil recruitment to the peritoneum. Slit2 also prevented neutrophil recruitment to the peritoneal cavity in response to other chemoattractant factors tested, including C5a and MIP-2. These data demonstrate that Slit2 acts as a potent inhibitor of chemotaxis for circulating neutrophils toward diverse inflammatory stimuli. Slit2 also inhibited infiltration of other leukocyte subsets, especially monocytes/macrophages (Tole et al., 2009).




Fig. 6. Slit2/Robo-1 signaling inhibits actin remodelling required for chemotaxis. Chemoattractant signaling induces the activation of Rho GTPases Cdc42 and Rac, allowing for actin remodeling and chemotaxis. The binding of the LRRs on Slit2 to Robo-1 recruits srGAPs to the membrane, converting active Rho GTPases to their inactive, GDP-bound, forms. Inactivation of Rho GTPases abolishes actin remodeling and prevents cell chemotaxis.

4. Conclusion and discussion

The LRRs contained in Slit proteins can inhibit the migration of diverse cells, including neuronal cells and vascular smooth muscle cells. The conserved structure of Slit proteins

also allows them to inhibit the migration of several subsets of leukocytes, including DCs and lymphocytes (Guan et al., 2003; Kanellis et al., 2004; Prasad et al., 2007). We have recently shown that Slit2 inhibits migration of neutrophils to diverse inflammatory attractants, *in vitro* and *in vivo*. Furthermore, we have demonstrated that this inhibition is mediated by inactivation of Rho-family GTPases, Rac and Cdc42 (Fig. 5). Excessive infiltration of leukocytes, particularly neutrophils, is associated with local tissue damage seen in inflammatory conditions such as rheumatoid arthritis and ischemia reperfusion injury (Weissmann et al., 1984; Kaminski et al., 2002). Thus, the protein structure of the conserved LRR regions contained in Slit proteins may be utilized as a novel therapeutic strategy to locally inhibit leukocyte recruitment.

Extensive glycosylation makes Slit2 a large and relatively "sticky" protein, potentially allowing it to maintain a high local concentration through adherence to extracellular matrix proteins such as glypican-1 (Ronca et al., 2001). Thus, after regional administration, Slit2 may be retained at sites of inflammation, such as joints and transplanted organs, thereby alleviating neutrophil-inflicted tissue injury associated with rheumatoid arthritis and ischemia reperfusion injury. As Slit2 blocks migration of several types of inflammatory cells, including neutrophils, T lymphocytes, macrophages, and dendritic cells, toward diverse chemoattractant signals, it could act as a highly effective anti-inflammatory agent (Guan et al., 2003; Kanellis et al., 2004; Prasad et al., 2007; Wu et al., 2001). Further studies are required to explore the therapeutic use of Slit2, or of a Slit-like compound containing the structurally critical LRRs, for prevention and treatment of localized inflammation.

5. Acknowledgment

This work was supported by the Canadian Institute of Health Research (L. A. R.). L. A. R. holds a Canada Research Chair, Tier 2.

6. References

- Allen, W. E., Jones, G. E., Pollard, J. W., & Ridley, A. J. (1997). Rho, rac and Cdc42 regulate actin organization and cell adhesion in macrophages. *Journal of Cell Science*, 110(6), 707-720.
- Alon, R., Hammer, D. A., & Springer, T. A. (1995). Lifetime of the P-selectin-carbohydrate bond and its response to tensile force in hydrodynamic flow. *Nature*, 374(6522), 539.
- Aspenström, P. (1999). Effectors for the rho GTPases. *Current Opinion in Cell Biology*, 11(1), 95.
- Astrof, N., Salas, A., Shimaoka, M., Chen, J., & Springer, T. (2006). Importance of force linkage in mechanochemistry of adhesion receptors. *Biochemistry*, 45(50), 15020.
- Bashaw, G. J., Kidd, T., Murray, D., Pawson, T., & Goodman, C. S. (2000). Repulsive axon guidance: Abelson and enabled play opposing roles downstream of the roundabout receptor. *Cell*, 101(7), 703-715.
- Battye, R., Stevens, A., & Jacobs, J. R. (1999). Axon repulsion from the midline of the drosophila CNS requires slit function. *Development*, 126(11), 2475-2481.
- Battye, R., Stevens, A., Perry, R. L., & Jacobs, J. R. (2001). Repellent signaling by slit requires the leucine-rich repeats. *The Journal of Neuroscience*, 21(12), 4290-4298.

- Benard, V., Bohl, B. P., Bokoch, G. M. (1999). Characterization of Rac and Cdc42 activation in chemoattractant-stimulated human neutrophils using a novel assay for active GTPases. *J. Biol. Chem.* 274, 13198-13204.
- Bertoli-Avella, A., Conte, M., Punzo, F., de Graaf, B., Lama, G., La Manna, A., et al. (2008). ROBO2 gene variants are associated with familial vesicoureteral reflux. *Journal of the American Society of Nephrology*, 19(4), 825-831.
- Beutler, B. (2004). Innate immunity: An overview. *Molecular Immunology*, 40(12), 845-859.
- Bokoch, G. M. (2005). Regulation of innate immunity by rho GTPases. *Trends in Cell Biology*, 15(3), 163.
- Bretscher, M. S. (1996). Getting membrane flow and the cytoskeleton to cooperate in moving cells. *Cell*, 87(4), 601.
- Brose, K., Bland, K. S., Wang, K. H., Arnott, D., Henzel, W., Goodman, C. S., et al. (1999). Slit proteins bind robo receptors and have an evolutionarily conserved role in repulsive axon guidance. *Cell*, 96(6), 795-806.
- Calderwood, D. A., Shattil, S. J., & Ginsberg, M. H. (2000). Integrins and actin filaments: Reciprocal regulation of cell adhesion and signaling. *The Journal of Biological Chemistry*, 275(30), 22607-22610.
- Campbell, J. J., Hedrick, J., Zlotnik, A., Siani, M. A., Thompson, D. A., & Butcher, E. C. (1998). Chemokines and the arrest of lymphocytes rolling under flow conditions. *Science*, 279(5349), 381.
- Casey, P. J., Solski, P. A., Der, C. J., & Buss, J. E. (1989). p21ras is modified by a farnesyl isoprenoid. *Proceedings of the National Academy of Sciences of the United States of America*, 86(21), 8323-8327.
- Cederqvist, K., T. Sorsa, et al. (2001). "Matrix metalloproteinases-2, -8, and -9 and TIMP-2 in tracheal aspirates from preterm infants with respiratory distress." *Pediatrics* 108(3): 686-92.
- Chen, J. H., Wen, L., Dupuis, S., Wu, J. Y., & Rao, Y. (2001). The N-terminal leucine-rich regions in slit are sufficient to repel olfactory bulb axons and subventricular zone neurons. *The Journal of Neuroscience*, 21(5), 1548-1556.
- Chen, J., Wu, W., Li, H., Fagaly, T., Zhou, L., Wu, J., et al. (2000). Embryonic expression and extracellular secretion of xenopus slit. *Neuroscience*, 96(1), 231.
- Clark, R. A., Volpp, B. D., Leidal, K. G., Nauseef, W. M. (1990). Two cytosolic components of the human neutrophil respiratory burst oxidase translocate to the plasma membrane during cell activation. *J. Clin. Invest.* 85, 714-721.
- Constantin, G., Majeed, M., Giagulli, C., Piccio, L., Kim, J. Y., Butcher, E. C., et al. (2000). Chemokines trigger immediate beta2 integrin affinity and mobility changes: Differential regulation and roles in lymphocyte arrest under flow. *Immunity*, 13(6), 759.
- Correia, I., Chu, D., Chou, Y., Goldman, R., & Matsudaira, P. (1999). Integrating the actin and vimentin cytoskeletons: Adhesion-dependent formation of fimbrin-vimentin complexes in macrophages. *The Journal of Cell Biology*, 146(4), 831.
- Critchley, D. R., Holt, M. R., Barry, S. T., Priddle, H., Hemmings, L., & Norman, J. (1999). Integrin-mediated cell adhesion: The cytoskeletal connection. *Biochemical Society Symposia*, 65, 79-99.
- Dallol, A., Da Silva, N., Viacava, P., Minna, J., Bieche, I., Maher, E., et al. (2002). SLIT2, a human homologue of the drosophila Slit2 gene, has tumor suppressor activity and is frequently inactivated in lung and breast cancers. *Cancer Research*, 62(20), 5874-5880.

- Dallol, A., Krex, D., Hesson, L., Eng, C., Maher, E., & Latif, F. (2003). Frequent epigenetic inactivation of the SLIT2 gene in gliomas. *Oncogene*, 22(29), 4611-4616.
- Dallol, A., Morton, D., Maher, E., & Latif, F. (2003). SLIT2 axon guidance molecule is frequently inactivated in colorectal cancer and suppresses growth of colorectal carcinoma cells. *Cancer Research*, 63(5), 1054-1058.
- DeBiasio, R. L., Wang, L. L., Fisher, G. W., & Taylor, D. L. (1988). The dynamic distribution of fluorescent analogues of actin and myosin in protrusions at the leading edge of migrating swiss 3T3 fibroblasts. *The Journal of Cell Biology*, 107(6), 2631-2645.
- Defacque, H., Egeberg, M., Habermann, A., Diakonova, M., Roy, C., Mangeat, P., et al. (2000). Involvement of ezrin/moesin in de novo actin assembly on phagosomal membranes. *EMBO Journal*, 19(2), 199-212.
- DeFife, K. M., Jenney, C. R., Colton, E., & Anderson, J. M. (1999). Cytoskeletal and adhesive structural polarizations accompany IL-13-induced human macrophage fusion. *The Journal of Histochemistry and Cytochemistry*, 47(1), 65-74.
- Dickinson, R., Dallol, A., Bieche, I., Krex, D., Morton, D., Maher, E., et al. (2004). Epigenetic inactivation of SLIT3 and SLIT1 genes in human cancers. *The British Journal of Cancer*, 91(12), 2071.
- Dunne, J., Ballantyne, C., Beaudet, A., & Ley, K. (2002). Control of leukocyte rolling velocity in TNF-alpha-induced inflammation by LFA-1 and mac-1. *Blood*, 99(1), 336.
- Dvorak, Ann., Feng, Dian. (2001). The vesiculo-vacuolar organelle (VVO): a new endothelial cell permeability organelle. *The Journal of Histochemistry & Cytochemistry*, 49(4), 419-431.
- Eltzschig, H. K. and C. D. Collard (2004). "Vascular ischaemia and reperfusion injury." *Br Med Bull* 70: 71-86.
- Eriksson, E. E., Xie, X., Werr, J., Thoren, P., & Lindbom, L. (2001). Importance of primary capture and L-selectin-dependent secondary capture in leukocyte accumulation in inflammation and atherosclerosis in vivo. *The Journal of Experimental Medicine*, 194(2), 205.
- Fauci, A. S., B. Haynes, et al. (1978). "The spectrum of vasculitis: clinical, pathologic, immunologic and therapeutic considerations." *Ann Intern Med* 89(5 Pt 1): 660-76.
- Fenteany, G., & Glogauer, M. (2004). Cytoskeletal remodeling in leukocyte function. *Current Opinion in Hematology*, 11(1), 15-24.
- Finger, E. B., Puri, K. D., Alon, R., Lawrence, M. B., von Andrian, U. H., & Springer, T. A. (1996). Adhesion through L-selectin requires a threshold hydrodynamic shear. *Nature*, 379(6562), 266.
- Frangogiannis, Nikolaos., Smith, Wayne., Entman, Mark. (2002). The inflammatory response in myocardial infarction. *Cardiovascular Research*, 53, 31-47.
- Fujishima, S., Aikawa, N. (1995). Neutrophil-mediated tissue injury and its modulation. *Intensive Care Medicine*, 21(3), 277-285.
- Fujiyama, A., & Tamanoi, F. (1990). RAS2 protein of *saccharomyces cerevisiae* undergoes removal of methionine at N terminus and removal of three amino acids at C terminus. *The Journal of Biological Chemistry*, 265(6), 3362-3368.
- Gamblin, S. J., & Smerdon, S. J. (1998). GTPase-activating proteins and their complexes. *Current Opinion in Structural Biology*, 8(2), 195-201.
- Ganz, T. (2003). Defensins: Antimicrobial peptides of innate immunity. *Nature Reviews Immunology*, 3(9), 710-720.

- Guan, H., Zu, G., Tang, H., Johnson, M., Xu, X., Kevil, C., Xiong, W-C., Elmets, C., Rao, Y., Wu, J. Y., Xu, H. (2003) Neuronal repellent Slit2 inhibits dendritic cell migration and the development of immune responses. *J. Immunol.* 171, 6519–6526.
- Garcia, J. G., H. L. James, et al. (1987). "Lower respiratory tract abnormalities in rheumatoid interstitial lung disease. Potential role of neutrophils in lung injury." *Am Rev Respir Dis* 136(4): 811-7.
- Gertler, F. B., Bennett, R. L., Clark, M. J., & Hoffmann, F. M. (1989). Drosophila abl tyrosine kinase in embryonic CNS axons: A role in axonogenesis is revealed through dosage-sensitive interactions with disabled. *Cell*, 58(1), 103-113.
- Glogauer, M., Marchal, C. C., Zhu, F., Worku, A., Clausen, B. E., Foerster, I., Marks, P., Downey, G. P., Dinauer, M., Kwiatkowski, D. J. (2003). Rac1 deletion in mouse neutrophils has selective effects on neutrophil function. *J. Immunol.* 170, 5652–5657.
- Guan, H., Zu, G., Xie, Y., Tang, H., Johnson, M., Xu, X., et al. (2003). Neuronal repellent Slit2 inhibits dendritic cell migration and the development of immune responses. *The Journal of Immunology*, 171(12), 6519-6526.
- Gu, Y., Filippi, M-D., Cancelas, J. A., Siefring, J. E., Williams, E. P., Jasti, A. C., Harris, C. E., Lee, A. W., Prabhakar, R., Atkinson, S. J., Kwiatkowski, D. J., Williams, D. A. (2003) Hematopoietic cell regulation by Rac1 and Rac2 guanosine triphosphatases. *Science* 302, 445–449.
- Gutierrez, L., Magee, A. I., Marshall, C. J., & Hancock, J. F. (1989). Post-translational processing of p21ras is two-step and involves carboxyl-methylation and carboxy-terminal proteolysis. *EMBO Journal*, 8(4), 1093-1098.
- Hao, J. C., Yu, T. W., Fujisawa, K., Culotti, J. G., Gengyo-Ando, K., Mitani, S., et al. (2001). C. elegans slit acts in midline, dorsal-ventral, and anterior-posterior guidance via the SAX-3/Robo receptor. *Neuron*, 32(1), 25-38.
- Haslam, P. L., C. W. Turton, et al. (1980). "Bronchoalveolar lavage in pulmonary fibrosis: comparison of cells obtained with lung biopsy and clinical features." *Thorax* 35(1): 9-18.
- Hirche, T. O., E. C. Crouch, et al. (2004). "Neutrophil serine proteinases inactivate surfactant protein D by cleaving within a conserved subregion of the carbohydrate recognition domain." *J Biol Chem* 279(26): 27688-98.
- Hirsch, E., Katanaev, V. L., Garlanda, C., Azzolino, O., Pirola, L., Silengo, L., et al. (2000). Central role for G protein-coupled phosphoinositide 3-kinase gamma in inflammation. *Science*, 287(5455), 1049-1053.
- Holdsworth, S. R. and R. Bellomo (1984). "Differential effects of steroids on leukocytemediated glomerulonephritis in the rabbit." *Kidney Int* 26(2): 162-9.
- Holmes, G. P., Negus, K., Burridge, L., Raman, S., Algar, E., Yamada, T., et al. (1998). Distinct but overlapping expression patterns of two vertebrate slit homologs implies functional roles in CNS development and organogenesis. *Mechanisms of Development*, 79(1-2), 57-72.
- Holmes, G., & Niswander, L. (2001). Expression of slit-2 and slit-3 during chick development. *Developmental Dynamics*, 222(2), 301-307.
- Huo, Y., Schober, A., Forlow, S. B., Smith, D., Hyman, M., Jung, S., et al. (2003). Circulating activated platelets exacerbate atherosclerosis in mice deficient in apolipoprotein E. *Nature Medicine*, 9(1), 61.
- Imhof, B., & Aurrand-Lions, M. (2004). Adhesion mechanisms regulating the migration of monocytes. *Nature Reviews.Immunology*, 4(6), 432.

- Ito, H. et al. (2006) Identification of ROBO1 as a novel hepatocellular carcinoma antigen and a potential therapeutic and diagnostic target. *Clin. Cancer Res.* 12, 3257–3264.
- Itoh, A., Miyabayashi, T., Ohno, M., & Sakano, S. (1998). Cloning and expressions of three mammalian homologues of drosophila slit suggest possible roles for slit in the formation and maintenance of the nervous system. *Molecular Brain Research*, 62(2), 175-186.
- Jin, T., Zhang, N., Long, Y., Parent, C. A., & Devreotes, P. N. (2000). Localization of the G protein betagamma complex in living cells during chemotaxis. *Science*, 287(5455), 1034-1036.
- Johnson, Z., Proudfoot, A. E., & Handel, T. M. (2005). Interaction of chemokines and glycosaminoglycans: A new twist in the regulation of chemokine function with opportunities for therapeutic intervention. *Cytokine Growth Factor Reviews*, 16(6), 625.
- Jones, G. E., Allen, W. E., & Ridley, A. J. (1998). The rho GTPases in macrophage motility and chemotaxis. *Cell Adhesion Communication*, 6(2-3), 237-245.
- Jones, C. A., Londin, N. R., Chen, H., Park, K. W., Sauvaget, D., Stockton, R. A., Wythe, J. D., Suh, W., Larrieu-Lahargue, F., Mukoutama, Y-S., Lindblom, P., Seth, P., Frias, A., Nishiya, N., Ginsberg, M. H., Gerhardt, H., Zhang, K., Li, D. Y. (2008) Robo4 stabilizes the vascular network by inhibiting pathologic angiogenesis and endothelial hyperpermeability. *Nat. Med.* 14, 448–453.
- Jones, C., Nishiya, N., London, N., Zhu, W., Sorensen, L., Chan, A., et al. (2009). Slit2-Robo4 signalling promotes vascular stability by blocking Arf6 activity. *Nature Cell Biology*, 11(11), 1325-1331.
- Jung, U., Norman, K. E., Scharffetter-Kochanek, K., Beaudet, A. L., & Ley, K. (1998). Transit time of leukocytes rolling through venules controls cytokine-induced inflammatory cell recruitment in vivo. *Journal of Clinical Investigation*, 102(8), 1526.
- Kaminski, K. A., Bonda, T. A., Korecki, J., Musial, W. J. (2002) Oxidative stress and neutrophil activation—the two keystones of ischemia/reperfusion injury. *Int. J. Cardiol.* 86, 41–59.
- Kanellis, J., Garcia, G., Ping, L., Parra, G., Wilson, C., Rao, Y., et al. (2004). Modulation of inflammation by slit protein in vivo in experimental crescentic glomerulonephritis. *The American Journal of Pathology*, 165(1), 341.
- Kang, T., J. Yi, et al. (2001). "Subcellular distribution and cytokine- and chemokine regulated secretion of leukolysin/MT6-MMP/MMP-25 in neutrophils." *J Biol Chem* 276(24): 21960-8.
- Kansas, G. S. (1996). Selectins and their ligands: Current concepts and controversies. *Blood*, 88(9), 3259.
- Kaur, S., Castellone, MD., Bedell, V., Konar, M., Gutkind, J., Ramchandran, R. (2006). Robo4 signaling in endothelial cells implies attraction guidance mechanisms. *Journal of Biological Chemistry*, 281(16), 11347.
- Keep, N. H., Barnes, M., Barsukov, I., Badii, R., Lian, L. Y., Segal, A. W., et al. (1997). A modulator of rho family G proteins, rhoGDI, binds these G proteins via an immunoglobulin-like domain and a flexible N-terminal arm. *Structure*, 5(5), 623-633.
- Kennedy, J., Kelner, G. S., Kleyensteuber, S., Schall, T. J., Weiss, M. C., Yssel, H., et al. (1995). Molecular cloning and functional characterization of human lymphotactin. *The Journal of Immunology*, 155(1), 203.


- Kennedy, T. E., Serafini, T., de la Torre, J. R., & Tessier-Lavigne, M. (1994). Netrins are diffusible chemotropic factors for commissural axons in the embryonic spinal cord. *Cell*, 78(3), 425-435.
- Kidd, T., Brose, K., Mitchell, K. J., Fetter, R. D., Tessier-Lavigne, M., Goodman, C. S., et al. (1998). Roundabout controls axon crossing of the CNS midline and defines a novel subfamily of evolutionarily conserved guidance receptors. *Cell*, 92(2), 205-215.
- Kintscher, U., Goetze, S., Wakino, S., Kim, S., Nagpal, S., Chandraratna, R. A., et al. (2000). Peroxisome proliferator-activated receptor and retinoid X receptor ligands inhibit monocyte chemotactic protein-1-directed migration of monocytes. *European Journal of Pharmacology*, 401(3), 259-270.
- Kjoller, L., & Hall, A. (1999). Signaling to rho GTPases. *Experimental Cell Research*, 253(1), 166-179.
- Kramer, S. G., Kidd, T., Simpson, J. H., & Goodman, C. S. (2001). Switching repulsion to attraction: Changing responses to slit during transition in mesoderm migration. *Science*, 292(5517), 737-740.
- Krugmann, S., Hawkins, P. T., Pryer, N., & Braselmann, S. (1999). Characterizing the interactions between the two subunits of the p101/p110gamma phosphoinositide 3-kinase and their role in the activation of this enzyme by G beta gamma subunits. *The Journal of Biological Chemistry*, 274(24), 17152-17158.
- Kunkel, E. J., & Ley, K. (1996). Distinct phenotype of E-selectin-deficient mice. E-selectin is required for slow leukocyte rolling in vivo. *Circulation Research*, 79(6), 1196.
- Lanier, L. M., Gates, M. A., Witke, W., Menzies, A. S., Wehman, A. M., Macklis, J. D., et al. (1999). Mena is required for neurulation and commissure formation. *Neuron*, 22(2), 313-325.
- Laudanna, C., Kim, J., Constantin, G., & Butcher, E. (2002). Rapid leukocyte integrin activation by chemokines. *Immunological Reviews*, 186, 37.
- Lawrence, M. B., Kansas, G. S., Kunkel, E. J., & Ley, K. (1997). Threshold levels of fluid shear promote leukocyte adhesion through selectins (CD62L,P,E). *The Journal of Cell Biology*, 136(3), 717.
- Lemanske, R. F., Jr., D. A. Guthman, et al. (1983). "The biologic activity of mast cell granules. VII. The effect of anti-neutrophil antibody-induced neutropenia on rat cutaneous late phase reactions." *J Immunol* 131(2): 929-33.
- Li, C. and R. M. Jackson (2002). "Reactive species mechanisms of cellular hypoxiareoxygenation injury." *Am J Physiol Cell Physiol* 282(2): C227-41.
- Li, Z., Jiang, H., Xie, W., Zhang, W., Smrcka, A., & Wu, D. (2000). Roles of PLC-2 and-3 and PI3K in chemoattractant-mediated signal transduction. *Science's STKE*, 287(5455), 1046.
- Liang, Y., Annan, R. S., Carr, S. A., Popp, S., Mevissen, M., Margolis, R. K., et al. (1999). Mammalian homologues of the drosophila slit protein are ligands of the heparan sulfate proteoglycan glypican-1 in brain. *The Journal of Biological Chemistry*, 274(25), 17885-17892.
- Liu, D., Hou, J., Hu, X., Wang, X., Xiao, Y., Mou, Y., et al. (2006). Neuronal chemorepellent Slit2 inhibits vascular smooth muscle cell migration by suppressing small GTPase Rac1 activation. *Circulation Research*, 98(4), 480-489.
- Linder, S., & Aepfelbacher, M. (2003). Podosomes: Adhesion hot-spots of invasive cells. *Trends in Cell Biology*, 13(7), 376-385.

- Luster, A. D. (1998). Chemokines--chemotactic cytokines that mediate inflammation. *New England Journal of Medicine*, *338*(7), 436.
- Martenson, C., Stone, K., Reedy, M., & Sheetz, M. (1993). Fast axonal transport is required for growth cone advance. *Nature*, *366*(6450), 66-69.
- Mattila, P., & Lappalainen, P. (2008). Filopodia: Molecular architecture and cellular functions. *Nature Reviews.Molecular Cell Biology*, *9*(6), 446-454.
- Marillat, V., Cases, O., Nguyen-Ba-Charvet, K. T., Tessier-Lavigne, M., Sotelo, C., & Chdotal, A. (2002). Spatiotemporal expression patterns of slit and robo genes in the rat brain. *Journal of Comparative Neurology*, *442*(2), 130-155.
- McDermott, David., Fong, Alan., Yang, Qiong., Sechler, Joan., Cupples, Adrienne., Merrell, Maya., Wilson, Peter., D'Agostino, Ralph., O'Donnell, Christopher., Patel, Dhavalkumar., and Murphy, Philip. (2003). Chemokine receptor mutant CX₃CR1-M280 has impaired adhesive function and correlates with protection from cardiovascular disease in humans. *Journal of Clinical Investigation*, *111*(8), 1241-1250.
- McEver, R. P., & Cummings, R. D. (1997). Perspectives series: Cell adhesion in vascular biology. role of PSGL-1 binding to selectins in leukocyte recruitment. *Journal of Clinical Investigation*, *100*(3), 485.
- Mitra, S., Hanson, D., & Schlaepfer, D. (2005). Focal adhesion kinase: In command and control of cell motility. *Nature Reviews.Molecular Cell Biology*, *6*(1), 56-68.
- Mohr, W., H. Westerhellweg, et al. (1981). "Polymorphonuclear granulocytes in rheumatic tissue destruction. III. an electron microscopic study of PMNs at the pannus-cartilage junction in rheumatoid arthritis." *Ann Rheum Dis* *40*(4): 396-9.
- Moore, K. J., Andersson, L. P., Ingalls, R. R., Monks, B. G., Li, R., Arnaout, M. A., et al. (2000). Divergent response to LPS and bacteria in CD14-deficient murine macrophages. *The Journal of Immunology*, *165*(8), 4272-4280.
- Morlot, C., Thielens, N., Ravelli, R. B. G., Hemrika, W., Romijn, R., Gros, P., et al. (2007). Structural insights into the slit-robo complex. *Proceedings of the National Academy of Sciences of the United States of America*, *104*(38), 14923-14928.
- Muller, W. A. (2003). Leukocyte-endothelial-cell interactions in leukocyte transmigration and the inflammatory response. *Trends in Immunology*, *24*(6), 326-333.
- Nassar, N., Hoffman, G. R., Manor, D., Clardy, J. C., & Cerione, R. A. (1998). Structures of Cdc42 bound to the active and catalytically compromised forms of Cdc42GAP. *Nature Structural Biology*, *5*(12), 1047-1052.
- Nguyen Ba-Charvet, K. T. N., Brose, K., Ma, L., Wang KH., Marillat, V., Sotelo, C., et al. (2001). Diversity and specificity of actions of Slit2 proteolytic fragments in axon guidance. *The Journal of Neuroscience*, *21*(12), 4281.
- Nguyen, D. H., Catling, A. D., Webb, D. J., Sankovic, M., Walker, L. A., Somlyo, A. V., et al. (1999). Myosin light chain kinase functions downstream of Ras/ERK to promote migration of urokinase-type plasminogen activator-stimulated cells in an integrin-selective manner. *The Journal of Cell Biology*, *146*(1), 149-164.
- Nobes, C. D., & Hall, A. (1995). Rho, rac, and cdc42 GTPases regulate the assembly of multimolecular focal complexes associated with actin stress fibers, lamellipodia, and filopodia. *Cell*, *81*(1), 53-62.
- Oloffson, B. (1999). Rho Guanine Dissociation Inhibitors - Pivotal Molecules in Cellular Signalling . *Cellular Signalling*, *11*(8), 545-554.
- Owen, C. A. and E. J. Campbell (1999). "The cell biology of leukocyte-mediated proteolysis." *J Leukoc Biol* *65*(2): 137-50.

- Panes, J., M. Perry, et al. (1999). "Leukocyte-endothelial cell adhesion: avenues for therapeutic intervention." *Br J Pharmacol* 126(3): 537-50.
- Piper, M., Georgas, K., Yamada, T., & Little, M. (2000). Expression of the vertebrate slit gene family and their putative receptors, the robo genes, in the developing murine kidney. *Mechanisms of Development*, 94(1-2), 213-217.
- Prasad, A., Fernandis, A., Rao, Y., & Ganju, R. (2004). Slit protein-mediated inhibition of CXCR4-induced chemotactic and chemoinvasive signaling pathways in breast cancer cells. *The Journal of Biological Chemistry*, 279(10), 9115-9124.
- Prasad, A., Qamri, Z., Wu, J., & Ganju, R. (2007). Slit-2/Robo-1 modulates the CXCL12/CXCR4-induced chemotaxis of T cells. *Journal of Leukocyte Biology*, 82(3), 465-476.
- Puri, D., Finger, B., Springer, A. (1995) Sialomucin CD34 is the major L-selectin ligand in human tonsil high endothelial venules. *The Journal of Cell Biology*, 131(1), 261-270.
- Ray, L. B. (2004). STKE: Slit and robo in kidney formation. *Science*, 304(5675), 1215c.
- Ridley, A. J. (2001). Rho proteins, PI 3-kinases, and monocyte/macrophage motility. *FEBS Letters*, 498(2-3), 168.
- Ridley, A. J., & Hall, A. (1992). The small GTP-binding protein rho regulates the assembly of focal adhesions and actin stress fibers in response to growth factors. *Cell*, 70(3), 389-399.
- Robinson, L. A., Nataraj, C., Thomas, D. W., Howell, D. N., Griffiths, R., Bautch, V., et al. (2000). A role for fractalkine and its receptor (CX3CR1) in cardiac allograft rejection. *The Journal of Immunology*, 165(11), 6067.
- Ronca, F., Andersen, J. S., Paech, V., & Margolis, R. U. (2001). Characterization of slit protein interactions with glypican-1. *The Journal of Biological Chemistry*, 276(31), 29141-29147.
- Rossi, D., & Zlotnik, A. (2000). The biology of chemokines and their receptors. *Annual Review of Immunology*, 18, 217-242.
- Rossmann, K., Der, C., & Sodek, J. (2005). GEF means go: Turning on RHO GTPases with guanine nucleotide-exchange factors. *Nature Reviews.Molecular Cell Biology*, 6(2), 167-180.
- Rothberg, J. M., & Artavanis-Tsakonas, S. (1992). Modularity of the slit protein. characterization of a conserved carboxy-terminal sequence in secreted proteins and a motif implicated in extracellular protein interactions. *Journal of Molecular Biology*, 227(2), 367-370.
- Rothberg, J. M., Hartley, D. A., Walther, Z., & Artavanis-Tsakonas, S. (1988). Slit: An EGF-homologous locus of *D. melanogaster* involved in the development of the embryonic central nervous system. *Cell*, 55(6), 1047-1059.
- Rothberg, J. M., Jacobs, J. R., Goodman, C. S., & Artavanis-Tsakonas, S. (1990). Slit: An extracellular protein necessary for development of midline glia and commissural axon pathways contains both EGF and LRR domains. *Genes Development*, 4(12A), 2169-2187.
- Rubio, F., J. Cooley, et al. (2004). "Linkage of neutrophil serine proteases and decreased surfactant protein-A (SP-A) levels in inflammatory lung disease." *Thorax* 59(4): 318-23.
- Salas, A., Shimaoka, M., Kogan, A., Harwood, C., von Andrian, U., & Springer, T. (2004). Rolling adhesion through an extended conformation of integrin alphaLbeta2 and relation to alpha I and beta I-like domain interaction. *Immunity*, 20(4), 393.

- Sasaki, T., Irie-Sasaki, J., Jones, R. G., Oliveira-dos-Santos, A. J., Stanford, W. L., Bolon, B., et al. (2000). Function of PI3K γ in thymocyte development, T cell activation, and neutrophil migration. *Science*, 287(5455), 1040-1046.
- Sehr, P., Joseph, G., Genth, H., Just, I., Pick, E., Aktories, K. (1998) Glucosylation and ADP ribosylation of Rho proteins: effects of nucleotide binding, GTPase activity, and effector coupling. *Biochemistry* 37, 5296-5304.
- Seki, M., Watanabe, A., Enomoto, S., Kawamura, T., Ito, H., Kodama, T., et al. (2010). Human ROBO1 is cleaved by metalloproteinases and γ -secretase and migrates to the nucleus in cancer cells. *FEBS Letters*, 584(13), 2909-2915.
- Servant, G., Weiner, O. D., Herzmark, P., Balla, T., Sedat, J. W., & Bourne, H. R. (2000). Polarization of chemoattractant receptor signaling during neutrophil chemotaxis. *Science*, 287(5455), 1037-1040.
- Shamri, R., Grabovsky, V., Gauguier, J., Feigelson, S., Manevich, E., Kolanus, W., et al. (2005). Lymphocyte arrest requires instantaneous induction of an extended LFA-1 conformation mediated by endothelium-bound chemokines. *Nature Immunology*, 6(5), 497.
- Shin, H. S., Snyderman, R., Friedman, E., Mellors, A., & Mayer, M. M. (1968). Chemotactic and anaphylatoxic fragment cleaved from the fifth component of guinea pig complement. *Science*, 162(851), 361.
- Sigal, A., Bleijs, D. A., Grabovsky, V., van Vliet, S. J., Dwir, O., Figdor, C. G., et al. (2000). The LFA-1 integrin supports rolling adhesions on ICAM-1 under physiological shear flow in a permissive cellular environment. *The Journal of Immunology*, 165(1), 442-452.
- Singer, S. J., & Kupfer, A. (1986). The directed migration of eukaryotic cells. *Annual Review of Cell Biology*, 2, 337-365.
- Srinivasan, S., Wang, F., Glavas, S., Ott, A., Hofmann, F., Aktories, K., Kalman, D., Bourne, H. R. (2003). Rac and Cdc42 play distinct roles in regulating PI(3,4,5)P3 and polarity during neutrophil chemotaxis. *J. Cell Biol.* 160, 375-385.
- Suchting, S., Heal, P., Tahtis, K., Stewart L., and Bicknell, R. (2005). Soluble Robo4 receptor inhibits in vivo angiogenesis and endothelial cell migration. *The FASEB Journal*, 19(1), 121-123.
- Sun, C. X., Downey, G. P., Zhu, F., Koh, A. L. Y., Thang, H., Glogauer, M. (2004). Rac1 is the small GTPase responsible for regulating the neutrophil chemotaxis compass. *Blood* 104, 3758-3765.
- Symons, M. H., & Mitchison, T. J. (1991). Control of actin polymerization in live and permeabilized fibroblasts. *The Journal of Cell Biology*, 114(3), 503-513.
- Tcherkezian, J., & Lamarche-Vane, N. (2007). Current knowledge of the large RhoGAP family of proteins. *Biology of the Cell*, 99(2), 67-86.
- Vanhaesebroeck, B., & Waterfield, M. D. (1999). Signaling by distinct classes of phosphoinositide 3-kinases. *Experimental Cell Research*, 253(1), 239-254.
- Vandivier, R. W., V. A. Fadok, et al. (2002). "Elastase-mediated phosphatidylserine receptor cleavage impairs apoptotic cell clearance in cystic fibrosis and bronchiectasis." *J Clin Invest* 109(5): 661-70.
- Vargesson, N., Luria, V., Messina, I., Erskine, L., & Laufer, E. (2001). Expression patterns of slit and robo family members during vertebrate limb development. *Mechanisms of Development*, 106(1-2), 175-180.
- Vetter, I. R., & Wittinghofer, A. (2001). The guanine nucleotide-binding switch in three dimensions. *Science*, 294(5545), 1299-1304.

- von Hundelshausen, P., Weber, K. S., Huo, Y., Proudfoot, A. E., Nelson, P. J., Ley, K., et al. (2001). RANTES deposition by platelets triggers monocyte arrest on inflamed and atherosclerotic endothelium. *Circulation*, 103(13), 1772.
- Wang, K. H., Brose, K., Arnott, D., Kidd, T., Goodman, C. S., Henzel, W., et al. (1999). Biochemical purification of a mammalian slit protein as a positive regulator of sensory axon elongation and branching. *Cell*, 96(6), 771-784.
- Wang, Y. L. (1985). Exchange of actin subunits at the leading edge of living fibroblasts: Possible role of treadmilling. *The Journal of Cell Biology*, 101(2), 597.
- Wehrle, R., Camand, E., Chedotal, A., Sotelo, C., & Dusart, I. (2005). Expression of netrin-1, slit-1 and slit-3 but not of slit-2 after cerebellar and spinal cord lesions. *European Journal of Neuroscience*, 22(9), 2134-2144.
- Weissmann, G. and H. Korchak (1984). "Rheumatoid arthritis. The role of neutrophil activation." *Inflammation* 8 Suppl: S3-14.
- Weissmann, G., R. B. Zurier, et al. (1971). "Mechanisms of lysosomal enzyme release from leukocytes exposed to immune complexes and other particles." *J Exp Med* 134(3 Pt2): 149s-165s.
- Wieland, T. and C. K. Chen (1999). "Regulators of G-protein signalling: a novel protein family involved in timely deactivation and desensitization of signalling via heterotrimeric G proteins." *Naunyn Schmiedebergs Arch Pharmacol* 360(1): 14-26.
- Wills, Z., Marr, L., Zinn, K., Goodman, C. S., & Van Vactor, D. (1999). Profilin and the abl tyrosine kinase are required for motor axon outgrowth in the drosophila embryo. *Neuron*, 22(2), 291-299.
- Wong, K., Ren, X. R., Huang, Y. Z., Xie, Y., Liu, G., Saito, H., et al. (2001). Signal transduction in neuronal migration: Roles of GTPase activating proteins and the small GTPase Cdc42 in the slit-robo pathway. *Cell*, 107(2), 209-221.
- Worthylake, R. A., Lemoine, S., Watson, J. M., & Burridge, K. (2001). RhoA is required for monocyte tail retraction during transendothelial migration. *The Journal of Cell Biology*, 154(1), 147-160.
- Wu, J. Y., Feng, L., Park, H. T., Havlioglu, N., Wen, L., Tang, H., et al. (2001). The neuronal repellent slit inhibits leukocyte chemotaxis induced by chemotactic factors. *Nature*, 410(6831), 948-952.
- Yamada, K. M., & Miyamoto, S. (1995). Integrin transmembrane signaling and cytoskeletal control. *Current Opinion in Cell Biology*, 7, 681.
- Yang, L., Froio, R., Sciuto, T., Dvorak, A., Alon, R., Luscinskas, F. (2005). ICAM-1 regulates neutrophil adhesion and transcellular migration of TNF- α -activated vascular endothelium under flow. *Blood*, 106(2), 584.
- Ye, BQ., Geng, Z., Ma, L., & Geng, J. (2010). Slit2 regulates attractive eosinophil and repulsive neutrophil chemotaxis through differential srGAP1 expression during lung inflammation. *The Journal of Immunology*, 185(10), 6294.
- Yuan, W., Zhou, L., Chen, J. H., Wu, J. Y., Rao, Y., & Ornitz, D. M. (1999). The mouse SLIT family: Secreted ligands for ROBO expressed in patterns that suggest a role in morphogenesis and axon guidance. *Developmental Biology*, 212(2), 290-306.
- Zallen, J. A., Yi, B. A., & Bargmann, C. I. (1998). The conserved immunoglobulin superfamily member SAX-3/Robo directs multiple aspects of axon guidance in *C. elegans*. *Cell*, 92(2), 217-227.
- Zigmond, S. H. (1974). Mechanisms of sensing chemical gradients by polymorphonuclear leukocytes. *Nature*, 249, 450.



Protein Structure

Edited by Dr. Eshel Faraggi

ISBN 978-953-51-0555-8

Hard cover, 396 pages

Publisher InTech

Published online 20, April, 2012

Published in print edition April, 2012

Since the dawn of recorded history, and probably even before, men and women have been grasping at the mechanisms by which they themselves exist. Only relatively recently, did this grasp yield anything of substance, and only within the last several decades did the proteins play a pivotal role in this existence. In this expose on the topic of protein structure some of the current issues in this scientific field are discussed. The aim is that a non-expert can gain some appreciation for the intricacies involved, and in the current state of affairs. The expert meanwhile, we hope, can gain a deeper understanding of the topic.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Ilya M. Mukovozov and Lisa A. Robinson (2012). Slit/Robo Signaling: Inhibition of Directional Leukocyte Migration, Protein Structure, Dr. Eshel Faraggi (Ed.), ISBN: 978-953-51-0555-8, InTech, Available from: <http://www.intechopen.com/books/protein-structure/slit-robo-signaling-inhibition-of-directional-leukocyte-migration>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen