

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,600

Open access books available

138,000

International authors and editors

175M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# Impacts of Carbon Dioxide Gas Leaks from Geological Storage Sites on Soil Ecology and Above-Ground Vegetation

Raveendra H. Patil

*Agricultural and Biological Engineering Department,  
Institute of Food and Agricultural Sciences, University of Florida, Gainesville  
USA*

## 1. Introduction

The global annual carbon dioxide (CO<sub>2</sub>) gas emissions, one of the major green house gases contributing to global warming, into atmosphere have increased to 31.5 Gt in 2008 (German renewable energy institute [IWR], 2009), of which little over two-third (~22 Gt) were emitted from manmade sources (Benson, 2005). Electricity production and transportation make up two-third of total manmade emissions while the rest is contributed from heating buildings and other industrial consumption. Our dependence on fossil fuels continues to be increasing with over 85% of the world's energy needs still coming from burning oil, coal and natural gas (Benson, 2005) and by the end of 21<sup>st</sup> century our demand for fossil fuel is projected to more than double. Some of the scenarios (e.g. with no action taken to limit emissions) suggest a doubling of CO<sub>2</sub> emissions by 2050. This increased energy demand is mainly driven by continued industrialization and improved quality of life in not only western countries (North America and Europe), but also in two of the largest populated countries in the world viz., China and India. This would further increase CO<sub>2</sub> gas emissions into atmosphere with a strong positive feedback on climate change. This demands urgent action to reduce or offset CO<sub>2</sub> emissions from fossil fuels. In this context, while the efforts are being made for cleaner and efficient use of fossil fuels, carbon capture and storage technology is being considered as one of the main short-term viable strategy to help mitigation climate change in coming decades (Bachu, 2000; Intergovernmental Panel on Climate Change [IPCC], 2005; Pacala et al., 2004; Van de Zwaan & Smekens, 2009).

### 1.1 Carbon capture and storage technology

The carbon capture and storage technology basically includes two approaches: (1) CO<sub>2</sub> gas is captured directly from the large and stationary source points (e.g. power plants, petroleum refineries, gas processing units & cement factories), concentrated into a nearly pure form, transported through pipelines and injected into geological storage sites (on- and off- shore) far below the ground surface (Bachu, 2000). This technology is called as "Carbon Capture and Storage" (CCS), and (2) atmospheric CO<sub>2</sub> gas is biologically fixed by growing vegetation (e.g. forest trees, biomass crops) and stored in above- and below-ground plant parts. This is referred to as "Carbon Sequestration" (CS) and these two approaches complement each

other as CCS approach, unlike CS, cannot capture CO<sub>2</sub> gas directly from the atmosphere. Hence, CCS technology is increasingly seen as one of the mechanisms that can make a useful contribution to the reduction of CO<sub>2</sub> emissions over the next 50 years (IPCC, 2005). Some estimates predict that CCS technology has the potential to capture and store CO<sub>2</sub> emissions from power generation by 80–90% (Department of Trade and Industry, UK [DTI], 2002). However, in order to have CCS that significant impact, CO<sub>2</sub> gas will have to be injected on a large scale (in Gt year<sup>-1</sup>).

### 1.2 The geological storage sites and their storage capacity

Geological formations suitable for CO<sub>2</sub> storage are located in sedimentary basins where thick sediments have accumulated over millions of years (Benson, 2005). Such sites are oil and gas reservoirs, deep saline aquifers with suitable caprocks and deep unmineable coal beds. Some estimates suggest that depleted oil and gas reservoirs can store as much as 800 Gt of CO<sub>2</sub> gas (Freund et al, 2003). These sites are known to trap buoyant fluids such as CO<sub>2</sub> and CH<sub>4</sub> underground for millions of years. Hence, CCS technology intends to adopt the methods already used by oil / gas exploration and production, and enhanced oil recovery (EOR) schemes. Under EOR schemes CO<sub>2</sub> gas is injected into oil reservoirs to make it dissolve into the oil which reduces the viscosity of oil while increasing its volume to enhance reservoir pressure and production (Benson, 2005). For example, EOR scheme at Williston Basin oilfield, Canada started injecting ~5000 tonnes of CO<sub>2</sub> day<sup>-1</sup> in October 2000. The CO<sub>2</sub> gas is being transported through 330 km long pipeline from the lignite-fired Dakota Gasification Company plant site in North Dakota, USA (DTI, 2005). The saline aquifers are the reservoirs deep underground and contain saline water (not suitable for drinking) and the global storage capacity of these sites is estimated to be between 400 and 10,000 Gt of CO<sub>2</sub> (Freund et al., 2003). Whereas, the unmineable coal beds, which are located too deep beneath the ground making them uneconomical to explore, can also be used to inject CO<sub>2</sub> gas where the injected gas is absorbed onto the coal. The global storage capacity of unmineable coal beds is estimated to be around 148 Gt of CO<sub>2</sub> (Freund et al., 2003).

### 1.3 Potential risks of failures of CCS technology

Presently existing CCS technology suggest that geological formations selected for CO<sub>2</sub> storage needs to be located at a minimum depth of 800 m so that CO<sub>2</sub> gas is stored at supercritical state to trap large amounts of gas in a small volume. However, in order to implement CCS technology on large and commercial scale, it is essential to assess all the potential risks and provide evidence to inform governments and the public that potential risks are well understood and impact assessments are studied for long-term safety and control measures (Wei et al., 2011).

As the CO<sub>2</sub> gas is captured from the large production sites located on land (e.g. power plants), the gas needs to be transported through pipelines over a long distance. The first source of potential leakage, therefore, would be pipeline failures or small leaks from joints. The scale of leakage and the potential impacts on surrounding environment from such leakages depends on whether the pipeline is laid over or under the ground and whether the pipeline pass through built-up area or near large drinking water sources (e.g. inland lakes or drinking water reservoirs). Pipeline corrosion over a period of time is another issue that might cause gas leakage. CO<sub>2</sub> gas transportation via pipelines in a supercritical state is

reported to cause corrosion of steel pipes @  $0.01 \text{ mm year}^{-1}$ . However, the corrosion rate further increases to  $0.7 \text{ mm year}^{-1}$  if free water was present in the pipeline (IPCC, 2005). If other gasses like hydrogen sulfide or hydrocarbons are mixed with  $\text{CO}_2$  gas, then the chances of leakage and corrosion increase further (Klusman, 2003). Therefore, the pipelines require continuous surveillance for leakage or third party intrusions or encroachments. For example, the Cortez pipeline in Colorado, USA which is buried 1 m below ground, but passes through build-up areas is being air monitored once every two weeks (IPCC, 2005).

The leakage from geological storage sites may also occur due to failure of the sealing cap of injection well or migration of gas through geologic media and lead to slow but large releases either due to over-pressurization or slow releases via faults and fractures (Heinrich et al., 2003; Klusman, 2003). While the past evidence from oil and gas fields shows that the natural gases and fluids, including  $\text{CO}_2$  gas, can be held intact underground for millions of years, incidences like the one of McElmo dome leakage (Gerlach et al., 1998; Stevens et al., 2000) demand complete evaluation of storage sites before selecting for gas injection (IPCC, 2005). Therefore, in order for CCS to be effective, the CCS technology must ensure that leakage is minimized from both sudden releases due to accidents or technical failures and slow leaks over longer period of time.

Perry (2005) reports that more than 600 natural gas storage reservoirs exist across the globe, but to-date on only 10 occasions significant leakage have occurred. These leakages were mainly attributed to failure of bore well integrity (5 times), leaks in the caprocks (4 times) and poor site selection (1 time). However, when it comes to long-term safety of CCS technology there are uncertainties as we lack the experience on the long-term fate (100 to 1000 years) and safety of large volume of  $\text{CO}_2$  gas to be injected into geological formations (Celia et al., 2002). Some of the naturally occurring  $\text{CO}_2$  springs and volcanic sites located across the world have been emitting  $\text{CO}_2$  gas with no severe effect on our ecosystems or population, but the risks of  $\text{CO}_2$  gas leakage from CCS transport pipelines or storage sites will be on larger scale as huge amounts of  $\text{CO}_2$  gas is being handled under CCS schemes (IPCC, 2005). If leaks were to occur, the CCS technology would defy the very purpose to help mitigate climate change (Heinrich et al., 2003). While the existing CCS technology claims to reduce the risks of such leakages by applying safety systems in place and selecting safe geological storage sites (both on- & off-shore), there are chances that slow but continuous releases from CCS sites could go unnoticed as leaking  $\text{CO}_2$  gas would quickly diffuse in the atmosphere (Heinrich et al., 2003). Therefore, it is important to understand the effects of  $\text{CO}_2$  gas leaks on surrounding environments viz., marine life if gas were to be injected into off-shore geological sites and, vegetation and soil ecosystems if gas were to be injected into on-shore geological sites.

#### **1.4 Effects of naturally occurring $\text{CO}_2$ leaks on soil geochemistry**

Elevated soil  $\text{CO}_2$  concentrations can cause changes in mineralogy composition together with changes in trace elements like As and Cr (Kruger et al., 2009; Stenhouse et al., 2009). Changes in cation exchange capacity (CEC), and the presence of oxides like CaO, MgO,  $\text{Fe}_2\text{O}_3$ , and  $\text{Mn}_3\text{O}_4$  have also been reported elsewhere (Blake et al., 2000; Billett et al., 1987; Goulding et al., 1998). Leakage of  $\text{CO}_2$  may also reduce groundwater pH besides affecting taste, color or smell and cause significant deterioration in the quality of potable groundwater by altering groundwater chemistry (Stenhouse et al., 2009).  $\text{CO}_2$  leakage rising

to sub-soil levels may also cause changes in subsurface microbial populations either by favoring some species or restricting others, depending on species type and site characteristics (Jossi et al., 2006; Tian et al., 2001).

### 1.5 Effects of naturally occurring CO<sub>2</sub> leaks on overlying vegetation

CO<sub>2</sub> is an odorless and non-toxic gas, and considered an integral part of our everyday lives. However, exposure to high concentrations of CO<sub>2</sub> poses danger to human beings, animals and surrounding environment, but such leaks may increase soil CO<sub>2</sub> concentrations in near surface and below vegetation canopies. Such a situation could have significant effects on above-ground vegetation and soil inhabiting organisms (e.g. earth worms, microorganisms), both in the short- and long-term. While plants in general are known to be more tolerant to elevated CO<sub>2</sub> gas, persistent leaks from geological storage sites may lead to accumulation of CO<sub>2</sub> gas in soil (near and sub surface). This may suppress root respiration, alter plant water / nutrient uptake capacity by altering soil pH towards acidity and ultimately affect above-ground biomass (Celia et al., 2002; Cook et al., 1998; Gahrooei, 1998; Miglietta et al., 1998; Sorey et al., 2000; Sowerby et al., 2000; Stephens & Hering, 2002). There are many studies where forest trees or perennial vegetation mortality from naturally occurring active volcanoes emitting CO<sub>2</sub> gas into atmosphere have also been documented (e.g. Macek et al., 2005; Stephens & Hering, 2002; Vodnik et al., 2006). In fact, to-date naturally occurring CO<sub>2</sub> springs and active volcanic sites are the ones quite extensively studied in Europe and America.

At Mammoth Mountain, USA, more than 30% by volume of soil CO<sub>2</sub> levels were measured at sites where tree mortality has occurred (Gerlach et al., 2001). At Bossoleto, Italy, soil CO<sub>2</sub> leaks increasing atmospheric CO<sub>2</sub> concentrations to as high as 75% by volume at night, but much lower during day time, were recorded (Van Gardingen et al., 1995). Low levels of CO<sub>2</sub> concentrations during day time, the period when plants are photosynthetically active, may have reduced the adverse effect of elevated CO<sub>2</sub> gas on trees and herbaceous plants or these plant species may well have adapted to elevated CO<sub>2</sub> gas after being exposed for long period of time (van Gardingen et al., 1995). While anaerobic conditions (anoxic or hypoxic) are harmful to plants, many species are known to adapt to such conditions, at least temporarily, by modifying their rooting system and supplying O<sub>2</sub> internally from leaves to roots (Vartapetian & Jackson, 1997). This may have been the case at Bossoleto, Italy where *Phragmites australis*, a wetland species adapted to temporary anoxic conditions was reported to be the most dominant plant cover (Van Gardingen et al., 1995).

Most of studies referred in above paragraphs including the ones reported elsewhere (Beaubien et al., 2008; Biondi & Fessenden, 1999; Vodnik et al., 2006;) have examined the effects of naturally occurring CO<sub>2</sub> leaks from active volcanoes / geothermal sites, and hot / cold CO<sub>2</sub> springs on ecosystems. At these environments the existing ecosystems have been exposed to elevated CO<sub>2</sub> for considerably long periods, thus the plant / tree species may have adapted (West et al., 2009). Therefore, the findings from these sites may not be representative of the effects of potential leakage from a CCS storage sites. Moreover, many of these sites release not only hot or cold CO<sub>2</sub> gas but also gas mixed with either volcanic ash or mineral particles or with other gasses like CH<sub>4</sub> (Bergfeld et al., 2006),

which makes it difficult to isolate the effect of CO<sub>2</sub> gas alone. Furthermore, the effects of slow release of CO<sub>2</sub> from CCS underground transport pipelines or geological storage sites on plant and soil inhabiting organisms are not well studied and needs better understanding (Lewicki et al., 2005). Past studies also suggest that not all the plant species respond similarly when exposed to elevated soil CO<sub>2</sub> concentrations (Van Gardingen et al., 1995), hence makes it difficult to define a “lethal CO<sub>2</sub> concentration level” as different vegetation types respond differently to anoxic stress or anaerobic soil environments due to severe dearth of O<sub>2</sub> level in the soil. Added to this, variations amongst the natural ecosystems and their surrounding environment make it difficult to gauge exposure levels (Sarah & Sjoersten, 2009).

Hence, understanding the risk of CO<sub>2</sub> leakage from storage sites and studying the effect of leaking gas on surface ecosystem is critical for multiple reasons. First, accurate data on the quantity of CO<sub>2</sub> that has been injected and stored is required for trading and accounting purposes; second, any leakage will negate the original purpose of the CCS technology and; third, the leaking CO<sub>2</sub> might damage surface ecosystems including above-ground vegetation and soil ecology. Findings from such studies would enable CCS technology to adopt full safety measures while transporting and injecting huge amounts of CO<sub>2</sub> gas into on-shore geological storage sites, and alleviate the public perceptions, if any, on long-term safety of CCS technology.

In order to test the response of overlying vegetation and soil ecology to gas leaks an experimental field facility, the Artificial Soil Gassing And Response Detection (ASGARD), has been established at the University of Nottingham, UK where gas can be artificially injected into the soil to simulate build up of gas concentrations and its slow release to soil surface (Photo 1). This facility has enabled to study some impacts of a controlled injection of CO<sub>2</sub> gas on a non-adapted pasture grass and field crops, and on soil ecology and chemistry. This chapter describes some of the main findings from studies carried out at the ASGARD site.

Between 2002 and 2005, the ASGARD site was used to investigate the effect of elevated concentrations of soil CH<sub>4</sub> on pasture grass, wheat and winter bean crops (Smith et al., 2005). The natural gas, a major source of energy supply in Europe, is mainly composed of CH<sub>4</sub> gas (78-95% by vol.). European natural gas sub-surface transportation pipelines are under regular helicopter surveillance to detect any gas leaks from pipeline joints, third party incursion and land slippage which may physically damage pipeline and gas supply (Smith et al., 2005). According to Baggott et al. (2003) the CH<sub>4</sub> gas leakage from UK natural gas transportation system alone amounts to 342 kilo tonnes year<sup>-1</sup>, which is equivalent to 10260 kilo tonnes of CO<sub>2</sub> in terms of global warming potential. Thus, early warning system to detect CH<sub>4</sub> gas leaks would enable the surveillance system to take immediate control measures. Therefore, Smith et al., (2005) artificially injected CH<sub>4</sub> gas into the soil at ASGARD site, to simulate CH<sub>4</sub> leakage and monitored above-ground vegetation stress symptoms using remote sensing technology.

Since 2006 the ASGARD site is being used to inject CO<sub>2</sub> gas, initially by the British Geological Survey, UK (West et al., 2009) and later by Patil et al. (2010) and Sarah and Sjoersten, (2009) in 2007-08. Therefore, this chapter while describing the ASGARD site in

detail presents main findings of previous studies undertaken at this site. These studies were undertaken with multiple objectives: (1) to develop a field site where CO<sub>2</sub> gas could be artificially injected at a targeted rate into the soil to simulate build up of soil CO<sub>2</sub> concentrations in near-surface and its leakage into the atmosphere, (2) to monitor temporal and spatial variations in soil gas concentrations under different land cover, and (3) to study the response of vegetation(s), soil properties, and soil inhabiting organisms to elevated soil CO<sub>2</sub> concentrations.



Photo 1. The ASgard site. Both pasture and fallow plots with gassing pipes and measuring tubes can be seen.

## 2. Methodology

### 2.1 The study location and the ASgard site

The ASgard research facility is located on a flat and open field of permanent pasture at the Sutton Bonington campus of the University of Nottingham (52° 49' 60 N, 1° 14' 60 E, 48 m a.s.l.), approximately 18 km south of central Nottingham, UK (Figure 1). The site was previously used as a sheep pasture and had remained grassland for over 10 years until the ASgard facility was laid out. Long term temperature average (1971–2000) at this site shows January as the coldest month with maximum and minimum temperatures of 6.9 and 1.2 °C, respectively and July as the warmest month with maximum and minimum temperatures of 21.3 and 11.4 °C, respectively. The mean annual precipitation is 606 mm; most of it is received as precipitation, and is fairly uniformly distributed throughout the year (Sarah & Sjogersten, 2009). The ASgard site is positioned in such a way that it is not influenced by shade from trees or fencing.




Fig. 1. Location of the ASgard field facility and the main experimental block. Green- main experimental plots with a total of 30 plots, and Yellow - 4 remote control area located >10 m away from the main experimental plots. Blue- gas cylinders and port-cabin which accommodated automated gas supplying hardware and software system.

## 2.2 Site mineralogy and soil characteristics

Geologically, the study area is characterized by up to 1.5 m of head deposit overlying mudstones of the Mercia Mudstone Group. Mineralogical analyses in 2006 showed that quartz was the dominant mineral (>90% of the total dry weight) followed by K-feldspar and albite as minor along with trace amounts of mica, kaolinite, chlorite and hematite (West et al., 2009). Mineralogical composition between the A horizon (0.15–0.30 m depth) and B horizon (0.45–0.50 m depth) were found to be the same (West et al., 2009). However, readers are directed to refer to Ford (2006) for detailed geological description of this site and its surrounding area. The soil type lies within the Worcester Series and comprises 0.3 m deep sandy clay loam over 0.7 m clay and marl horizon. The top soil layer (~ 0.1 m) contains 8.91% clay, 22.89% silt and 68.2% sand (West et al., 2009).

## 2.3 Experimental plots and gas supply system

In the year 2006 a total of 30 plots each of 2.5 m × 2.5 m in size were laid out in a rectangular grid patterns (5 × 6 plots) with 0.5 m pathways between each plot (Figure 2). Ten plots were kept under already established pasture grass and rest were planted with agricultural crops after removing the pasture grass from those plots (West et al., 2009). CO<sub>2</sub> gas was delivered into the pasture plots at a constant rate of 3 liters min<sup>-1</sup> for 19 weeks (May to September 2006). In the year 2007, Sarah and Sjogersten (2009) used 8 plots to plant commercial turf, previously planted with agricultural crops, but left untreated for more than 8 months before they began the study. The turf grass was composed of *Lolium perenne*, *Festuca rubra*, *Festuca rubra commutate* and *Poa pratensis* types. The CO<sub>2</sub> gassing was started 6 weeks after planting the turf grass. Of the total 8 turf grass plots, 4 plots were injected with CO<sub>2</sub> gas at a constant



rate of 1 liter  $\text{min}^{-1}$  for 10 weeks (June to August 2007). The remaining 4 plots were left un-gassed (control plots). Patil et al. (2010) used a total of 16 plots: 8 pasture grass plots and 8 fallow plots which were previously cultivated with agricultural crops, but left untreated for more than 6 months. Pasture and fallow plots were chosen to represent two land use types: perennial pasture grass and fallow land (bare soil). Eight out of these 16 plots (4 pasture & 4 fallow), chosen randomly, were equipped for controlled  $\text{CO}_2$  release at the center of the plots at a constant rate of 1 liter  $\text{min}^{-1}$  for 9 months (May 2007 through January 2008). This subsurface gas injection combined with the flat terrain of the field site prevented build up of air  $\text{CO}_2$  concentration at the site.

During all these studies pure industrial  $\text{CO}_2$  gas was supplied via cryogenic cylinders (supplied by British Oil Company [BOC], UK) and the gas flow to each one of those gassing plots was controlled automatically by individual mass flow controllers (Photo 2). The  $\text{CO}_2$  gas was delivered from the cryogenic cylinder using a 32 mm polyethylene gas pipe. Automated system in turn released the gas at a pre-determined rate into 15-mm copper tubes, one tube to each gassing plot. These copper tubes carried the gas up to experimental plots, where these tubes were separately connected to 22-mm internal diameter medium density polyethylene (MDPE) gas pipes, sealed at the far end (the end which went into soil at the centre of plot). To avoid obstructing the measurement area above-ground within each gassing plot, the MDPE gas pipes were inserted into augered holes at an angle of  $45^\circ$  to the vertical. These pipes were drilled with twenty-six 5-mm holes at the far end 0.1 m of the tube (the end that went into soil) to deliver  $\text{CO}_2$  gas into the soil at the center of gassing plot at 0.5–0.6 m depth (Figure 3).



Photo 2.  $\text{CO}_2$  gas delivery hardware and software system installed at the ASGAR site. Top left: automated system to control delivery of gas at a targeted rate, bottom left: copper tubes with automatically controlled valves to supply gas, bottom right: copper tubes are connected to yellow plastic pipes which, in turn, carry gas to individual plots and top right: pasture plots at the ASGAR site with measuring tubes and gas delivery pipes in each plots, and in the back ground the cryogenic gas cylinders and port-cabin can be seen.




Fig. 2. Layout of 30 plots within the main experimental block. Light green plots- pasture grass, yellow plots- fallow and agricultural crops, and dark green - turf grass. G = gassed, C = un-gassed control and T = plots used only for testing the gassing system and all the T plots were not used to impose any of the treatments during study period.

The injection rate for all the gassed plots at ASGAR site since CO<sub>2</sub> gassing studies began varied from 1–3 liters min<sup>-1</sup>, which equates to an annual injection rate of around 1–3 tonnes year<sup>-1</sup>. This rate of gas injection is far less than the amount injected at some of the already existing off-shore storage sites. For example, Sleipner west field beneath the North Sea, Norway injects 1 million tonnes of CO<sub>2</sub> year<sup>-1</sup>. Nevertheless, the injection rates selected at the ASGAR project site were constrained by the funding and the logistics put in place as well as for practical purposes.

#### 2.4 Measurements on soil gas concentrations

Plastic tubes of 1 m long with 0.2 mm internal diameter were installed vertically into the soil to a depth of 0.3 m and at different distances from the center of plots on a diagonal transect in all the plots (gassed & un-gassed; Figure 3). The bottom end of each sampling tube (at 0.3 m depth) was sealed and the lower 0.15 m of the tube was drilled with 14 equally spaced holes (4.5 mm diameter). This portion of the tube was covered with fine meshed cloth from inside the tube to prevent outside soil clogging the holes. These holes enabled free diffusion of gas from surrounding soil into the tube so as to attain equilibrium with the soil gas concentration at 0.15 to 0.30 m depth. The top end of the tube was sealed with a plastic on/off valve connectable to a portable GA2000 Landfill Gas Analyzer (Geotechnical Instruments UK Limited) to measure soil CO<sub>2</sub> and O<sub>2</sub> concentrations (in % of total 100% by vol.). In addition, Patil et al. (2010) measured soil surface CO<sub>2</sub> efflux from pasture grass and fallow plots (ppm hr<sup>-1</sup>) using Draeger tubes (Draeger Safety AG & Co., Germany,) placed in a grid spacing of 0.5 m × 0.5 m on the surface to monitor gas diffusive pattern within each plot and its horizontal spread. The Draeger tube system is an established method for measuring and detecting contaminants in the soil, water and air.




Fig. 3. Details of individual experimental plot showing the position of gas delivery pipe (A = above-ground portion, B = below-ground portion), gas measurement spots (C; one at 0.15 m & the other at 0.7 m away from center) and above-ground grass biomass sampling areas only in pasture plot (each of 0.2 × 0.2 m in size). This holds true for un-gassed plots but for the absence of soil gas measuring tube at 0.7 m away from centre of the plot.

## 2.5 Measurements on soil chemistry

West et al. (2009) analyzed soil samples from 0.15–0.30 m and 0.45–0.60 m depth, before and after gassing, for mineralogical (e.g. particle fraction) and geochemical (pH, organic carbon) content and changes due to gassing. Whereas, Patil et al. (2010) collected the soil samples from the top 0.3 m depth, one composite sample from each plot, three times during the gassing period at equal interval, and analyzed for pH and organic matter content. The holes left open after removing augered soil samples were re-filled immediately with local soil and marked to avoid repeated use of the same spots during subsequent sampling periods. Wei et al. (2011), on the other hand, used the top 0.12 m layer soil from the ASGARD site and incubated with CO<sub>2</sub> gas at varied levels of soil moisture under laboratory conditions to study the effect of CO<sub>2</sub> gassing on soil chemical properties and availability of mineral elements.

## 2.6 Measurements on soil inhabiting organisms

To study the effect of CO<sub>2</sub> gassing on soil-inhabiting organisms, earthworm activities were monitored by counting the number of their castings on the surface (Patil et al., 2010). Sarah and Sjoersten (2009) collected soil samples from the turf grass plots from the top 0.1 m depth and at different distance from the center of the plot. They did this on 3 occasions: before gassing began, and at 5 and 10 weeks after gassing. These soil samples were used to determine microbial biomass and microbial activity. West et al. (2009) also looked at soil bacterial population and adenosine triphosphate (ATP) concentrations, the latter as a measure of microbial activity in the soil.

## 2.7 Measurements on vegetation composition and growth

While West et al. (2009) did not collect destructive biomass samples, Sarah and Sjoergersten (2009) collected turf grass above- and below-ground biomass samples to record dry biomass. West et al. (2009) instead monitored botanical composition (% plant species cover) of the pasture plots before and after the gassing period both in gassed and un-gassed plots. Patil et al. (2010) monitored pasture growth by collecting only its above-ground biomass from  $0.2 \times 0.2$  m patch ( $0.04 \text{ m}^2$ ) above soil surface taken at distances of 0.3, 0.6 and 0.9 m from the plot center in all four directions (as shown in Figure 2), making up four samples within each plot from each distance interval. Each time after collecting biomass samples the pasture grass plots were mowed and let the grass grow again, whereas, Sarah and Sjoergersten (2009) did not mow the turf grass. Between April through October 2007, the fallow plots had only bare soil with no vegetation cover and only soil gas concentrations were recorded using GA2000 Landfill Gas Analyzer as described in above paragraphs. On November 1, 2007 all the eight fallow plots were sown with winter bean (*Vicia faba* Cv. Clipper). The seeds were hand dibbled at 45 seeds per  $\text{m}^2$ . First germination count (number of seeds emerged per plot) was recorded on December 3, 2007 and the same was repeated at regular interval until the germination / emergence process was complete or no additional seeds emerged from both gassed and un-gassed plots.

## 2.8 Measurements on pasture grass stress responses

Pasture and turf grass stress symptoms were also monitored by recording visual appearance of grass on the surface (e.g. drying, brown / yellow coloration) both in gassed and un-gassed plots. Furthermore, Patil et al. (2010) monitored physiological stress responses to  $\text{CO}_2$  gassing by measuring moisture content in above-ground grass biomass after each sampling (as a difference between fresh and oven dried biomass) and leaf chlorophyll content. To measure leaf chlorophyll content without any destructive sampling, the SPAD 502 Chlorophyll Meter (Spectrum Technologies Inc., USA) was used. The SPAD meter instantly measures the amount of chlorophyll content in leaves, a key indicator of plant health. The SPAD meter was clamped over pasture grass leaf part for few seconds and the meter displays an indexed chlorophyll content reading (0–99.9). Lower the value means higher stress. A more detailed description of these methodologies and measurements are given elsewhere (Patil et al., 2010; Sarah & Sjoergersten, 2009; West et al., 2009).

## 3. Results and discussion

### 3.1 Soil gas leakage and migration pattern

The first objective of developing ASGARD field gassing facility was to achieve control over  $\text{CO}_2$  gas injection into the soil at a targeted rate all through the study period. This was achieved successfully with the hardware and software logistics installed at the site. Furthermore, artificial gas injection at 0.5–0.6 m depth simulated gas diffusion and migration in the gassed plots (Patil et al., 2010; West et al., 2009). West et al. (2009), who injected  $\text{CO}_2$  gas @  $3 \text{ liters min}^{-1}$  for three months in 2006, reported horizontal migration of gas at a roughly similar rate in all directions in the gassed plots. They also observed that while the injected  $\text{CO}_2$  gas had clearly migrated upwards throughout all the gassed plots, larger lateral movement was recorded at depth. This may have been probably influenced by

the relative permeability of soil and the topographies of the boundaries between plots (West et al., 2009). The injected gas tended to move laterally beneath the soil beyond the boundaries of gassed plots as only the one-third amount of total injected gas was emitted at the surface from the gassed plots (West et al., 2009). However, in 2007-08 when CO<sub>2</sub> gas was injected at lower rates ( 1 liter min<sup>-1</sup>) the lateral spread of diffused gas was relatively small, although the spread was much more in pasture plots than in fallow ones (Figure 4 & 5). This suggests that rate of leakage and the land use type (whether covered with pasture or left fallow) does influence the extent of surface flux and lateral migration of leaking gas. Hoeks (1972) suggests that the sub-surface injected CO<sub>2</sub> gas would migrate isotropically towards soil surface and creates a spherically symmetric gas plume, but that was not the case at ASGARD site (Figure 4 & 5). This may have been caused by the physical disturbance and loosening of soil along the path of inserted plastic pipe (Patil et al., 2010). Therefore, surface CO<sub>2</sub> flux rates recorded and its adverse effect on overlying vegetation were much higher in and around the entry point of plastic pipe (Photo 3). The diffusion and migration pattern of CO<sub>2</sub> gas seems to follow the pattern of CH<sub>4</sub> gas as reported by Adams and Ellis (1960) and Smith et al. (2005).




Fig. 4. Mean CO<sub>2</sub> efflux (ppm hr<sup>-1</sup>) measured at 0.5 m × 0.5 m grid intervals at the soil-air interface in pasture plots using the Draeger tubes and shown in a 3-D contour map (source: Patil et al., 2010).




Fig. 5. Mean CO<sub>2</sub> efflux (ppm hr<sup>-1</sup>) measured at 0.5 m × 0.5 m grid intervals at the soil-air interface in fallow plots using the Draeger tubes and shown in a 3-D contour map (source: Patil et al., 2010).



Photo 3. Vegetation in gassed pasture plots (top two) showing visual stress symptoms compared with non stressed un-gassed pasture plots (middle two), and winter bean emergence and seedlings growth pattern in gassed (bottom right) and un-gassed plots (bottom left).

### 3.2 Effect of land vegetation cover on soil gas concentrations

Gassing significantly increased the soil CO<sub>2</sub> concentrations in both pasture and fallow plots compared with their respective un-gassed pasture and fallow plots (Patil et al., 2010). In the gassed pasture plots, the maximum soil CO<sub>2</sub> concentrations were in the range of 19.5–76.3% with a seasonal mean of 45.7% (by vol. of the total 100%) from the measuring tubes that were 0.15 m from the center of the plots and significantly less, which ranged between 6.1–29.2% with mean of 11.5%, in measurement tubes at 0.7 m from the center as lesser amount of CO<sub>2</sub> migrated laterally below the soil surface. Similar trend was recorded from gassed fallow (bare soil) plots, although the seasonal high, low and mean CO<sub>2</sub> concentrations were on much lower side compared with gassed pasture plots. In contrast, soil CO<sub>2</sub> concentrations recorded from un-gassed plots were very low. From un-gassed pasture and fallow plots seasonal mean CO<sub>2</sub> concentrations of 1.9% and 0.7%, respectively, were recorded, but the soil O<sub>2</sub> concentrations reached as high as 20.8%. This clearly showed

a significant negative relationship ( $r^2=-0.95$ ) between soil CO<sub>2</sub> and O<sub>2</sub> concentrations as soil O<sub>2</sub> gas was displaced by injected CO<sub>2</sub> gas, since the latter one is heavier than former one (Patil et al., 2010). The difference in soil CO<sub>2</sub> concentrations between gassed pasture and gassed fallow plots also suggests that build up of injected CO<sub>2</sub> gas in near-surface and its diffusion / lateral migration are influenced by the land use type. Presence of vegetation tended to hold more gas due to the presence of root biomass, where as bare soil tended to release the leaking gas quickly into the atmosphere. This may have been to do with difference in soil aggregation. Pasture plots were never ploughed or tilled for over 10 years. This may have helped the pasture to develop thickly matted rooting system closing the bigger soil pores while creating larger proportion of micro pores compared with the broken soil (with more of macro pores) in fallow plots due to ploughing and cultivating with agricultural crops. Wei et al. (2011) reported that when soil was exposed to 100% CO<sub>2</sub>, the gas is either absorbed by the soil water present in the pore space or filled the micro pores between the soil particles. This suggests that pasture soil with more of micro pores tends to hold more gas and build up CO<sub>2</sub> concentrations compared with loosely aggregated soil from fallow plots or the soil cultivated with seasonal crops.

When it comes to CO<sub>2</sub> gas migration beyond the plots of gassing, Patil et al. (2010) recorded slightly higher levels of soil CO<sub>2</sub> concentrations in both un-gassed pasture and fallow plots compared with the pasture plots located far away from the main treatment plots (as remote control plots). This could only be possible with sub-surface lateral migration of diffusing CO<sub>2</sub> gas from gassed plots to adjacent un-gassed plots (Patil et al., 2010), which again shows that even with low levels of sub-surface gas leakage (e.g. 1 liter min<sup>-1</sup>) the diffusing gas can move laterally beneath the surface over quite a distance and affect the overlying vegetation and soil ecology. Therefore, if CO<sub>2</sub> leaks were to occur from geological storage sites located >800 m deep via cracks / faults in caprocks the gas would start diffusing upwards and spread in a funnel like shape occupying large area by the time gas reaches soil surface. This leak, if goes unnoticed, could affect large patch of land and its surrounding ecosystem.

### 3.3 Effect CO<sub>2</sub> gassing on pasture composition

The baseline botanical characterization, recorded in 2006 before gassing study was initiated, showed a range of monocotyledonous (e.g. grasses) and dicotyledonous (e.g. dandelions, thistle, plantain, chickweed, mallow, clover) plant species in all the pasture plots. However, three months after gassing the % composition changed only in gassed plots (West et al., 2009). Grass was the dominant species at the center of the plot where the soil CO<sub>2</sub> concentrations were much higher (> 75% of the total 100% by vol.) than at the edges of the plots. Whereas, towards the edges of the gassed plots where CO<sub>2</sub> concentrations were around 45% at 0.2 m depth, grass was the dominant species. In contrast, un-gassed (control) pasture plots had more species composition and higher proportions of minor plant groups than the gassed plots even at the end of the injection period. This suggests that grass species tended to be more tolerant to higher soil CO<sub>2</sub> concentrations than the non-grass or broad leafed species found in this perennial pasture (West et al. 2009). Similar responses were reported from a Mediterranean pasture at a natural CO<sub>2</sub> vent (Latera, Italy) where Beaubien et al. (2008) studied the effects of venting CO<sub>2</sub> gas on the shallow ecosystem.



### 3.4 Effect of CO<sub>2</sub> gassing on pasture grass stress symptoms and growth

Despite being tolerant to elevated CO<sub>2</sub> concentrations, the grass turned yellow or brown and patches of bare earth was exposed as no vegetation (including perennial grass) grew in and around the point of entry of gassing pipeline, the patch where the soil CO<sub>2</sub> concentrations reached as high as 75% at 0.2 m depth (West et al., 2009). Similar visual stress symptoms on pasture and turf grass, respectively, were observed by Patil et al. (2010) and Sarah and Sjoergersten (2009) when CO<sub>2</sub> gas was injected @ 1 liter min<sup>-1</sup> and also when CH<sub>4</sub> gas was injected @ 100 liters hr<sup>-1</sup> in 2002-03 (Smith et al., 2005). While the grass close to the point of entry of gassing pipe turned yellow and died, a small patch of grass in a circular pattern around the dead grass showed much shorter growth, looked yellow and dry, but not dead yet. Whereas, the pasture towards the edges of the plots looked green and less affected as the soil CO<sub>2</sub> gas concentrations recorded towards the edges of the plot were much lower.

The observations on above-ground biomass, chlorophyll content and moisture content of pasture grass followed the same pattern; significantly lower values at the centre and higher values at the edges of gassed plots (Patil et al., 2010). This pattern of effect on overlying vegetation very much followed the pattern of soil CO<sub>2</sub> gas migration below the surface (Figure 4 & 5). Previous studies carried out at sites of naturally occurring CO<sub>2</sub> springs / vents and or volcanic sites have shown that elevated soil CO<sub>2</sub> concentrations reduce plant growth, disrupt plant photosynthesis, inhibit root respiration mainly due to severe dearth of soil O<sub>2</sub> levels, and even kill the vegetation (Cook et al., 1998; Macek et al., 2005; Miglietta et al., 1998; Pfanz et al., 2007; Vartapetian & Jackson, 1997; Vodnik et al., 2006). Previous studies on leaking CH<sub>4</sub> gas have also caused the same stress symptoms on above-ground vegetation as both CO<sub>2</sub> and CH<sub>4</sub> gases displace O<sub>2</sub> gas from the soil (Arthur et al., 1985; Hoeks, 1972; Smith et al., 2005; Smith, 2002) thus depriving the plant roots off O<sub>2</sub> for respiration, which in turn affects other plant functions viz., water and nutrient uptake, evapotranspiration, photosynthesis and ultimately plant growth. In fact, Adamse et al. (1972) suggested that for the proper functioning of a healthy root system, a minimum soil O<sub>2</sub> concentration of 12-14% is required, whereas at ASGARD site in the gassed plots this was not the case. Macek et al. (2005), Vodnik et al. (2006) and Pfanz et al. (2007) studied plant responses in relation to measured soil CO<sub>2</sub> concentrations at a natural CO<sub>2</sub> spring in Slovenia. In their studies soil CO<sub>2</sub> concentrations reached as high as 100% near vents, but atmospheric concentrations barely exceeded ambient concentrations (360-500 ppm) due to fast dispersion of leaking CO<sub>2</sub> by winds on a flat terrain. The study by Macek et al. (2005) considered root respiration in seven plant species and found that only when exposed to very high CO<sub>2</sub> concentrations did it inhibit root respiration, but the effects were highly variable amongst different plant species. This suggests that sensitivity of plants to elevated soil CO<sub>2</sub> concentrations differs with plant species as some species are more sensitive than others. The study by Vodnik et al. (2006) noted that leaf chlorophyll content was negatively correlated with soil CO<sub>2</sub> concentration, whereas Pfanz et al. (2007) noted that plants exposed to high soil and air CO<sub>2</sub> concentrations contained lower levels of nutrients in their vegetative parts and did not flower.

### 3.5 Effect of CO<sub>2</sub> gassing on winter bean crop

Smith et al. (2005) while studying the effect of leaking CH<sub>4</sub> gas on crops (wheat and winter bean) observed that in the plot area with the highest CH<sub>4</sub> concentrations most seeds

germinated, but did not grow further. However, towards edges of plots, an area approximately 0.5 m in diameter, showed reduced growth with yellow leaves. Similarly, in this study significant adverse effect of CH<sub>4</sub> gas on chlorophyll content and leaf area of grass, wheat and bean was also reported. When it comes to CO<sub>2</sub> gassing, significantly lower number of winter bean seeds emerged / grew into seedlings in gassed plots compared with un-gassed plots (Patil et al., 2010). Even a very low level of leakage with 1 liter min<sup>-1</sup> gas injection had lethal effect on bean seeds emergence. Plants are known to be more sensitive to anaerobic conditions especially during early growth stages (Hoeks, 1972) and that might be the reason why CO<sub>2</sub> gassing had lethal effect on winter bean seed emergence and seedling growth compared with the response of well established pasture grass observed by Patil et al. (2010). Schollenberger (1930) while studying the effect of CH<sub>4</sub> leaks on crops noted that leaking gas pipeline killed all the oat seedlings within the range of 1.0–1.3 m while stunting the seedling growth up to 4–5 m away from pipeline and beyond which no injury appeared. Godwin et al. (1990) observed wheat being more tolerant to CH<sub>4</sub> leaks compared with oil seed rape, which again suggests that crops differ in their sensitivity to gas leaks. In fact, this differential sensitivity could be used in early warning system of gas leaks, especially from under-ground pipeline, by growing highly sensitive crops along the path of transportation pipeline.

### 3.6 Effect of CO<sub>2</sub> gassing on soil mineralogy and chemistry

No significant changes in the mineralogical composition of soils from gassed and non-gassed plots were recorded during the 16 weeks injection period (West et al, 2009). However, soil Ca concentrations, in general, decreased, but the largest reductions were recorded in the soil samples collected close to the injection point, which also recorded the highest CO<sub>2</sub> concentrations (West et al, 2009). Wei et al. (2011) suggested that elevated soil CO<sub>2</sub> concentrations enhance weathering of minerals, thus it would be possible to assess the impact of CO<sub>2</sub> leaks on soil mineralogy only if the studies were carried out for longer period of time (in years). With regards to soil pH, soil CO<sub>2</sub> gassing @ 3 liters min<sup>-1</sup> for only 16 weeks reduced the soil pH in gassed plots and the drop in soil pH was drastic near the point of entry of gas delivering pipeline (West et al., 2009). Among the soil layers, soil samples collected from gassed 'A horizon' recorded the largest reduction in pH (by 0.5 units) after only 16 weeks of gassing (West et al, 2009). Similarly, Patil et al. (2010) also recorded lower soil pH both in gassed pasture and fallow plots in comparison with their respective un-gassed control plots. This could be attributed to acidification of soil water due to dissolution of leaking CO<sub>2</sub> gas (Celia et al., 2002) and was corroborated by the findings of Wei et al. (2011), where the latter reported that absorption of CO<sub>2</sub> gas would occur either by reacting with soil pore water or by filling the pore space, thus increased soil moisture increases soil CO<sub>2</sub> build up in the soil. This process further leads to the formation of H<sub>2</sub>CO<sub>3</sub> which, in turn, lowers the pH of soil solution. Soils in general have a buffering capacity and when pH of soil solution was lowered in the presence of absorbed CO<sub>2</sub>, clay minerals start to weather to neutralize the lowering pH by releasing minerals (to exchange with H<sup>+</sup> ions) leading to an increase in the CaCl<sub>2</sub>-exchangeable concentration of Al. Thus, increase in uptake of CO<sub>2</sub> by the soil solution leads to increased Al mobilization in moist soil (Wei et al, 2011). However, Al concentrations reported in their study were much lower than plant tolerance limits, which range between 40 μM and 60 μM depending on species (Poschenrieder et al, 2008; Taylor et al, 1998). In addition to Al, Wei et al. (2011) also reported increase in CaCl<sub>2</sub>-

exchangeable concentrations for Mg, K, Ti, V, Cr, Mn, Fe, Co, Cu, Rb, Sr, Mo, Cs, Ba, Pb, Th and U, while the metal concentrations for Zn and Cd decreased. Whereas, total organic carbon (TOC) concentrations in the 'A horizon' increased in both gassed and non-gassed plots by the end of the summer growing season, but at the injection depth TOC concentrations were found to be on lower side in the gassed plot compared with the un-gassed plots (West et al., 2009). This suggests that elevated soil CO<sub>2</sub> concentrations lower soil pH and TOC, although different soil horizons responded differently depending on their buffering capacity to changes in pH and chemical components. However, these gassing studies were carried out for a very short period (few months) and variations in soil carbon observed in these studies show only the rough indications of changes in soil chemistry which, in this case, might have been of temporary in nature influenced by factors not controlled during the study period (e.g. seasonal temperature, above-ground vegetation and other climatic conditions). Therefore, further investigation by continuing gassing over long period of time needs to be undertaken.

### 3.7 Effect of CO<sub>2</sub> gassing on soil inhabiting organisms

CO<sub>2</sub> gassing significantly increased the earth worm casts in gassed pasture plots compared with un-gassed pasture plots, but no earth worm activities were noticed on both gassed and un-gassed fallow plots (Patil et al., 2010). Smith et al. (2005) also reported higher worm casts in grass plots injected with natural gas. While the higher earth worm casting on gassed pasture plots indicate increased activities, but surprisingly enough the absence of earth worm casting on both gassed and un-gassed fallow plots begs question which the authors fail to clarify. Therefore, this needs to be looked at in detail to understand the causation. When it comes to soil bacterial population CO<sub>2</sub> gassing drastically reduced their number in gassed plots. The ATP concentrations, a measure of microbiological activity, were also reported to be below detection limits at the center of the gassed plots where CO<sub>2</sub> concentrations reached maximum of 87% at 0.15–0.30 m depth (West et al, 2009).

Sarah and Sjögersten (2009) observed a tendency for reduced respiration in the soil exposed to elevated CO<sub>2</sub> concentrations, not significant though, compared with the soil in un-gassed plots within a period of 10 weeks. Soil moisture content seemed to have negative significant influence on microbial respiration. This negative effect of higher soil moisture on soil respiration, Sarah and Sjögersten (2009) report, was not anticipated as increased soil moisture generally results in increased microbial respiration up to the point of anoxia (Wardle and Parkinson 1990; Davidson et al. 2000). One possible explanation could be that higher soil water content reduced the O<sub>2</sub> supply (anoxic condition) while increasing the share of CO<sub>2</sub> concentrations in soil pores owing to the latter's leakage from the center of the plot. Such a soil environment would inhibit microbial respiration in the same way when soil water content reaches saturation (Davidson et al. 2000). Sarah and Sjögersten (2009) also looked at carbon source utilization rate as an estimate of change in microbial community metabolism, which however, did not differ between the soil from gassed and un-gassed plots in a 72 hr incubation study.

However, findings from the atmospheric CO<sub>2</sub> concentrations enrichment studies (not the soil concentrations) suggest that longer period of time is required to see any noticeable changes in microbial communities. For example, Griffiths et al. (1998) recorded no difference in soil microbial communities in a less than a month long study whereas, Grayston et al.

(1998) observed changes in bacterial community in a study carried out for four long years. Therefore, to understand the actual and long lasting effects on soil chemistry, organic matter, microbial population / their diversity, the gassing needs to be continued for longer period.

#### 4. Conclusions

The ASGARD site developed at the University of Nottingham, UK successfully injected CO<sub>2</sub> gas into the soil at a targeted rate, simulated build up of soil CO<sub>2</sub> concentrations near soil surface and enabled measurement of soil concentrations and its effect on vegetation and soil biogeochemistry. The CO<sub>2</sub> gassing studies from the ASGARD site, despite being carried out for short period of time, showed the potential ecological risks of CO<sub>2</sub> leakage from geological storage sites on agro-ecosystems. Surprisingly enough, even the low levels of gas leakage with 1–3 liters of CO<sub>2</sub> gas injection min<sup>-1</sup> significantly increased the soil CO<sub>2</sub> concentrations in a very short period of time by displacing the soil O<sub>2</sub> and adversely affected the growth of pasture and turf grass, severely hindered the germination and establishment of winter bean crop, lowered soil pH, TOC and microbial population while increasing the activities of earthworms. These studies also showed that different plant species have different tolerance capacity to soil CO<sub>2</sub> concentrations. Monocotyledonous species (e.g. grass types) tended to be more tolerant compared with dicotyledonous species (e.g. beans & broad leaf species). Therefore, some of the more sensitive plant species (e.g. non-grassy species) could be used to grow along the path of CCS transportation pipelines in early warning system to detect gas leaks.

Another limitation of studies carried out at the ASGARD site was that the CO<sub>2</sub> gas was injected at a very low rate (1–3 liters min<sup>-1</sup>) and at a shallow depth (0.5–0.6 m) compared with CCS schemes where amount of gas injected each minutes would be in tonnes and at > 800 m deep. Therefore, while the ASGARD site based studies increased our understanding of the effects of below ground CO<sub>2</sub> leaks on agro-environment, long-term studies to evaluate the potential long-term consequences on ecosystem are needed to be carried out on larger scale for decision making and management. In CCS schemes the upwards movement of gas from the deep storage sites, possibly via cracks and faults in the caprocks, and its further diffusion and migration before reaching the soil surface would naturally enable the gas to spread on much larger area. Moreover, under such a situation factors like quality of selected site, amount of gas injected, the state and pressure under which it is held beneath the caprocks, geochemistry of the path of gas movement and issues related to hydrology needs to be studied in detail in addition to the impacts on overlying vegetation and environment. Therefore, further gassing studies needs to be carried out on larger scale and continued for longer period to better understand the long-term impacts.

#### 5. Acknowledgments

The fellowship and research funding awarded to the author of this chapter by the Royal Society, London International Incoming Fellowship program 2006/R2 is duly acknowledged. The author wish to acknowledge the host institution project leaders Prof. Jeremy J. Colls (now retired while writing this chapter) and Prof. Michael D. Steven, University of Nottingham for initiating and leading the ASGARD project site. Thanks are

also due to the British Geological Survey, UK for extending logistics, resources and technical support while developing the ASGARD site and for undertaking baseline studies to start with.

## 6. References

- Adams, R.J., & Ellis, R.J. (1960). Some physical and chemical changes in the soil brought about by saturation with natural gas. *Soil Science Society of America Proceedings* 24, pp. 41–44.
- Adamse, A.D., Hoeks, J., DeBont, J.A.M., & van Hessel, J.F., (1972). Microbial activities in soil near natural gas leaks. *Archive Microbiology* 83, pp. 32–51.
- Arthur, J., Leone, I., & Flower, F. (1985). The response of tomato plants to simulated landfill gas mixtures. *Journal of Environmental Science and Health* 20, pp. 913–925.
- Bachu, S. (2000). Sequestration of CO<sub>2</sub> in geologic media: approaches for site selection in response to climate change. *Energy Conversion and Management* 41, pp. 953–970.
- Baggott, S.L., Davidson, I., Dore, C., Doogwin, J., Milne, R., Murrels, T.P., Rose, M., Watterson, J.D., & Underwood, B. (2003). UK Greenhouse Gas Inventory, 1990 to 2001. *Annual report for submission under the Framework Convention on Climate Change*, <http://www.naei.org.uk/reports.php> (accessed February 10, 2008)
- Beaubien, S. E., Ciotoli, G., Coombs, P., Dictor, M.C., Kruger, M., Lombardi, S., Pearce J. M., & West J. M. (2008). The impact of a naturally-occurring CO<sub>2</sub> gas vent on the shallow ecosystem and soil chemistry of a Mediterranean pasture (Latera, Italy). *Journal of Greenhouse Gas Control* 2, pp. 373–387.
- Benson, S.M. (2005). Carbon dioxide capture for storage in deep geological formations – results from the CO<sub>2</sub> capture project. In *Geological Storage of Carbon Dioxide with Monitoring and Verification* Vol. 2, Benson, S.M. (Ed.), pp. 654, Elsevier Publishing, UK.
- Bergfeld, D., Evans, W.C., Howle, J.F., Farrar, C.D. (2006). Carbon dioxide emissions from vegetation-kill zones around the resurgent dome of Long Valley caldera, eastern California, USA. *Journal of Volcanology and Geothermal Research* 152, pp. 140–56.
- Billett, M.F., Parker-Jervis, F., FitzPatrick, E.A., & Cresser, M.S. (1990). Forest soil chemical changes between 1949/50 and 1987. *Journal of Soil Science*, 41, pp.133–145.
- Biondi, F., & Fessenden, J. E. (1999). Response of lodgepole pine growth to CO<sub>2</sub> degassing at Mammoth Mountain, California. *Ecology* 80(7), pp. 2420–2426.
- Blake, L., Goulding, K.W.T., Mott, C.J.B., & Poulton, P.R. (2000). Temporal changes in chemical properties of air-dried stored soils and their interpretation for long-term experiments. *European Journal of Soil Science*, 51 (2), pp. 345–353.
- Celia, M.A., Peters, C.A., & Bachu, S. (2002). Geological storage of CO<sub>2</sub>: leakage pathways and environmental risks. *American Geophysical Union, Spring Meeting 2002*, abstract #GC32A-03 in SAO/NASA Astrophysics Data System.
- Cook, A.D., Tissue, D.T., Roberts, S.W., Oechel, W.C. (1998). Effects of long-term elevated CO<sub>2</sub> from natural CO<sub>2</sub> springs on *Nardus stricta*: photosynthesis, biochemistry, growth and phenology. *Plant, Cell and Environment* 21, pp. 417–425.

- Davidson, E.A., Verchot, L.V., Cattanio, J.H., Ackerman, I.L., & Carvalho, J.E.M. (2000). Effects of soil water content on soil respiration in forests and cattle pastures in eastern Amazonia. *Biogeochemistry* 48, pp. 53–69.
- DTI, Department of Trade and Industry, UK, (2002). The feasibility of carbon dioxide capture and storage in the UK. *In project definition and scoping paper*. p. 12, DTI, London, UK.
- DTI, Department of Trade and Industry, UK, (2005). Geoscientific monitoring of the Weyburn CO<sub>2</sub> flood. A multidisciplinary approach to CO<sub>2</sub> monitoring. *Project summary 281 to Cleaner fossil fuels program*, DTI, London, UK.
- Ford, R. (2006). The geology of the ASGARD site, Sutton Bonington. *British Geological Survey Internal Report IR/06/049R*.
- Freund, P., Gale, J., Davison J., & Riley, N. (2003). Capture and geological storage of carbon dioxide – a status report on the technology. *IEA Greenhouse Gas R & D Program*. COAL R223, British Geological Survey, London, UK.
- Gahrooe, F.R., (1998). Impacts of elevated atmospheric CO<sub>2</sub> on litter quality, litter decomposability, and nitrogen turnover rate of two oak species in a Mediterranean forest ecosystem. *Global Change Biology* 4, pp. 667–677.
- Gelrach, T., Doukas, M., McGee, K., & Lessler, R. (1998). Three-year decline of magmatic CO<sub>2</sub> emissions from soils of a Mammoth Mountain tree kill: Horseshoe Lake, CA, 1995–1997. *Geophysical Research Letters*, 25, pp. 1947–1950.
- Gerlach, T.M., Douglas, M.P., McGee, K.A., & Kessler, R. (2001). Soil efflux and total emission rates of magmatic CO<sub>2</sub> at the Horseshoe Lake tree kill, Mammoth Mountain, California, 1995–1997. *Chemical Geology* 177, pp. 101–116.
- Godwin, R., Abouguendia, Z., & Thorpe, J. (1990). Response of soils and plants to natural gas migration from two wells in the Lloydminster area, *Saskatchewan Research Council Report (No. E- 2510-3-E-90)*, pp. 85.
- Goulding, K.W.T., Bailey, N.J., Bradbury, N.J., Hargreaves, P., Howe, M., Murphy, D.V., Poulton, P. R., & Willison, T.W. (1998). Nitrogen Deposition and its Contribution to Nitrogen Cycling and Associated Soil Processes. *New Phytologist*, 139 (1), pp. 49–58.
- Grayston, S.J., Campbell, C.D., Lutze, J.L., & Gifford, R.M. (1998). Impact of elevated CO<sub>2</sub> on the metabolic diversity of microbial communities in N-limited grass swards. *Plant Soil* 203, pp. 289–300.
- Griffiths, B.S., Ritz, K., Ebbelwhite, N., Paterson, E., & Killham, K. (1998). Ryegrass rhizosphere microbial community structure under elevated carbon dioxide concentrations, with observation on wheat rhizosphere. *Soil Biology & Biochemistry* 30, pp. 315–321.
- Heinrich, J.J., Herzog, H.J., & Reiner, D.M. (2003). Environmental assessment of geologic storage of CO<sub>2</sub>. Laboratory for Energy and the Environment, MIT, presented at the *Second National Conference on Carbon Sequestration*, Washington D.C., May 5–8, 2003.
- Hoeks, J. (1972). Changes in the composition of soil air near leaks in natural gas mains. *Soil Science* 113, pp. 46–54.
- IPCC, Intergovernmental Panel on Climate Change, (2005). Carbon Dioxide Capture and Storage. Special report prepared by Working Group III of the Intergovernmental Panel on Climate Change, Metz, B., Davidson, O., de Coninck, H.C., Loos, M., &

- Meyer, L.A. (Eds), p. 442, Cambridge, United Kingdom: Cambridge University Press; and New York, NY, USA: Cambridge University Press.
- IWR, Institute for Renewable Energy, (2009). Climate protection just a lot of hot air – Global CO<sub>2</sub> emissions at record levels, Press Release, August 10, 2009, Internationales Wirtschaftsforum Regenerative Energien, Muenster, Germany (<http://www.iwrpressedienst.de/iwr/Global-CO2-emissions-2008-renewable-energy-investment-plan.pdf>; as on July 5, 2010)
- Jossi, M., Fromin, N., Tarnawski, S., Kohler, F., Gillet, F., Aragno, M. & Hamelin, J. (2006). How elevated pCO<sub>2</sub> modifies total and metabolically active bacterial communities in the Rhizosphere of two perennial grasses grown under field conditions. *FEMS Microbiology Ecology*, 55, pp. 339–350.
- Klusman, R.W. (2003). Evaluation of leakage potential from a carbon dioxide EOR/sequestration project. *Energy Conversion and Management*, 44, pp. 1921–1940.
- Kruger, M., West, J., Frerichs, J., Oppermann, B., Dictor, M.C., Joulain, C., Jones, D., Coombs, P., Green, K., Pearce, J., May, F., & Moller, I. (2009). Ecosystem effects of elevated CO<sub>2</sub> concentrations on microbial populations at a terrestrial CO<sub>2</sub> vent at Laacher See, Germany. *GHGT-9, Energy Procedia* 1, pp. 1933-1939.
- Lewicki, J.L., Hilley, G.E., & Oldenburg, C.M. (2005). An improved strategy to detect CO<sub>2</sub> leakage for verification of geologic carbon sequestration, *Geophysical Research Letters*, 32 (19), L19403.
- Macek, I., Pfan, H., Francetic, V., Batic, F., & Vodnik, D. (2005). Root respiration response to high CO<sub>2</sub> concentrations in plants from natural CO<sub>2</sub> springs. *Environmental and Experimental Botany* 54, pp. 90–99.
- Miglietta, F., Berrardini, I., Raschi, A., Korner, C., & Vaccari, F.P. (1998). Isotope discrimination and photosynthesis of vegetation growing in the Bossoleto CO<sub>2</sub> spring. *Chemosphere* 36, pp. 771–776.
- Pacala, S., & Socolow, R. (2004). Stabilization Wedges: Solving the Climate Problem for the Next 50 Years with Current Technologies. *Science*, 305(5686), pp. 968–972.
- Patil, R.H., Colls, J.J., & Steven, M.D. (2010). Effects of CO<sub>2</sub> gas leaks from geological storage sites on agro-ecosystems. *Energy*, 35, 4587–4591.
- Perry, K. (2005). Natural gas storage industry experience and technology: Potential application to CO<sub>2</sub> – do photosynthetic and productivity data from FACE experiments support early predictions? *New Phytologist*, 162, pp. 253–280.
- Pfan, H., Vodnik, D., Whittman, C., Aschan, G., Batic, F., Turk, B., & Macek, I. (2007). Photosynthetic performance (CO<sub>2</sub> compensation point, carboxylation efficiency, and net photosynthesis) of timothy grass (*Phleum pratense* L.) is affected by elevated carbon dioxide in post-volcanic mofette areas. *Environmental and Experimental Botany* 61, 41–48.
- Poschenrieder, C., Gunse, B., Corrales, I., & Barcelo, J. (2008). A glance into aluminum toxicity and resistance in plants. *Science of the Total Environment* 400, pp. 356–368.
- Sarah, P., & Sjogersten, S. (2009). Effects of below ground CO<sub>2</sub> emissions on plant and microbial communities. *Plant Soil*, 325, pp. 197–205.
- Scholenberger, C. (1930). Effect of leaking natural gas upon the soil. *Soil Science*, 29, pp. 260–266.


- Smith, K.L., Steven, M.D., & Colls, J.J. (2005). Use of hyper spectral derivative ratios in the red-edge region to identify plant stress responses to gas leaks. *Remote Sensing of Environment* 92, 207–217.
- Smith, K.L., (2002). Remote sensing of leaf responses to leaking underground natural gas. PhD Thesis, p. 225, University of Nottingham, UK.
- Sorey, M.L., Farrar, C.D., Gerlach, T.M., McGee, K.A., Evans, W.C., Colvard, E.M., Hill, D.P., Bailey, R.A., Rogie, J.D., Hendley, J.W., II, & Stauffer, P.H. (2000). Invisible CO<sub>2</sub> gas killing trees at Mammoth Mountain, California: *U. S. Geological Survey Fact Sheet FS 0172-96*, p. 2.
- Sowerby, A., Ball, A.S., Gray, T.G.R., Newton, P.C.D., & Clark, H. (2000). Elevated atmospheric CO<sub>2</sub> concentration from a natural soda spring affects the initial mineralization rates of naturally senesced C<sub>3</sub> and C<sub>4</sub> leaf litter. *Soil Biology & Biochemistry*, 32, pp. 1323–1327.
- Stenhouse, M., Arthur, R., & Zhou, W. (2009). Assessing Environmental Impacts from Geological CO<sub>2</sub> Storage. *Energy Procedia*, 1(1), pp. 1895–1902.
- Stephens, J.C., & Hering, J.G. (2002). Comparative characterization of volcanic ash soils exposed to decade-long elevated carbon dioxide concentrations at Mammoth Mountain, California, *Chemical Geology*, 186, pp. 301–313.
- Stevens, S., Fox, C., & Melzer, L. (2000). McElmo dome and St. Johns natural CO<sub>2</sub> deposits: Analogs for geologic sequestration. In *proceedings of the Fifth International Conference on Greenhouse Gas Control Technologies (GHGT5)*, Williams, D.J., Durie, R.A., McMullan, P., Paulson, C.A.J., & Smith, A.Y. (Eds), pp. 328–333, CSIRO Publishing, Collingwood, VIC, Australia.
- Taylor, G.J., Blamey, F.P.C., & Edwards, D.G. (1998). Antagonistic and synergistic interactions between aluminum and manganese on growth of *Vigna unguiculata* at low ionic strength. *Physiologia Plantarum*, 104(2), pp. 183–194.
- Tian, S., Fan, Q., Xu, Y., Wang, Y., & Jiang, A. (2001). Evaluation of the use of high CO<sub>2</sub> concentrations and cold storage to control of *Monilinia fructicola* on sweet cherries. *Postharvest Biology and Technology*, 22(1), pp. 53–60.
- Van der Zwaan, B., & Smekens, K. (2009). CO<sub>2</sub> capture and storage with leakage in an energy-climate model. *Environmental Modeling & Assessment*, 14(2), pp. 135–148.
- Van Gardingen, P.R., Grace, J., Harkness, D.D., Miglietta, F., & Raschi, A. (1995). Carbon dioxide emissions at an Italian mineral spring: measurements of average CO<sub>2</sub> concentration and air temperature. *Agricultural & Forest Meteorology*, 73, pp.17–27.
- Vartapetian, B.B., & Jackson, M.B. (1997). Plant adaptations to anaerobic stress. *Annals of Botany (London)* 79(Suppl. A), pp.3–20.
- Vodnik, D., Kastelec, D., Pfanž H., & Macek, I. B. T. (2006). Small-scale spatial variation in soil CO<sub>2</sub> concentration in a natural carbon dioxide spring and some related plant responses. *Geoderma*. 133, pp. 309–319.
- Wardle, D.A., & Parkinson, D. (1990). Interactions between microclimatic variables and the soil microbial biomass. *Biology and Fertility of Soils*, 9, pp. 273–280.
- Wei, Y., Maroto-Valer, M., & Steven, M.D. (2011). Environmental consequences of potential leaks of CO<sub>2</sub> in soil. *Energy Procedia*, 4, pp. 3224–3230.



West, J.M., Pearce, J.M., Coombs, P., Ford, J.R., Scheib, C., Colls, J.J., Smith, K.L., & Steven, M.D. (2009). The impact of controlled injection of CO<sub>2</sub> on the soil ecosystem and chemistry of an English lowland pasture. *Energy Procedia*, 1, pp. 1863–1870.

IntechOpen

IntechOpen



## **Diversity of Ecosystems**

Edited by Prof. Mahamane Ali

ISBN 978-953-51-0572-5

Hard cover, 484 pages

**Publisher** InTech

**Published online** 27, April, 2012

**Published in print edition** April, 2012

The ecosystems present a great diversity worldwide and use various functionalities according to ecologic regions. In this new context of variability and climatic changes, these ecosystems undergo notable modifications amplified by domestic uses of which it was subjected to. Indeed the ecosystems render diverse services to humanity from their composition and structure but the tolerable levels are unknown. The preservation of these ecosystemic services needs a clear understanding of their complexity. The role of research is not only to characterise the ecosystems but also to clearly define the tolerable usage levels. Their characterisation proves to be important not only for the local populations that use it but also for the conservation of biodiversity. Hence, the measurement, management and protection of ecosystems need innovative and diverse methods. For all these reasons, the aim of this book is to bring out a general view on the function of ecosystems, modelling, sampling strategies, invading species, the response of organisms to modifications, the carbon dynamics, the mathematical models and theories that can be applied in diverse conditions.

### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Raveendra H. Patil (2012). Impacts of Carbon Dioxide Gas Leaks from Geological Storage Sites on Soil Ecology and Above-Ground Vegetation, Diversity of Ecosystems, Prof. Mahamane Ali (Ed.), ISBN: 978-953-51-0572-5, InTech, Available from: <http://www.intechopen.com/books/diversity-of-ecosystems/impacts-of-carbon-dioxide-gas-leaks-from-geological-storage-sites-on-soil-ecology-and-terrestrial-ec>

**INTECH**  
open science | open minds

#### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166  
[www.intechopen.com](http://www.intechopen.com)

#### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen