

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,600

Open access books available

138,000

International authors and editors

175M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Molecular Genetics and its Applications in Forensic Sciences

Raluca Dumache, Veronica Ciocan,
Camelia-Oana Muresan and Alexandra Enache

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/63530>

Abstract

The way to medico legal identification was open at the end of the twenty-first century by the “digital fingerprinting” represented by the multifactorial phenotypical trait, determined by both polygenic and environmental factors, followed by group-specific antigens, or with specificity for blood and tissue, and ending with the DNA molecule in use today. Because of this aspect, the framework of modern forensic medicine includes a new field, that of forensic genetics, that mostly involves working with investigations that have human genotype identification as a goal.

Keywords: forensic genetics, human identification, DNA genotype, polymerase chain reaction (PCR), short tandem repeats (STR), single-nucleotide polymorphisms (SNPs)

1. Introduction

Forensic identification is a universal method used to establish the truth in the process of forensic investigation. The evidence includes, among other components, the identification but without being mistaken for it or being reduced to just this.

Both medico-legal and criminalities identification are integrative parts of forensic identification, having probative value. The value of an identification method resides in the expert's ability to compare traces left at the crime scene with traces found on other materials such as reference evidence. Through this method, one can compare: traces of blood left at the crime scene with those found on a suspects' clothes and with samples from the victim. Furthermore,

the rifling can be compared with fingerprints left on the weapon and with the rifling of other weapons.

Medico-legal identification is based on intrinsic scientific methods or scientific methods absorbed from other sciences, usually bio-medical sciences. Scientific progress from the last 30–40 years has highlighted and continues to highlight the role of the specialists in identification. Their role proves its importance in cases that have to do with civil and criminal law, family law, as well as in cases of catastrophes with numerous victims (natural, accidents, wars, terrorist attacks).

Together with the discovery by Mullis in 1983 of the polymerized chain reaction (PCR), Sir Alex Jeffreys will introduce in the field of forensic genetics this technique by studying a set of DNA fragments that proved to have unique characteristics, which were nonrecurring and inherent for each individual, the only exception being monozygotic twins. Alec Jeffreys named these reaction products “genetic fingerprints”.

In this chapter, we will present the utility of molecular genetics techniques in the case of forensic identification, as well as in criminalistics cases.

2. Short history of forensic genetics

In 1880, British anthropologist Sir Francis Galton published the first studies on digital fingerprinting as an identification method of a certain person.

Another important step in forensic identification was discovering the protein polymorphism of the AB0 blood groups by the Austrian doctor Landsteiner at the beginning of the twentieth century.

After year 1950, in the forensic serology, laboratories were tested a number of blood and tissue antigens, culminating with researching the major histocompatibility complex (HLA).

Sir Alec Jeffreys introduced for the first time the DNA fingerprinting in the field of forensic genetics, proving that some regions from the DNA contain repetitive sequences which are variable among individuals [1]. He was the first to prove the importance of using genetic fingerprinting in the case of forensic personal identification (crimes, filiation, consanguinity, sexual abuse, immigration).

Due to this discovery, the first case of forensic genetics could be solved using the DNA analysis [2]. After murdering of two girls in 1983 and 1986, the police organized the blood sample collection from 5000 men living in the area where the murder took place, and finally found the killer by his DNA profile [3].

In 1983, Kary Mullis developed the polymerase chain reaction (PCR) technique which opened new ways in DNA analysis in the forensic genetics. Thus, from each biological human trace or micro-trace containing nucleated cells, the DNA is extracted to be subjected to amplification reactions. The method is an enzymatic process by which regions of the DNA are replicated

(multiplied) 28–34 times, generating about one billion (10^9) copies. This technique highlights the number of repetitions of the base unit and turns them into alpha-numeric values, known as genetic profiles.

Genetic analysis of a very small number of nucleated cells, namely a very small amount of biological material, is made by a different approach from the usual situations and aims to generate, through the reaction of amplification, the sufficient quantity of copies of DNA fragments to obtain an exploitable genetic profile. This method is applicable to biological samples with a DNA matrix containing <50 pg., to biological micro-traces, to biological samples that are in an advanced stage of decay, with single source or multiple sources.

For the last two decades, the results of the DNA analysis have been accepted as evidence in the court in many countries [4]. Since February 1992 when The European Council issued the **Recommendation No. 92**, regarding the use of the DNA analysis in the criminal justice the DNA test is accepted in the Court [5].

3. Biological sample collection

3.1. Samples

Biological samples consisting of nucleated cells are essential for forensic genetic profiling [6].

Biological samples belonging to this category are as follows:

- Liquid blood or dry deposited on supports;
- Liquid semen or dry deposited on supports;
- Various biological secretions (saliva, semen) or mixtures of secretions originating after sexual acts;
- Hard tissues (bone, teeth);
- Hairs with follicles (“root”);
- Slides and cytological smears.

3.2. Sample collection, storage, and characterization for DNA analysis

Nowadays, whole blood is considered to be the most widely used source of DNA. It is harvested on anticoagulant EDTA (ethylenediaminetetraacetic acid). It can be conserved at 4°C for a maximum of 5–7 days; after this time, the DNA sample being kept at -20°C for a few weeks, or at -80°C for longer periods of time [7].

In the case of blood harvested during autopsy, it does not need anticoagulant and it can be harvested from the following organs:

- Heart chambers;

- Skeletal muscles or heart muscle;
- Peripheral vessels;
- Long bones;
- Other tissue types.

Apart from whole blood, epithelial cells from oral mucosa as well as hair are considered to be important sources for DNA analysis, being frequently used. In the case of epithelial cells, they are harvested with a sterile bud or brush. After harvesting, they are wrapped in a paper envelope and put to dry. Hair is taken by pulling, 5–10 with the root being enough for the analysis.

The hair is wrapped in paper envelopes or plastic wrapping and is kept in a dry environment, at room temperature [8].

In comparison with the biological samples mentioned earlier, in the case of tissue, muscles, organs, or skin, the harvested volume needs to be between 2 and 4 cm. Right after harvesting, the DNA extraction and isolation follows because both its quantity and quality decrease in time and depend on the storage conditions.

3.3. Characterization of the DNA analysis

DNA analysis is a complex process which consists in the following phases:

- a. DNA extraction;
- b. DNA quantification;
- c. DNA amplification;
- d. Detection of the DNA amplified products.

4. Methods used in forensic for human identification

4.1. Autosomal STR profiling

In forensic DNA typing, short tandem repeats (STR) or microsatellites are the most frequently genotyped in order to distinguish between individuals, to tie an individual to a crime or to exonerate the innocent. STRs were discovered in the 1980s [9] and since then, they are the “gold standard” in human identification in forensic investigations. They consist of mono-, di-, tri-, tetra-, penta-, and hexa-nucleotide repeat. An individual can be either homozygous (with the same number of repeats) or heterozygous (different number of repeats) in a certain locus. Tetra-nucleotide repeats are used for genotyping in forensic DNA analysis [10].

STR profiles obtained from biological samples found at crime scenes are compared with other profiles of known suspects and are identified by police or are included in a national forensic DNA database [11]. Also, the STR profiling is used in paternity/maternity testing, disaster

victim identification (DVI), rape perpetrators identification, and kinship testing [12]. Due to the STR profiles, in numerous cases, the persons have been excluded from involvement in crimes and have been exonerate.

A main advantage of the STRs markers consists in the fact that they can test in a rapid, simple, and simultaneously way more than 10 STR loci by multiplexing [13]. Due to this characteristic, it offers an increased degree in the identification of different biological samples.

The tetra-nucleotide and penta-nucleotide systems are included in the multiplex analysis kits because they can provide results with an increased index of exclusion. The nomenclature of the STR loci and the allelic variants was established in 1993 by the DNA Commission of the International Society of Forensic Genetics (ISFG). Except their usefulness in forensic DNA analysis, STRs became used in medical genetic research, because it has been demonstrated that the trinucleotide STR loci is associated with some genetic disorders. The DNA profile refers to the genotype (the number of repeats found in each allele of the analyzed STR marker) of a suspect, victim, or crime scene sample.

In the development of STR typing system, in 1997, the Federal Bureau of Investigations (FBI) introduced the database named CODIS (Combined DNA Index System) that included 13 autosomal loci and the amelogenin sex test [14]. These loci are highly polymorphic, localized in non-coding regions which are on different chromosomes. As an improvement in their efficiency, the new multiplexes that amplify 16 loci or more, in a single reaction (including amelogenin too), have been introduced in the last years [15]. The most common STR kits used in the forensic laboratories for the identification are manufactured by three companies: Life Technologies, Promega and Qiagen.

The forensic DNA analysis is made through multiplex PCR amplification of 10–16 STRs or more, followed by automated sequencing equipment, such as capillary electrophoresis (CE) [16].

The STR-based forensic DNA analysis has been well accepted by population and professionals, as an important tool in human identifications and in the criminal justice.

Other sources of genetic variations that have been demonstrated to present more specialized uses in forensic identification are as follows: autosomal SNPs, the markers on the Y chromosome and mitochondrial DNA (mtDNA).

4.2. Analysis of the Y-chromosome

In forensic medicine, the Y-chromosome has only one useful property: It is present only in males. Thus, in crime cases, the investigators expect to find Y-chromosome at the scene. Also, when talking about male–female ratio in body fluid mixtures, such as rapes, by analyzing the Y-STR component, the investigators can obtain more information regarding the male component. It is well known that vasectomized or azoospermic rapists do not leave sperm traces, and it is impossible to find spermatozoa on the microscopic examination. In such cases, the Y-STR profiling is very useful, offering information regarding the identity of the rapist [17].

4.3. Analysis of mitochondrial DNA (mtDNA)

Mitochondrial DNA (mtDNA) is inherited on the maternal line; thus, all the members of a matrilineal share the same haplotype. Its advantage relies on the fact that it has a number of 200–1700 copies per cell and has an increased probability of survival compared to the nuclear DNA. Therefore, the forensic applications for mtDNA include analysis of biological samples that are old or severely degraded, and analysis of biological samples which contain a low amount of DNA (e.g., hair shafts). MtDNA has been used to identify the Tsar Nicholas II and his brother Georgij Romanov [18].

4.4. Autosomal single-nucleotide polymorphisms (SNP) typing

Single-nucleotide polymorphisms (SNPs) have a lower heterozygosity when compared to STRs. The advantage of SNP typing consists in the fact that the DNA template size can be as large as 50 bp, compared to the STRs which need the DNA template size of 300 bp, to obtain a good STR profiling [19].

Due to this, the SNPs became important tools in analyzing degraded samples. Thus in 2001, SNP typing was used in identifying the victims from the World Trade Center disaster [20, 21]. Related to the use of SNP analysis in degraded biological samples, the European Network of Forensic Science Institutes (ENFSI) and the US FBI Scientific Working Group on DNA Methods (SWGDM) are working on recommendations regarding the standardization of the SNPs [22].

5. Impact of genetic identification on justice

At the present time, genetic testing using DNA has wide applicability to the field of justice, this method being used for the following:

- Identification of suspects and confirmation of guilt;
- Exculpation of innocent parties;
- Discovering of persons who commit crimes, or serial killers;
- Researching biological filiation;
- Establishing consanguinity in more complex cases;
- Establishing biological bloodline relations that serve in finding the historical truth (the Romanov case, the case of Louis XVII of France, the Nazi doctor Joseph Mengele)
- Identification of victims of terrorist attacks or natural catastrophes;

- Identification of certain probes for medical testing (tissue for histology testing, blood alcohol content testing, other toxicological tests).

6. Emerging biomarkers in forensic identification

Forensic genetics will continue to develop and improve its methods, due to the advances in the technical development. The key element in forensic protocols is to identify the origin of biological traces found at the scene. In the last 5 years, some studies have demonstrated that messenger RNA (mRNA) can be useful in forensic identification. Zubakov et al. [23] have identified some mRNA stable in body fluids, such as blood and saliva.

In recent years, the European DNA Profiling Group (EDNAP) has performed some studies on the RNA/DNA co-extraction and proved once again the usefulness of mRNA as a tool for the identification of semen and saliva in forensic cases, compared to the DNA methodology [24].

Another important application of mRNA is in the forensic identification of the skin, using the RT-PCR methods [25].

Since 2009 Hanson et al. [26] have been explored miRNA for their use in forensically body fluids identification. The advantage of these new markers over the mRNAs markers consists in the fact that they have a smaller size containing around 18–22 bp, thus being more stable than mRNA to degradation conditions.

Since last decade new steps have been done in “forensic molecular pathology” and in “post-mortem pharmacogenetics”.

Studies regarding the molecular diagnosis of genetic cardiac arrhythmia or long QT syndrome which leads to sudden death [27] and the post-mortem analysis of gene CYP2D6, which encodes a drug metabolizing enzyme whose variation leads to adverse drug effects and finally to death [28] are new tools which evolve in forensics.

7. Conclusions

Currently, the DNA genotyping of all types of biological traces or micro-traces containing nucleated cells is possible if they are not entirely destroyed, either chemically or bacterial. The DNA analysis represents an important tool in solving caseworks in forensic medicine, such as establishing the custody of a child through paternity or maternity tests, identifying victims from disasters or crimes, exonerating innocent people convicted to prison.

Due to the recent advances in molecular genetics, other biomarkers have been proposed to be used in forensic body fluids identifications, such as messenger RNA (mRNA), microRNA (miRNA), and DNA methylation.

Author details

Raluca Dumache^{1,2*}, Veronica Ciocan^{1,2}, Camelia-Oana Muresan^{1,2} and Alexandra Enache^{1,2}

*Address all correspondence to: raluca.dumache@umft.ro

1 Victor Babes University of Medicine and Pharmacy, Timisoara, Romania

2 Institute of Legal Medicine, Timisoara, Romania

References

- [1] Gill P, Jeffreys AJ, Werrett DJ. Forensic application of DNA 'fingerprints'. *Nature*. 318:577–579 (1985).
- [2] Tamarin HR. *Principles of genetics*. New York: McGraw-Hill (2002).
- [3] Butler JM. *Fundamentals of Forensic DNA Typing*. San Diego, CA: Elsevier Academic Press (2010).
- [4] Kaye DH. *The double helix and the law of evidence*. Cambridge, MA: Harvard University Press (2010).
- [5] Council of Europe. Committee of Ministers'. Recommendation No. R (92) on the use of analysis of deoxyribonucleic acid (DNA) within the framework of the criminal justice system. 470th Meeting of the Minister's Deputies (1992).
- [6] Kuperus WR, et al. Crime scene links through DNA evidence: the practical experience from Saskatchewan casework. *Canadian Society of Forensic Science Journal* 36:19–28 (2003).
- [7] Bond JW, Hammond C. The value of DNA material recovered from crime scenes. *J. Forensic Sci.* 53:797–801 (2008).
- [8] Graham EAM, et al. Room temperature DNA preservation of soft tissue for rapid DNA extraction: an addition to the disaster victim identification investigators toolkit? *Forensic Sci. Int. Genet.* 2:29–34 (2008).
- [9] Reynolds R, Sensabaugh G, Blake E. Analysis of genetic markers in forensic DNA samples using the polymerase chain reaction. *Anal. Chem.* 63:2–15 (1991).

- [10] Goodwin W, Linacre A, Hadi S. An introduction to forensic genetics. Hoboken, NJ: Wiley-Blackwell (2011).
- [11] Jobling MA, Gill P. Encoded evidence: DNA in the forensic analysis. *Nat. Rev. Genet.* 5:739–751 (2004).
- [12] Yoshida K, Yayama K, Hatanaka A, Tamaki K. Efficacy of extended kinship analysis utilizing commercial STR kit in establishing personal identification. *Legal Med.* 13:12–15 (2011).
- [13] Morretti TR, et al. Validation of short tandem repeats (STRs) for forensic usage: performance testing of fluorescent multiplex STR systems and analysis of authentic and simulated forensic samples. *J. Forensic Sci.* 46:647–660 (2001).
- [14] Sullivan KM, Mannucci A, Kimpton CP, Gill P. A rapid and quantitative DNA sex test: fluorescence-based PCR analysis of X–Y homologous gene amelogenin. *Biotechniques.* 15:636–641 (1993).
- [15] Dixon LA, et al. Validation of a 21-locus autosomal SNP multiplex for forensic identification purposes. *Forensic Sci. Int.* 154:62–77 (2005).
- [16] von Heeren F, Thormann W. Capillary electrophoresis in clinical and forensic analysis. *Electrophoresis.* 18:12–13:2415–2426 (1997).
- [17] Shewale JG, Sikka SC, Schneida E, Sinha SK. DNA profiling of azoospermic semen samples from vasectomized males by using Y-PLEX 6 amplification kit. *J. Forensic Sci.* 48:127–129 (2003).
- [18] Ivanov PL, et al. Mitochondrial DNA sequence heteroplasmy in the Grand Duke of Russia Georgij Romanov establishes the authenticity of the remains of Tsar Nicholas II. *Nat. Genet.* 12:417–420 (1996).
- [19] Butler JM, Shen Y, McCord BR. The development of reduced size STR amplicons as tools for analysis of degraded DNA. *J. Forensic Sci.* 48:1054–1064 (2003).
- [20] Brenner CH, Weir BS. Issues and strategies in the identification of World Trade Centre victims. *Theor. Popul. Biol.* 63:173–178 (2003).
- [21] Marchi E. Methods developed to identify victims of the World Trade Centre disaster. *Am. Labor.* 36:30–36 (2004).
- [22] Gill P, Werrett DJ, Budowle B, Guerrieri R. An assessment of whether SNPs will replace STRs in national DNA databases-joint considerations of the DNA working group of the European Network of Forensic Science Institutes (ENFSI) and the Scientific Working Group on DNA analysis methods (SWGDM). *Sci. Justice.* 44:51–53 (2004).
- [23] Zubakov D, Hanekamp E, Kokshoorn M, van Ijcken W, Kayser M. Stable RNA markers for identification of blood and saliva stains revealed from whole genome expression analysis of time-wise degraded samples. *Int. J. Legal Med.* 122:135–142 (2008).

- [24] Haas C, Hanson E, Anjos MJ, et al. RNA/DNA co-analysis from human saliva and semen stains. Results of a third collaborative EDNAP exercise. *Forensic Sci. Int. Genet.* 7:230–239 (2013).
- [25] Visser M, Zubakov D, Ballantyne KN, Kayser M. mRNA-based skin identification for forensic applications. *Int. J. Legal Med.*(2011). DOI: 10.1007/s00414-010-0545-2
- [26] Hanson EK, Lubenove H, Ballantyne I. Identification of forensically relevant body fluids using a panel of differentially expressed microRNAs. *Anal. Biochem.* 387:303–314 (2009).
- [27] Ackerman MJ, Tester DJ, Driscoll DJ. Molecular autopsy of sudden unexplained death in a young. *Am. J. Forensic Med. Pathol.* 22:105–111 (2001).
- [28] Levo A, Koski A, Ojanprena I, Vuori E, Sajantila A. Post-mortem SNP analysis of CYP2D6 gene reveals correlation between genotype and opioid drug(tramadol) metabolite ratio in blood. *Forensic Sci. Int.* 135:9–15 (2003).