We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,300

Open access books available

172,000

International authors and editors

190M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities

WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com

Introductory Chapter: Crosstalk Approach for a Deeper Understanding of the Biological Processes

Mohamed A. El-Esawi

1. Introduction

Molecular signaling has been widely studied in the recent years in order to investigate the different biological processes in living organisms. Information provided by this approach has been utilized to unravel the various functions of different molecules or organs in cells, which in turn facilitate the understanding of the molecular mechanisms underlying the physiological and biochemical processes in these organisms. Therefore, nowadays it is much easier to understand how the biological processes are regulated and controlled inside the organism cells. Understanding the crosstalk and molecular signaling pathways could also help to understand the gene regulatory networking. In plants, studying the signaling processes and crosstalk at the physiological, biochemical, and molecular levels would definitely help to improve the plant growth, development, survival, and productivity as well as to adapt plant crops to the challenging environmental conditions including abiotic and biotic stresses [1–5]. Furthermore, in animals and human, revealing the crosstalk in the biological processes leads to understanding how diseases can be controlled and treated. Therefore, more studies should discuss this matter at the different levels within living organisms. Such kind of information will definitely help to develop different advanced strategies to understand and control the cellular biological processes at different levels.

2. Crosstalk in biological processes

Several earlier studies have reported the crosstalk approach in understanding biological processes in different living organisms. For example, in plants, El-Esawi et al. [1] revealed that Trp triad substitution mutants at W400F and W324F positions can be photoreduced in whole cell extracts, albeit with reduced efficiency. The flavin signaling state (FADH°) has been shown to be stabilized in an in vivo context. These results confirmed that in vivo modulation by metabolites in the cellular environment could have a key role in cryptochrome signaling and is discussed with regard to the possible impacts on the conformation of the C-terminal domain to create the biologically active conformational state. Furthermore, El-Esawi et al. [2] addressed that the blue-light induced biosynthesis of reactive oxygen species may contribute to the signaling mechanism of *Arabidopsis* cryptochrome. El-Esawi et al. [3] also addressed the processes of micropropagation technology

and its applications in crop improvement. Nonzygotic embryogenesis and somatic hybridization processes have been explained and assisted in plant development and crop improvement [4, 5]. Moreover, studying the physiological, biochemical, and molecular processes in plants helped to understand the plant development and to develop improved crop varieties tolerant to different environmental stresses [5–15]. In addition, earlier studies reported the importance of crosstalk in understanding the biological processes in other living organisms such as animals and humans. These approaches and processes could be discussed for further understanding and improvement.

Author details

Mohamed A. El-Esawi

Botany Department, Faculty of Science, Tanta University, Tanta, Egypt

*Address all correspondence to: mohamed.elesawi@science.tanta.edu.eg

IntechOpen

© 2020 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

- [1] El-Esawi M, Glascoe A, Engle D, Ritz T, Link J, Ahmad M. Cellular metabolites modulate *in vivo* signaling of *Arabidopsis* cryptochrome-1. *Plant Signaling & Behavior*. 2015;**10**:e1063758
- [2] El-Esawi M, Arthaut L, Jourdan N, d'Harlingue A, Martino C, Ahmad M. Blue-light induced biosynthesis of ROS contributes to the signaling mechanism of *Arabidopsis* cryptochrome. *Scientific Reports*. 2017;**7**:13875
- [3] El-Esawi MA. Micropropagation technology and its applications for crop improvement. In: Anis M, Ahmad N, editors. *Plant Tissue Culture: Propagation, Conservation and Crop Improvement*. Singapore: Springer; 2016. pp. 523-545
- [4] El-Esawi MA. Nonzygotic embryogenesis for plant development. In: Anis M, Ahmad N, editors. *Plant Tissue Culture: Propagation, Conservation and Crop Improvement*. Singapore: Springer; 2016. pp. 583-598
- [5] El-Esawi MA. Somatic hybridization and microspore culture in *Brassica* improvement. In: Anis M, Ahmad N, editors. *Plant Tissue Culture: Propagation, Conservation and Crop Improvement*. Singapore: Springer; 2016. pp. 599-609
- [6] El-Esawi MA, Alayafi AA. Overexpression of rice *Rab7* gene improves drought and heat tolerance and increases grain yield in rice (*Oryza sativa* L.). *Genes*. 2019;**10**:56
- [7] El-Esawi MA, Al-Ghamdi AA, Ali HM, Alayafi AA, Witczak J, Ahmad M. Analysis of genetic variation and enhancement of salt tolerance in French pea (*Pisum sativum* L.). *International Journal of Molecular Sciences*. 2018;**19**:2433
- [8] El-Esawi MA, Alaraidh IA, Alsahli AA, Ali HM, Alayafi AA, Witczak J, et al. Genetic variation and alleviation of salinity stress in barley (*Hordeum vulgare* L.). *Molecules*. 2018;**23**:2488
- [9] El-Esawi MA, Alaraidh IA, Alsahli AA, Alamri SA, Ali HM, Alayafi AA. *Bacillus firmus* (SW5) augments salt tolerance in soybean (*Glycine max* L.) by modulating root system architecture, antioxidant defense systems and stress-responsive genes expression. *Plant Physiology and Biochemistry*. 2018;**132**:375-384
- [10] El-Esawi MA, Alaraidh IA, Alsahli AA, Alzahrani SM, Ali HM, Alayafi AA, et al. *Serratia liquefaciens* KM4 improves salt stress tolerance in maize by regulating redox potential, ion homeostasis, leaf gas exchange and stress-related gene expression. *International Journal of Molecular Sciences*. 2018;**19**:3310
- [11] El-Esawi MA, Al-Ghamdi AA, Ali HM, Alayafi AA. *Azospirillum lipoferum* FK1 confers improved salt tolerance in chickpea (*Cicer arietinum* L.) by modulating osmolytes, antioxidant machinery and stress-related genes expression. *Environmental and Experimental Botany*. 2019;**159**:55-65
- [12] Jourdan N, Martino C, El-Esawi M, Witczak J, Bouchet P-E, d'Harlingue A, et al. Blue light dependent ROS formation by *Arabidopsis* Cryptochrome-2 may contribute towards its signaling role. *Plant Signaling & Behavior*. 2015;**10**:e1042647
- [13] Vwioko E, Adinkwu O, El-Esawi MA. Comparative physiological, biochemical and genetic responses to prolonged waterlogging stress in okra and maize given exogenous ethylene priming. *Frontiers in Physiology*. 2017;**8**:632

[14] El-Esawi MA, Al-Ghamdi AA, Ali HM, Ahmad M. Overexpression of *AtWRKY30* transcription factor enhances heat and drought stress tolerance in wheat (*Triticum aestivum* L.). *Genes*. 2019;**10**(2):163

[15] El-Esawi MA, Alayafi AA. Overexpression of *StDREB2* transcription factor enhances drought stress tolerance in cotton (*Gossypium barbadense* L.). *Genes*. 2019;**10**:142