

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,800

Open access books available

142,000

International authors and editors

180M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Changing Climate and Advances on Weeds Utilization as Forage: Provisions, Nutritional Quality and Implications

Muhammad Aamir Iqbal, Sajid Ali, Ayman El Sabagh, Zahoor Ahmad and Muzammil H. Siddiqui

Abstract

Under changing climate, growth and prevalence of many invasive and indigenous weeds are expected to boost up owing to their greater genetic diversity, competitive superiority and better plant architecture. Atmospheric CO₂ enrichment and elevating global temperature are causing weeds invasion to new localities making prevalent weed management strategies ineffective. Weed utilization as forage for ruminants provided that their nutritional profile is available and can be a biologically feasible and economically viable approach compared to existing management system of eliminating them from agro-ecological systems. Different weeds like Bermuda grass (*Cynodon dactylon*), Johnson grass (*Sorghum halepense*), canary grass (*Phalaris minor*), nut sedge (*Cyperus rotundus*), yellow duck (*Rumex crispus*), drooping brome (*Bromus tectorum*), burr clover (*Medicago polymorpha*), button weed (*Diodia scandens*), and purslane (*Portulaca oleracea*) had acceptable nutritional profile with organic matter (89.0–91.3%), protein (7.1–19.5%) and fats (2.1–3.7%). Those were also rich in micro-nutrients (calcium, magnesium and zinc), while anti-nutritional factors (saponins, tannins, phytates and oxalates) were in safer limits for dairy animals. Lack of nutritional profiling and presence of anti-nutritional factors decreased feed intake and led to malnutrition, while higher concentration of tannins caused digestibility depression in small ruminants. There is need to conduct further studies for nutritional profiling of local weed species and development of techniques for reducing their anti-nutritional factors.

Keywords: anti-nutritional factors, global warming, protein content, saponins, tannins

1. Introduction

Climate change has been feared to incur frequent drought spells and floods, while temperature fluctuations and shifting of rainfall patterns are projected to alter growth habits of weeds flora globally. There is an emerging rhetoric that most of invasive and indigenous weeds have the potential and botanical superiority to adjust and acclimatize to atmospheric CO₂ enrichment through the optimization of photosynthesis process leading to significant boost in their biomass production.

In this way, some of the weed species can increase their establishment and dominance in indigenous agro-ecosystems along with invading adjacent as well as far flung terrestrial ecosystems. Thus contrary to notion that weeds are menace and agro-ecological systems must be kept free of indigenous and invasive weeds, their utilization as forage for dairy animals has the potential to become the most feasible and pro-environment strategy [1–3].

Globally, large ruminant's performance is directly influenced by the nutritional value of feed which accounts for over 50% of total expenditures. It deserves mentioning that dairy animals confront forage shortage owing to temperature extremes leading to drastic fall in milk production especially in developing countries. The shortage of forage and rising population of dairy animals has necessitated identifying and evaluating alternate feed resources which are cheap and can also fulfill animal's dietary needs. It has been established that weeds can inflict drastic influence on crops productivity and use of chemical herbicides for keeping them below threshold level, has led to serious concerns pertaining to their residual persistence in crops, soil and environment. Weeds utilization as animal feed holds potential because these are cheap owing to their abundance on field paths and water channels. Weeds harvested from cropped and non-cropped area may constitute an effective and biologically viable approach to keep weeds below the threshold level. Weeds utilization for feeding animals can also reduce herbicides use in agricultural lands which has the potential to curb environmental pollution. Many weeds have been reported to be resistant and better adapt to dynamic environmental conditions, and thus making them less prone to drastic impacts of climate change. In addition, it was reported that animals preferred naturally grown mixtures of weeds over crop residues and roughage during dry season. Furthermore, rapid regeneration favors many weed species for their inclusion as a source of vegetable protein in animal's diet [4–10].

Weeds such as Bermuda grass (*Cynodon dactylon*), Johnson grass (*Sorghum halepense*), canary grass (*Phalaris minor*), nut sedge (*Cyperus rotundus*), cheat-grass or drooping brome (*Bromus tectorum*), burr clover (*Medicago polymorpha*) and pigweed (*Amaranthus viridis*) contained organic matter over 90% indicating that these weeds can fulfill the dry matter requirement of animals. In addition, spotted knapweed (*Centaurea stoebe ssp. micranthos*), a weed of rangelands in Northern America was reported to displace local plant species, degraded wildlife habitats, altered biogeochemistry of soil and triggered soil erosion and thus its control through grazing was found to be biologically and economically viable. Similarly, broom snakeweed and medusa-head (*Taeniatherum caput-medusae*) were effectively controlled through controlled grazing with reasonably good palatability [11–17].

Along with substantial quantity, nutritional quality of weeds is of the utmost importance for dairy animals in order to produce milk on sustainable basis. Field bindweed (*Convolvulus arvensis*) and yellow duck (*Rumex crispus*) were reported to have significantly higher protein content (27 and 22% respectively) [11] which were greater than all cereal forages and most of the legumes, while button weed (*Diodia scandens*) contained 7.7% protein [18]. Although, a number of species belonging to *Commelinaceae* family such as climbing dayflower (*Commelina diffusa* L.), tropical spiderwort/wandering jew (*Commelina benghalensis* L.), Asiatic dayflower (*Commelina communis* L.), African dayflower (*Commelina africana* L.), white mouth dayflower/slender dayflower (*Commelina erecta*) are considered weed but constituted a major chunk of animal feed in Tanzania [19], rural regions of Mauritius [20], USA [21] and in Kenya owing to reasonably good palatability [14, 18, 22, 23].

It is pertinent to mention that anti-nutritional factors (saponins, tannins, oxalates, etc.) of weeds constitute as the most crucial concern as far as animal nutrition is concerned. Tick weed (*Cleomea viscosa*) recorded safer limits of anti-nutritional factors such as condensed tannins (0.0491%), saponins (0.23%), phytates (1.2%)

and oxalates (3.3%), but unfortunately in-depth studies are lacking in this context. Furthermore, narrow leaf weeds (nut sedge, wild oat, etc.) were recommended to be a good source of fiber, while broad leaf weeds (pigweed, field bindweed, etc.) provided cheap vegetable protein to dairy animals [1, 11, 13, 20].

To date, very few studies have been done to assess the nutritional status, digestibility and intake of indigenous and exotic (invasive) weeds and their utilization in sustainable ruminant's production systems, but not a single study has so far synthesized and evaluated the literature on weeds utilization as forage. This chapter attempts to synthesize as well as assess the potential of weeds for supplementing traditional feedstuffs (forages, crop residues and concentrates) partially without compromising the productivity of large ruminants in terms of milk production. Weeds mineral constituents and anti-nutritional factors and various implications in weeds utilization as animal feed have also been evaluated.

2. Materials and methods

In order to synthesize published findings pertaining to nutritional quality of weeds, search was performed on Google Scholar (<http://scholar.google.com>) and PubMed (<http://www.PubMed.gov>) using the below mentioned search strings:

1. Weeds and forages.
2. Weeds nutritional value.
3. Weeds, animal feed.
4. Anti-nutritional factors in weeds.

The search was time-restricted to 2000–2019, however owing to limitation of published findings; it was later on relaxed to 1990–2019.

The research studies were screened based on following criteria;

1. Reporting at least one weed's biomass production under changing climate.
2. Describing one or more nutritional quality parameters such as protein content of indigenous or invasive weeds.
3. Stating anti-nutritional factors of weeds species.
4. Reporting mineral constituents of at least one or more weeds.

The screening process resulted in 55 studies which fully fit in the objectives and selection criteria.

3. Weeds under changing climate

The rising temperature and carbon dioxide level along with the rapidly altering dynamics of rainfall and evaporation are the most important factors for determining management and utilization of weeds under changing climate. Weeds have been reported to have a greater genetic diversity compared to crops and thus can respond positively to agro-environmental changes. Owing to CO₂ enrichment of atmosphere and rising temperature, some of the weed species can invade new

geographical localities while making the existing weed management strategies ineffective. In addition, weeds can have superiority over crop plants by virtue of better plant architecture and incorporating nitrogen and carbon in seeds. Rag weed (*Ambrosia artemisiifolia*) developed more number of branches and leaf area along with producing greater number of pollens under increased temperature. Similarly, comparatively higher production of spines by Canada thistle (*Cirsium arvense*) in response to elevated CO₂ level was reported [10, 24–27].

In addition, biomass production of bitter vine/American rope (*Mikania mikrantha*), creeping oxeye (*Wedelia trilobata*) and Cairo morning glory (*Ipomea cairica*) was enhanced with increasing CO₂ level [28]. Spurred anoda (*Anoda cristata*) gave the highest green biomass at CO₂ fertilization up to 700 ppm and 32°C temperature, while barnyard grass (*Echinochloa crusgalli*) and Indian goose-grass/wire grass/crowfoot grass (*Eleusine indica*) remained non-responsive to elevated temperature and CO₂ concentration. It was concluded that elevated CO₂ effectively enhanced the photosynthetic process even under water limited conditions indicating higher water use efficiency of weeds under drought stress which led to higher biomass production. However, weeds response to elevated CO₂ and temperature under well watered conditions continues to remain an unexplored aspect which demands further research to determine the physiological plasticity of different weed species [25, 26].

The temperature elevation as a result of global warming is feared to trigger weeds migration. Cogon grass (*Imperata cylindrica*) prickly acacia (*Acacia nilotica*) and witch weed (*Striga asiatica*) were reported to invade cooler areas of Europe owing to global warming, while some of the invasive weed species such as mesquite (*Prosopis juliflora*) can become more hardy and difficult to control owing to greater partitioning of assimilates to roots under elevated temperature particularly under agro-ecological conditions of Indo-Pak subcontinent [27, 29–32].

4. Nutritional quality of weeds

Although, weeds presence in and around the cultivated fields has never been deemed desirable, but these can contribute significantly to the production of quality organic feed for dairy animals. The nutritional profile of weeds determines feasibility and scope for their inclusion in ruminant's feed. The nutritional quality of weeds encompasses digestibility, chemical composition, energy and extent of presence of anti-nutritional factors and such information can assist to determine the allowable proportion of weeds in ruminant's feed [10, 33–35].

Dry matter digestibility has direct relationship with the quality of forage. Different weeds such as Barnyard grass (*Echinochloa crusgalli*) and Jerusalem artichoke (*Helianthus tuberosus*) had significantly higher digestibility compared to many cereal forages. The comparative dry matter digestibility of many weeds and forages crops is presented in **Table 1**. Digestibility was reported to be an important indicator of any forage's quality, while Lamb-squarters (*Chenopodium album*), barn-yard-grass (*Echinochloa crus-galli*), dandelion (*Taraxacum officinale*), Jerusalem artichoke (*Helianthus tuberosus*), yellow foxtail (*Setaria glauca*), perennial sowthistle (*Sonchus arvensis*) and Canada thistle (*C. arvense*) had in-vitro dry matter digestibility equal to alfalfa (*Medicago sativa*). In addition, common ragweed (*Ambrosia artemisiifolia*) and redroot pigweed (*Amaranthus retroflexus*) had even greater in-vitro dry matter digestibility compared to alfalfa [11, 39–41]. There is dire need to determine the digestibility of local weed flora in order to find out their suitability as an alternate animal feed.

Protein (CP) is the most important nutritional quality attribute having direct impact on milk production. Testing of 102 weed species belonging to *Poacea*,

Weeds	DMD (%)	Forage crops	DMD (%)
Barnyard grass (<i>Echinochloa crusgalli</i>)		Alfalfa (<i>Medicago sativa</i>)	64–75
Canada thistle (<i>Cirsium arvense</i>)	68–74	Sorghum (<i>Sorghum bicolor</i>)	59–61
Dandelion (<i>Taraxacum officinale</i>)	78–84	Maize (<i>Zea mays</i>)	63–68
Sowthistle (<i>Sonchus arvensis</i>)	76–82	Oat (<i>Avena sativa</i>)	60–63
Swamp smartweed/knotweed/tanwed (<i>Polygonum amphibium</i>)	54–62	Barley (<i>Hordeum vulgare</i>)	59–64
Quackgrass (<i>Elymus repens</i>)	58–68	Pearl millet (<i>Cenchrus americanus</i>)	58–60
Brome grass (<i>Bromus tectorum</i>)	66–76	Cowpea (<i>Vigna unguiculata</i>)	68–76
Curly dock (<i>Rumex crispus</i>)	50–58	Soybean (<i>Glycine max</i>)	70–76
Jerusalem artichoke (<i>Helianthus tuberosus</i>)	81–86	Cluster bean (<i>Cyamopsis tetragonoloba</i>)	70–79

Table 1.
 Dry matter digestibility of some weeds and common forage crops grown under varied agro-climatic conditions [11, 28, 33, 36–38].

Asteraceae, *Fabaceae* and *Euphorbiaceae* families commonly found in central Mexico revealed that only 25 had balanced nutritional profile. Weeds CP content depend on growth stage as matured weeds recorded lesser protein compared to harvestings done at pre-bloom stage. Asthma plant (*Eurphobia hirta*) recorded 16.7% protein content while tick weed or Asian spider-flower (*Cleomea viscosa*) with 14.7% followed it, while yellow nutsedge or nut grass (*Cyperus esculentus*) and button weed (*Diodia scandens*) contained 9.8 and 7.7% CP respectively. Bluegrass (*Poa annua*) was found to have over 14% which is higher than maize, sorghum and oat, while common purslane (*Portulaca oleracea*) (8%) was also suggested to be an equally good forage weed as far as CP content is concerned. Another study suggested that weeds including bush sunflower (*Simsia amplexicaulis*), creeping false holly (*Jaltomata procumbens*) and mosquito flower weed (*Lopezia racemosa*) contained CP in the range of 6.5–16.9% and could be used solely or as supplementary feed mixed with maize straw to feed dairy cattle. Mixtures of weeds (*Commelinaceae* + *Amaranthaceae*) recorded crude protein twice than most of the roughages. Another study reported that bush sunflower (*Simsia amplexicaulis*) weed supplemented with maize straw based animal diets resulted in higher protein content successfully met dairy animals dietary needs. Similarly, climbing dayflower (*Commelina diffusa* L.) recorded appreciably higher content of protein (17.7%) which is comparable to commonly used forage crops. In addition, its rumen degradability of protein was recorded over 72% making it forage with balanced nutrition [11, 36, 42–44].

Higher content of fiber increases the bulkiness of feed which results in reduced intake. The lowest crude fiber content was recorded by button weed (18.7%) and nut grass yielded the highest fiber (27%). The minimum lignin content (9.6%) of asthma plant favored its inclusion in animal feed [18]. Similarly, common dandelion (*Taraxacum officinale*) recorded significantly lower crude fiber content (15%), while Bermuda grass (*Cynodon dactylon*) gave the lowest fiber content of just over 6% [11]. In contrast, barnyard grass (*Echinochloa crusgalli*) recorded the highest neutral detergent fiber (NDF) compared to trans-pecos drymary (*Drymaria laxiflora*) [45]. It was reported that climbing dayflower (*Commelina diffusa* L.) recorded 36% and 22% NDF and acid detergent fiber (ADF) respectively and thus compares well to commonly used grasses such as sorghum-Sudan grass and napier

grass (*Pennisetum purpureum*) [9]. Similarly, spiderwort (*Tripogandra purpuracens*), a weed of South America recorded reasonably good concentration of carbohydrates which was higher compared to *Tridax coronopifolia* and was recommended to be fed to dairy animals [10, 11, 13, 19, 37, 46].

Digestibility is an important indicator of any forage's quality. Lamb-squarters (*Chenopodium album*), barn-yard-grass (*Echinochloa crus-galli*), dandelion (*Taraxacum officinale*), Jerusalem artichoke (*Helianthus tuberosus*), yellow foxtail (*Setaria glauca*), perennial sowthistle (*Sonchus arvensis*) and Canada thistle (*C. arvense*) had in-vitro dry matter digestibility equal to alfalfa (*Medicago sativa*). In addition, common ragweed (*Ambrosia artemisiifolia*) and redroot pigweed (*Amaranthus retroflexus*) had even greater in-vitro dry matter digestibility compared to alfalfa. Similarly, it was reported that high protein and low fiber contents are indicative of high energy and high productive value feeds. Field bindweed (*Convolvulus arvensis* L.) and common amaranth (*Amaranthus retrofl exus* L.) recorded the highest protein (18.8 and 13.0% respectively) and the lowest fiber (14.7 and 17.6% respectively) which was comparable to alfalfa (*Medicago sativa*) hay having 16.9% protein and 27% fiber. In addition, especial emphasis was paid to palatability of weeds as high nutritional value becomes irrelevant if animals have little likelihood for the weeds species. The hay of different weeds was given to sheep to determine their palatability by using cafeteria of manger technique and biomass consumed in 15 minutes was recorded. Alfalfa had the highest palatability followed by field bindweed (*Convolvulus arvensis* L.)

Weeds	OM (%)	CP (%)	ADF (%)	NDF (%)	F (%)	A (%)
Knapweed (<i>Centaurea stoebe</i>) [43]	—	19.5	—	29.5	—	—
Nut sedge (<i>Cyperus rotundus</i>) [11]	91.03	16.3	57.8	64.5	—	12.8
Red dead-nettle (<i>Lamium purpureum</i>) [43]	—	9.7	25.8	—	2.1	9.8
Field bindweed (<i>Convolvulus arvensis</i>) [11]	90.30	27.0	41.0	35.5	—	10.4
Pigweed (<i>Amaranthus viridis</i>) [11]	91.00	26.2	57.7	31.0	—	13.2
Johnson grass (<i>Sorghum halepense</i>) [46]	—	5.3	30.2	—	1.5	5.5
Field mustard (<i>Brassica rapa</i>) [46]	—	9.8	49.5	63.7	—	—
Chicory (<i>Cichorium intybus</i>) [46]	—	7.1	35.2	—	3.34	7.5
Bermuda grass (<i>Cynodon dactylon</i>) [11]	90.90	13.5	47.0	76.5	—	13.3
Mexican aster (<i>Cosmos bipinnatus</i>) [40]	—	10.5	43.5	41.7	—	—
Tick clover (<i>Desmodium molliculum</i>) [40]	—	16.2	42.9	41.6	—	—
Yellow foxtail (<i>Setaria glauca</i>) [10]	—	20	30.0	—	—	—

Table 2. Nutritional quality (organic matter, crude protein CP, acid detergent fiber ADF, neutral detergent fiber NDF, fats F, total ash A) of weeds [11, 17, 21, 31, 35, 39, 41, 48, 49].

and common amaranth (*Amaranthus retrofl exus* L.) owing to higher protein and lesser fiber contents. Thus, it was inferred that protein and fiber content of feeds can be used as predictors of palatability and it was also concluded that weeds leaves had 2–3 times higher protein than stems and thus leafy weeds such as field bindweed (*Convolvulus arvensis* L.) recorded higher palatability [11, 45, 47]. The nutritional quality of some weeds has been presented **Table 2**.

5. Mineral constituents of weeds

Minerals in appropriate quantity are essential for dairy animals to be utilized in various metabolic processes, for boosting immunity level against diseases and reproductive health. Asthma plant (*Eurphobia hirta*) was recommended to be included in animal feed for having reasonably higher concentrations of major minerals including calcium (Ca) (13.6%), magnesium (Mg) (3.0%) and potassium (K) (2.5%), along with many trace elements such as iron (Fe) (0.7%), copper (Cu) (0.1%) and manganese (Mn) (0.1%). Common chicory (*Cichorium intybus*) was also suggested as forage weed for having a comparable mineral composition including Ca (6%), Mg (2%), Fe (0.5%) and Cu (0.06%). In addition, pink sorrel (*Oxalis debilis*) was found to be poor on animal nutrition scale for being deficient in Ca (4%), Mg (2.3%), Fe (2.4) and Zinc (Zn) (0.15) compared to other forage weeds [10, 21, 48]. Very limited information has been reported so far regarding mineral constituents of weeds which limit their utilization as a feed source for ruminants. **Table 3** contains mineral constitution of some weeds.

Weeds	Ca	Mg	Zn
Wild oat (<i>Avena fatua</i>) [11]	1.8	1.10	0.06
Burr clover (<i>Medicago polymorpha</i>) [11]	10.2	2.42	0.14
Morning glory (<i>Ipomoea purpurea</i>) [40]	9.0	0.63	2.99
Yellow duck (<i>Rumex crispus</i>) [11]	4.7	2.70	0.20
Cheese weed (<i>Malva parviflora</i>) [40]	19.3	1.22	4.59
Wood sorrel (<i>Oxalis decaphyllai</i>) [40]	5.1	1.43	2.76

Table 3. Mineral constituents (calcium Ca, magnesium Mg and zinc Zn) of different weeds grown under varied agro-climatic conditions [11, 18, 22, 32, 40].

6. Anti-nutritional contents of weeds

Condensed tannins, saponins, phytate and oxalate are some of the anti-nutritional factors which reduce the nutritional quality and even impart toxicity to animal feeds. It was reported that button weed (0.029–0.052%) had the lowest tannin content, while nutsedge recorded the maximum tannin content. It was suggested that weeds having tannins 2–4% of dry matter did not pose a life threatening situation rather were found to be effective in improving protein flow towards duodenum which led to higher weight gain and reduced the parasitic infections. It was suggested that effective drying of weeds has the potential to significantly reduce the condensed tannins of weeds. Similarly, saponins which are generally produced by defense systems of weeds in response to pathogenic attacks impart a bitter taste and reduce nitrogen

digestibility leading to lower palatability of weeds. Tick weed and nutsedge recorded similar saponins (0.22%), while button weed contained higher saponins (0.35%).

Phytates produce phytic acid which acts as a chelator of various macro-minerals (calcium and magnesium) and trace mineral (iron and zinc) leading to a severe deficiency of these minerals. Button weed with 1.18% phytates remained superior to nut-sedge and asthma plants. Oxalate is another important anti-nutritional factor which binds with calcium to form calcium oxalate leading to calcium unavailability. Asthma plant recorded the lowest oxalate concentration (2.36%) while button weed had the maximum oxalate concentration (2.92).

It was suggested that weeds such as Arizona sunflower (*Tithonia tubiformis*), wood sorrel (*Oxalis divergens*) and bush sunflower (*Simsia amplexicaulis*) commonly found in America, Mexico, Argentina and Chile contained tannins in safer limits and might be utilized to feed dairy animals. In contrast, a fatty acid called malvalic acid was isolated from cheese weed (*Malva parviflora* L.) which caused deaths of dairy animals. Similarly, different phenolic compounds were reported to be the major reason behind low palatability of many weed species. In addition, the presence of phytochemicals and free oxygen metabolites in weeds contributed to mastitis and ultimately led to udder edema along with deteriorating the reproductive performance of cattle. Similarly, spotted knapweed (*Centaurea stoebe* ssp. *micranthos*) in North American rangelands contained an allelo-chemical named cnicin (a sesquiterpene lactone compound) which reduced its palatability by imparting bitter taste and deterred grazing. In contrast, knapweed was readily consumed by small ruminants' preferably at rosette and bolting stages compared to flowering and seed-set phonological stages. This preference was associated with higher protein and lower fiber content at rosette stage in comparison to flowering or seed-set stages without any link between cnicin content and knapweed palatability [28, 49–51].

7. Nutritional comparison of weeds and forage crops

To the best of our knowledge, no comprehensive studies have been reported pertaining to qualitative analyses of different native and exotic weeds with forage crops. Some of the weeds such as Canada thistle, spotted knapweed, white-top, Russian knapweed and pigweeds contain protein in the range of 15–22% while typical grasses has only 2–11% protein, thus have the potential to become cost-free source of plant protein. In addition, higher leaf-stem ratio in weeds impart them superiority over grasses in terms of higher digestibility. Moreover, weeds provide nutrients rumen microbes which enable dairy animals to digest lower quality forages and thereby reducing overall feed cost. Weeds are always available even during periods of drought in arid areas and their utilization can help to obtain sustainable supplies of milk throughout the years with minimum cost. Dairy animals being fed on protein rich weeds tend to gain weight more rapidly and that too with no additional cost.

8. Limitations and implications

The studies regarding nutritional quality, presence of anti-nutritional or toxic substances and palatability of most of the native weed species is lacking. However, one of the most important limitations in utilizing tropical weeds for dairy animals is the presence of anti-nutritional factors such as tannins which are harmful and toxic to ruminants [13, 38, 51]. Animals being fed on weeds having high concentration of tannins witnessed digestibility depression. Weeds having exceptionally higher lignin content caused a sharp decline in feed intake leading to serious malnutrition in dairy

animals. Similarly, significantly less palatability of weeds reduce their intake, which led to animal's weight loss along with sharp decline in milk production [52–54].

Moreover, some of the weeds species have thorns and spines due to which animal gets its mouths injured along with irritating of eyes which leads to pinkeye. In addition, there could be some weeds which can impart unpleasant odor and taste to milk and meat. Lastly, although weeds offer cost-free source of animal feed but centuries-long war against weeds has made it difficult to change the mind of ranchers and dairy farmers to utilize this precious source of plant protein which needs to be changed.

9. Conclusions

The exceptional resistance to drought, higher biomass production under unfavorable pedo-climatic conditions, rapid regeneration capacity, and acceptable nutritional quality at all phenological stages suggests that there are opportunities to utilize weeds as forage for all types of ruminants. Weeds availability throughout the year warrants their potential to fulfill essential dietary needs of animals and favors their inclusion as supplementary forage especially during extreme weather conditions. However, controlled field investigations for determining the appropriate growth timings, nutritional quality, anti-nutritional factors and biomass production potential of different native weeds must be done in rangelands while maintaining a balanced and healthy rangeland ecosystem.

Author details

Muhammad Aamir Iqbal^{1*}, Sajid Ali², Ayman El Sabagh^{3,4}, Zahoor Ahmad⁵
and Muzammil H. Siddiqui^{1*}

1 Department of Agronomy, Faculty of Agriculture, University of Poonch Rawalakot (AJK), Pakistan

2 Institute of Agricultural Sciences, University of the Punjab, Lahore, Pakistan

3 Department of Field Crops, Faculty of Agriculture, Siirt University, Turkey

4 Department of Agronomy, Faculty of Agriculture, Kafrelsheikh University, Egypt

5 Department of Field Crops, Faculty of Agriculture, Cukurova University, Turkey

*Address all correspondence to: muhammadaamir@upr.edu.pk and muzammilagro@upr.edu.pk

IntechOpen

© 2020 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Ziska LH, Faulkner SS, Lydon J. Changes in biomass and root:shoot ratio of field grown Canada thistle (*Cirsium arvense*), a noxious, invasive weed, with elevated CO₂. *Weed Science*. 2004;**47**:608-615
- [2] Kalita P, Mukhopadhyay PK, Mukherjee AK. Evaluation of the nutritional quality of four unexplored aquatic weeds from Northeast India for the formulation of cost-effective fish feeds. *Food Chemistry*. 2007;**103**:204-209
- [3] Iqbal MA, Muzammil HS, Sher A, Zahoor A, Qaiser M, Rana DK. Forage productivity of cowpea [*Vigna unguiculata* (L.) Walp] cultivars improves by optimization of spatial arrangements. *Revista Mexicana De Ciencias Pecuarias*. 2018;**9**:203-219
- [4] Iqbal MA, Asif I, Muzammil HS, Zahid M. Bio-agronomic evaluation of forage sorghum-legumes binary crops on Haplic Yermosol soil of Pakistan. *Pakistan Journal of Botany*. 2018;**50**:1991-1997
- [5] Iqbal MA, Asif I, Zahid M, Zahoor A, Essa A, Muzammil HS, et al. Revamping soil quality and correlation studies for yield and yield attributes in sorghum-legumes intercropping systems. *Bioscience Journal*. 2018;**34**:1165-1176
- [6] Iqbal MA, Asif I, Rana NA. Spatio-temporal reconciliation to lessen losses in yield and quality of forage soybean (*Glycine max* L.) in soybean-sorghum intercropping systems. *Bragantia*. 2018;**77**:283-291
- [7] Iqbal A, Muhammad AI, Imtiaz H, Muzammil HS, Muhammad N, Jamshed A, et al. Seed blending of oat (*Avena sativa* L.) and canola (*Brassica napus* L.) under variable seed proportions enhanced forage productivity and nutritional quality. *Pakistan Journal of Botany*. 2018;**50**:1985-1990
- [8] Iqbal MA, Abdul H, Imtiaz H, Muzammil HS, Tanveer A, Abdul K, et al. Competitive indices in cereal and legume mixtures in a South Asian environment. *Agronomy Journal*. 2019;**111**:242-249. DOI: 10.2134/agronj2017.11.0658
- [9] Lanyasunya TP, Wang HR, Abdulrazak SA, Mukisira EA, Zhang J. The potential of the weed, *Commelina diffusa* L., as a fodder crop for ruminants. *South African Journal of Animal Science*. 2006;**36**:27-31
- [10] Ziska LH. Evaluation of yield loss in field sorghum from a C3 and C4 weed with increasing CO₂. *Weed Science*. 2003;**51**:914-918
- [11] Khan R, Khan MA, Sultan S, Marwat KB, Khan I, Hassan G, et al. Nutritional quality of sixteen terrestrial weeds for the formulation of cost-effective animal feed. *Journal of Animal and Plant Sciences*. 2013;**23**:75-79
- [12] Thrift BD, Mosley JC, Brewer TK, Roeder BL, Olson BE, Kott RW. Prescribed sheep grazing to suppress spotted knapweed on foothill rangeland. *Rangeland Ecology and Management*. 2008;**61**:18-25
- [13] LeJeune KD, Seastedt TR. *Centaurea* species: The forb that won the west. *Conservation Biology*. 2001;**15**:1568-1574
- [14] Olson BE, Lacey JR. Sheep: A method for controlling rangeland weeds. *Journal of Sheep and Goat Research*. 1994;**10**:105-112
- [15] Wright AL, Kelsey RG. Effects of spotted knapweed on cervid winter-spring range in Idaho. *Journal of Range Management*. 1997;**50**:487-496

- [16] Olson BE, Kelsey RG. Effect of *Centaurea maculosa* on sheep rumen microbial activity and mass in vitro. *Journal of Chemistry and Ecology*. 1997;**23**:1131-1144
- [17] Ralphs MH, Banks JE. Cattle grazing as a biological control for broom snakeweed: Vegetation response. *Rangeland Ecology and Management*. 2009;**62**:38-43
- [18] Akinfemi A, Mako AA. Assessment of nutritive value of four dominant weed species in north central Nigeria. *Livestock Research for Rural Development*. 2012;**24**. Available from: <http://www.lrrd.org/lrrd24/11/akin24208.htm>
- [19] Ingratubun GF, Owen E, Massawe NF, Mtenga LA, Mtengeti EG. Effect of upgrading small East African goats on feed resource utilisation in the Uluguru mountains in Tanzania; a farmers' perspective. *Livestock Research for Rural Development*. 2000;**12**:13-19
- [20] Boodoo AA, Ramjee R, Hulman B, Dolberg F, Rowe JB. Evaluation of the basal forage diet of village cows. *Livestock Research for Rural Development*. 1990;**1990**:2. Available from: <http://www.lrrd.org/lrrd24/11/akin24208.htm>
- [21] Mullahey JJ. Tropical soda apple, wetland nightshade, and Turkey berry. In: Van Driesche R et al., editors. *Biological Control of Invasive Plants in the Eastern United States*. Florida, USA: USDA Forest Service Publication; 2002. pp. 413-423
- [22] Walker JW, Kronberg SL, Al-Rowaily SL, West NE. Comparison of sheep and goat preference for leafy spurge. *Journal of Range Management*. 1994;**47**:429-434
- [23] Aggarwal GC, Pimentel D, Mario G. Weed benefits and costs in rice and wheat production in India. *Agricultura, Ecosystems & Environment*. 1992;**39**:235-244
- [24] Hyvonen T, Glemnitz M, Radics L, Hoffmann J. Impact of climate and land use type on the distribution of finish casual arable weeds in Europe. *Weed Research*. 2010;**51**:201-208
- [25] Kriticos DJ, Sutherst RW, Brown JR, Adkins SW, Maywald GF. Climate change and the potential distribution of an invasive alien plant: *Acacia nilotica* spp. *indica* in Australia. *Journal of Applied Ecology*. 2003;**40**:111-124
- [26] Patterson DT. Weeds in a changing climate. *Weed Science*. 1995;**43**:685-701
- [27] Malarkodi N, Manikandan N, Ramaraj AP. Impact of climate change on weeds and weed management—A review. *Journal of Innovative Agriculture*. 2017;**4**:1-6
- [28] Song L, Wu J, Furong LC, Peng S, Chen B. Different responses of invasive and native species to elevated CO₂ concentration. *Acta Ecologica*. 2009;**35**:128-135
- [29] Chaudhury UN, Kirkham MB, Kanemasu ET. Carbon dioxide and water level effects on yield and water use of winter wheat. *Agronomy Journal*. 1990;**82**:637-641
- [30] Ziska LH. Influence of rising atmospheric CO₂ since 1900 on early growth and photosynthetic response of a noxious invasive weed, Canada thistle (*Cirsium arvense*). *Functional Plant Biology*. 2002;**29**:1387-1392
- [31] Ziska LH, George K. Rising carbon dioxide and invasive, noxious plants: Potential threats and consequences. *World Resource Review*. 2004;**16**:427-447
- [32] Tang J, Xu L, Chen X, Hu S. Interaction between C4 barnyard grass

and C3 upland rice under elevated CO₂: Impact of mycorrhizae. *Acta Ecologica*. 2009;**35**:227-235

[33] Castelan O, Estrada J, Carretero L, Vieyra A, Martínez N, Cárdenas S, et al. Degradation characteristics of maize weeds used as forage in smallholder maize-livestock production systems of central México, in different growing periods. *Tropical Subtropical Agroecosystems*. 2003;**3**:115-119

[34] De-Bruijn SL, Bork EW. Biological control of Canada thistle in temperate pastures using high density rotational cattle grazing. *Biological Control*. 2006;**36**:305-315

[35] Di-Tomaso JM, Kyser GB, Goerge MR, Doran MP, Laca EA. Control of Medusahead (*Taeniatherum caput-medusae*) using timely sheep grazing. *Invasive Plant Science Management*. 2008;**1**:241-247

[36] Holst PJ, Allan CJ, Campbell MH, Gilmour AR. Grazing of pasture weeds by goats and sheep, pt. 1: Nodding thistle (*Carduus nutans*). *Australian Journal of Experimental Agriculture*. 2004;**44**:547-551

[37] Lamers J, Burkert A, Makkar HPS, Von Oppen M, Becker K. Biomass production, feed and economic value of fodder weeds as by-products of millet cropping in a Sahelian farming system. *Experimental Agriculture*. 1996;**32**(3):317-326

[38] Ziska LH, Caulfield FA. Rising CO₂ and pollen production of common ragweed (*Ambrosia artemisiifolia*), a known allergy inducing species: Implications for public health. *Australian Journal of Plant Physiology*. 2000;**27**:893-898

[39] Di-Tomaso JM. Invasive weeds in rangelands: Species, impacts, and management. *Weed Science*. 2000;**48**:255-265

[40] Dora G, Mendoza S, Serrano V, Bah M, Pelz R, Balderas R, et al. Proximate composition, mineral content, and antioxidant properties of 14 Mexican weeds used as fodder. *Weed Biology and Management*. 2008;**8**:291-296

[41] Frost RA, Launchbaugh KL. Prescription grazing for rangeland weed management. *Rangelands*. 2003;**25**:43-47

[42] Frost RA, Wilson LM, Launchbaugh KL, Hovde EM. Seasonal change in forage value of rangeland weeds in Northern Idaho. *Invasive Plant Science and Management*. 2008;**1**:343-351

[43] Ganguli AC, Michael BH, Karen LL. Seasonal change in nutrient composition of spotted knapweed and preference by sheep. *Small Ruminant Research*. 2010;**89**:47-50

[44] Gutman M, Perevolotsky A, Sternberg M. Grazing effects on a perennial legume, *Bituminaria bituminosa* (L.) Stirton, in a mediterranean rangeland. *Cahiers Options Mediterranean*. 2000;**45**:299-303

[45] Martinez-Loperena R, Castelan-Ortega OA, Gonzalez-Ronquillo M, Estrada-Flores JG. Nutritive value, in vitro fermentation and secondary metabolites of weeds and maize straw used for feeding dairy cattle. *Tropical and Subtropical Agroecosystems*. 2011;**14**:525-536

[46] Kirilov A, Georgieva N, Stoycheva I. Determination of composition and palatability of certain weeds. *International Journal of Agricultural Science and Food Technology*. 2016;**2**:41-43

[47] Miller JK, Brzezinska-Slebodzinska E. Oxidative stress, antioxidants, and animal

function. Journal of Dairy Science.
1993;**76**:2812-2823

[48] Olson BE, Wallander RT, Lacey JR. Effects of sheep grazing on spotted knapweed-infested Idaho fescue community. Journal of Range Management. 1997;**50**:386-390

[49] Iqbal MA, Asif I, Nadeem A, Haroon ZK, Rana NA. A study on feed stuffs role in enhancing the productivity of milch animals in Pakistan-existing scenario and future prospect. Global Veterinária. 2015;**14**:23-33

[50] Shale TL, Stirk WA, Van SJ. Variation in antibacterial and anti-inflammatory activity of different growth forms of *Malva parviflora* and evidence for synergism of the anti-inflammatory compounds. Journal of Ethnopharmacology. 2005;**96**:325-330

[51] Tremmel DC, Patterson DT. Responses of soybean and five weeds to CO₂ enrichment under two temperature regimes. Canadian Journal of Plant Science. 1993;**73**:1249-1260

[52] Iqbal MA, Abdul H, Tanveer A, Imtiaz H, Sajid A, Ansar A, et al. Forage sorghum-legumes intercropping: Effect on growth, yields, nutritional quality and economic returns. Bragantia. 2019;**78**:82-95

[53] Iqbal MA, Abdul H, Muzammil HS, Imtiaz H, Tanveer A, Saira I, et al. A meta-analysis of the impact of foliar feeding of micronutrients on productivity and revenue generation of forage crops. Planta Daninha. 2019;**37**:e019189237

[54] Akdeniz H, Mehmet AB, Ibrahim H, Mohammad SI, Akbar H, Mabrouk S, et al. Yield and nutritive value of Italian ryegrass (*Lolium italicum* L.) is influenced by different levels of nitrogenous fertilizer. Fresenius Environmental Bulletin. 2019;**28**:8986-8992