

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,900

Open access books available

186,000

International authors and editors

200M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



## Chapter

# *Lecanicillium* spp. for the Management of Aphids, Whiteflies, Thrips, Scales and Mealy Bugs: Review

Sajjalavarahalli Gangireddy Eswara Reddy

## Abstract

*Lecanicillium* spp. are potential microbial bio-control agent mainly used for the management of sucking insect pests such as aphids, whiteflies, scales, mealy bugs etc. and gaining much importance at present for management of pests. Due to indiscriminate use of chemical pesticides which results in development of resistance, resurgence, outbreak of pests and residue problem, the farmers/growers are forced to use bio-pesticides for sustainable agriculture. *Lecanicillium* spp. is promising biocontrol agent against sucking insect pests and can be used as one of the components in integrated pest management (IPM). However, optimum temperature and relative humidity are the major environmental factors, for the performance of *Lecanicillium* spp. under protected/field conditions. The present review is mainly focused on nomenclature of *Lecanicillium* spp., mode of infection, natural occurrence, influence of temperature and humidity on the growth, factors influencing the efficacy, virulence/pathogenicity to target pests, substrates used for mass production, safety to non-target organisms, compatibility with agrochemicals and commercially available products. This review is mainly useful for the researchers/students to plan their future work on *Lecanicillium* spp.

**Keywords:** entomopathogenic fungus, aphid, whitefly, virulence, mass production, safety

## 1. Introduction

The increased use of conventional chemical pesticides over the years has not only contributed to an increase in food production, but also has resulted in adverse effects on the environment and non-target organisms. In view of these side effects, the necessity for sustainable crop production through ecofriendly pest management technique is being largely felt in the recent times. Few biopesticides are available in the market, among them *Lecanicillium* spp. based microbial bio-pesticide gaining much importance for sucking pests for organic and sustainable agriculture [1–4]. Myco pesticides are potential microbial alternative to chemical pesticides and offer a number of benefits such as facility of growth on a variety of substrates, high virulence, trans cuticular penetration, broad host range, less expensive, safe to humans, animals and the environment. Therefore, this review is prepared by compiling the research work done on *Lecanicillium* spp. by various research groups on various

aspects viz., nomenclature, mode of infection, natural occurrence, effect of temperature and humidity for the growth, factors influencing its efficacy, virulence and pathogenicity against target pests under laboratory/greenhouse/field, substrates used for mass production, safety to non-target organisms, compatibility with agrochemicals and commercially available products were discussed and presented.

## 2. Nomenclature of *Lecanicillium* spp.


The genus *Verticillium* contains diverse host ranges including arthropods, nematodes, plants and fungi [5]. The genus has been redefined using rDNA sequencing, grouping insect pathogens into the new genus *Lecanicillium* which includes *L. attenuatum*, *L. lecanii*, *L. longisporum*, *L. muscarium* and *L. nodulosum*, which were all formerly classified as *V. lecanii* [5–7].

## 3. Mode of infection


When *L. lecanii* conidia comes in contact with the host integument, it gets adhere to the epicuticle and germinate. Germinated conidia form germ tubes that penetrate cuticle directly or grow over the surface of the epicuticle. The germ tube penetrates by lysing both the epicuticle and the procuticle [8, 9]. This is accomplished by the mechanical pressure exerted by appressorium (penetration peg) and secretion of enzymes viz., proteases, chitinases and esterase's which plays an important role during cuticle penetration of insect host and also serve as cuticle degrading enzymes. The fungus proliferates throughout the insect's body, draining the insect of nutrients, and eventually killing it in around 48–72 hours. The mycotoxins produced by *L. lecanii* are bassianolide [10, 11], vertilecanin-A1, decenedioic acid and 10-hydroxy-8-decenoic acid) [12–14]. As the host nutrients are depleted, the blastopores' differentiate into elongated hyphae which extend outward from the body forming a mycelial mat of conidiophores over the surface of the integument resulting in mummification. Under favourable environmental condition, conidiophores mature giving rise to conidia which continues the disease cycle further.

## 4. R & D publications on different aspects of *Lecanicillium* spp.


The number of publications related to *Lecanicillium* spp. from 1971 to 2020 was presented in the **Figures 1** and **2**. The data clearly indicated that, during 1971–80<sup>s</sup>



**Figure 1.**  
Per cent R & D publications related to *Lecanicillium* spp.



**Figure 2.**  
 R & D publications on *Lecanicillium spp.*



**Figure 3.**  
 R & D publications on various aspects of *Lecanicillium spp.*

the publications were completely nil, but during 1981–91<sup>s</sup>, the R&D work has been initiated in the entire world and the publications were increased gradually reaching 58% during 2001–2020 (**Figure 2**). While, considering the number publications on various aspects of *Lecanicillium spp.*, more research work has been done on virulence and pathogenicity (**Figure 3**) followed by biotechnology and biochemistry as compared to morphology, diversity, ecology, mass production. The number of publications was meagre on effect of environmental factors (temperature and humidity), safety to natural enemies and compatibility with pesticides [15].

### 5. Natural occurrence of *Lecanicillium spp.*

*Lecanicillium spp.* is the most widely distributed and generally found on infected insects both in temperate and tropical areas throughout the world. There are number of reports on natural infection of *Lecanicillium spp.* on different insect pests but out of the reported insects and pests, maximum are sucking pests belonging to Hemiptera, Thysanoptera and Acarina which indicates its possible spectrum for use as a biocontrol agent for pest management. Reports of natural occurrence of *Lecanicillium spp.* on sucking insects presented in the **Table 1**.

Strain/isolate	Host	Location	Reference
<i>L. lecanii</i> (Is-2, Is-5)	<i>M. persicae</i>	Israel	[14]
<i>L. lecanii</i> (Is-6)	<i>Acrithosiphon pisum</i>	Israel	[14]
<i>L. lecanii</i> (R-1)	<i>T. vaporariorum</i>	Russia	[14]

Strain/isolate	Host	Location	Reference
<i>L. lecanii</i> (V16063)	<i>T. vaporariorum</i>	Halifax, Canada	[2]
<i>L. lecanii</i> (V0175)	<i>B. tabaci</i>	Guangdong, China	[2]
<i>L. lecanii</i> (Vp28)	<i>Pseudococcus</i> sp.	Guangdong, China	[2]
<i>L. lecanii</i> (ICAL4)	<i>Nasonovia ribisnigri</i> in lettuce	Madrid	[16]
<i>L. lecanii</i> (ICAL6)	<i>M. persicae</i> in pepper	Madrid	[16]
<i>L. lecanii</i> (41185)	<i>M. persicae</i> , <i>T. vaporariorum</i>	Korea	[17]
<i>L. longisporum</i> (6541)	<i>Aphis gossypii</i>	UK	[17]
<i>L. longisporum</i> (6543)	<i>M. persicae</i>	UK	[17]
<i>L. longisporum</i> (4078)	<i>M. persicae</i>	Denmark	[17]
<i>L. longisporum</i> (HRI 1.72)	<i>Macrosiphoniella sanbornii</i>	UK	[18]
<i>L. lecanii</i> (ARSEF 7207)	<i>T. vaporariorum</i>	Argentina	[16]
<i>L. longisporum</i> (ARSEF 7461)	<i>T. vaporariorum</i>	Argentina	[16]
<i>L. muscarium</i> (ARSEF 7460)	<i>T. vaporariorum</i>	Argentina	[16]
<i>L. lecanii</i> (ICAL3)	<i>Macrosiphum euphorbiae</i> in tomato	Madrid, Spain	[19]
<i>L. lecanii</i> (ITEM 3757)	<i>Brevicorne brassicae</i> in Cabbage	Bari, Italy	[20]
<i>L. lecanii</i>	<i>S. bispinosus</i> on tea	Tamil Nadu, India	[21]
<i>L. sabanense</i> sp. nov	<i>Pulvinaria caballeroramose</i>	Bogota (Columbia)	[22]
<i>L. attenuatum</i> ZJLSP07 and <i>L. psalliotae</i> ZJLA08	<i>Diaphorina citri</i>	Taizhou (Zhejiang Province, China)	[23]
<i>L. lecanii</i> (FI 2482) and <i>L. muscarium</i> (FI 2481)	<i>Thaumastocoris peregrinus</i>	South-East Uruguay	[24]

**Table 1.**  
Natural occurrence of *Lecanicillium* spp. on different sucking insect pests.

## 6. Effect of temperature and humidity on the growth of *Lecanicillium* spp.

Temperature and humidity are the main factors influencing the growth of the fungus. Effect of different temperature on conidial germination, growth rate, colony size and mycelial growth was discussed and presented in **Table 2**.

Temperature	5 °C	10 °C	15 °C	20 °C	25 °C	30 °C
Water activity (aw)	—	0.985	0.99	0.98	0.975	[25]
Strain/isolate	% Conidial germination					
<i>L. longisporum</i> (Vertalec)	—	—	98	98	28.7	—
<i>L. muscarium</i> (Mycotal)	—	20.6	98	98	98	—
PFC 1	—	—	—	50.6	47	—
PFC 3	—	—	—	49.7	86.6	—

Temperature	5 °C	10 °C	15 °C	20 °C	25 °C	30 °C	
PFC 10	—	64.7	47.7	14.7	—	—	
PFC 11	—	—	98	98	49.3	—	
PFC 13	—	88	98	98	98	—	
Mean radial growth rates (mm/day)							
Strain/isolate	5 °C	10 °C	15 °C	20 °C	25 °C	30 °C	[25]
<i>L. longisporum</i> (Vertalec)	0.21	0.66	1.10	1.31	1.86	0.55	
<i>L. muscarium</i> (Mycotal)	0.22	0.59	1.03	1.59	2.03	0.59	
PFC 1	0.16	0.43	0.90	1.13	1.64	0.69	
PFC 3	0.15	0.54	1.03	1.35	1.86	0.83	
PFC 10	0.18	0.63	1.02	1.40	2.07	0.05	
PFC 11	0.17	0.58	1.03	1.25	2.05	0.05	
Mean colony size (diameter: mm)							
Strain/isolate	5 °C	10 °C	15 °C	20 °C	25 °C	30 °C	[26]
Vertalec	5.0	18.6	34.1	50.2	52.1	5.0	
Mycotal	12.1	20.5	31.5	42.1	47.3	8.3	
B-2	11.6	21.3	25.4	46.2	53.6	26.9	

**Table 2.**  
 Effect of temperature on growth of *Lecanicillium spp.*

## 6.1 Temperature

Temperature affects the *Lecanicillium spp.* in different ways by influencing the germination, growth and viability of the fungus in the host insect and environment. High temperature inactivates the fungus before contact with the pest insect or may reduce or accelerate the growth within an insect depending on the temperature requirements of the fungus and the host insect. In contrast, low temperatures reduce or stop the germination and growth. Optimal germination and growth rates of *Lecanicillium spp.* range between 23°C and 28°C, growth rapidly slows >30°C and ceases at 34 to 37°C. Similarly, conidial germination is adversely affected by temperatures above 30°C. Temperature below 16°C increasingly slows germination and growth and thus affects efficacy in terms of a longer survival of the target population. *Lecanicillium* strains showed optimum growth at 25°C; the aerial conidia of *Lecanicillium* strains germinate in a broad temperature range (15–30°C) and *L. lecanii* 41,185 was the only strain with conidial germination at 35°C [27].

Effect of different temperature on conidial germination, growth rate, colony size and mycelial growth of *L. lecanii* was discussed and presented in **Table 2**. At 25°C and 0.975  $a_w$  (water activity) conidial germination occurred in all the isolates ranging from 28.7 to 98% whereas isolate PFC 10 no conidial germination had. Per cent germination decreased from highest values at 25°C to the lowest trend at 10°C in Mycotol (20.6°C). Maximum germination of conidia was observed between 15 and 25°C [25]. Most of the isolates showed growth at 5 and 30°C and mean growth rate increased as temperature increased. Optimum growth rate occurred at 25°C (1.64 to 2.07 mm) for all isolates) [25]. Colony size of the fungus was influenced

by the temperature, the colony growth is maximum at 25°C (47.3 to 53.6 mm) as compared to the temperature between 5 to 20°C [26]. The optimum temperature for the mycelial growth of *L. lecanii* CA-1-G was 23°C (37.57 mg/cm<sup>2</sup>) and 26°C (39.43 mg/cm<sup>2</sup>) as compared to 20°C (29.43 mg/cm<sup>2</sup>), 29°C (20.7 mg/cm<sup>2</sup>) and 32°C (20.63 mg/cm<sup>2</sup>). Similarly, *L. lecanii* grew and sporulated over a wide range of temperatures (20–32°C). The optimum temperature for growth was 23°C (46.45 x10<sup>5</sup> conidia cm<sup>-2</sup>) or 26°C (33.76 x10<sup>5</sup> conidia cm<sup>-2</sup>) for *L. lecanii* CA-1-G [28]. Virulence of *Lecanicillium* spp. isolates was evaluated against third instar *T. vaporariorum* on tomato plants at 23°C. Colony radial growth, conidial production and germination decreased with the reduction in water activity, while 32°C was extremely detrimental for all fungal isolates. However, some isolates were able to grow and produce conidia at low water activity and high temperature [29]. *L. muscarium* can multiply in temperature range of 15–30°C but optimum temperature against *M. persicae* between 20 to 30°C [30].

## 6.2 Humidity

Humidity is another important environmental factor affecting the efficacy and survival of *Lecanicillium*. Spore germination on the insect cuticle and sporulation after outgrowth of the dead host insect require high moisture. Generally high humidity is required for germination of spores under in vitro, insects can become infected at much lower humidity. Under fluctuating humidity, daily saturated humidity requirement of at least 16 h for causing death in *Trialeurodes vaporariorum* (Westwood) infected with *L. lecanii* [31]. Several previous studies provided evidence that a threshold time period at high humidity was required for infection. Conidia of *L. lecanii* required at least 72 h at 100% RH and 20°C before removal to 70% RH to reach >90% infectivity of *Myzus persicae* (Sulzer) [32]. Similarly, at 25°C temperature and 75% relative humidity (RH), *L. lecanii* 41,185 showed highly virulent pathogenicity (100%) against *M. persicae* and *Aphis gossypii* Glover [27]. Application of *L. longisporum* against *A. gossypii* on cucumber in controlled environment (Temperature; 19–26°C and humidity; 80–98%) resulted in 100% mortality [32, 33]. *L. muscarium* grow at optimum temperature but higher mortality observed against *M. persicae* between 55 and 90% humidity [30].

## 7. Factors influencing the efficacy of *Lecanicillium* spp. against sucking insect pests

The virulence and pathogenicity of *Lecanicillium* spp. vary with strain, stage of the insect and dose of the fungus.

### 7.1 Strains

Virulence of the *Lecanicillium* spp. varies with strain to strain or isolate to isolate. The isolate ICAL6 was more virulent (LC<sub>50</sub> = 1.05 x 10<sup>7</sup> conidia mL<sup>-1</sup>) to nymphs of *M. persicae* than *Macrosiphum euphorbiae* (Thomas) (LC<sub>50</sub> = 1.26 x 10<sup>7</sup> conidia mL<sup>-1</sup>) and *Nasonovia ribisnigri* (Mosley) (LC<sub>50</sub> = 2.78 x 10<sup>7</sup> conidia mL<sup>-1</sup>) [19]. The strain VI 6063 imported from Canada was more virulent to *Bemisia tabaci* (Gennadius) (2.57 x 10<sup>5</sup> conidia mL<sup>-1</sup>) than the domestic strains V3450 and Vp28 (LC<sub>50</sub> = 6.03 x 10<sup>5</sup> conidia mL<sup>-1</sup>) [2]. *L. lecanii* @ 1x10<sup>7</sup> conidia mL<sup>-1</sup> is more effective against nymphs of *Plannococcus citri* (84% mortality) after six days

of treatment as compared to *L. longisporum* (59% mortality) [34]. *L. muscarium* isolate FI 2481 @  $1 \times 10^7$  conidia  $\text{mL}^{-1}$  was more effective against *Thaumastocoris peregrinus* (72% mortality) as compared to *L. lecanii* isolate FI 2482 which reported 50% mortality [24]. Similarly, *L. lecanii* hybrid strain 2aF4 was more promising ( $\text{LC}_{50} = 5.3 \times 10^4$  conidia  $\text{mL}^{-1}$ ) for the management of *Trialeurodes vaporariorum* than *L. lecanii* 2aF4 ( $\text{LC}_{50} = 7.8 \times 10^4$  conidia  $\text{mL}^{-1}$ ) [35].

## 7.2 Stage of the insect

Stage of host plays important role in the success of *Lecanicillium* spp. and not all stages of insect life cycle are equally susceptible to fungal infection. So, the fungal application can be successful against the particular pest when it can be done at the condition where the susceptible stage or weaker stages of the particular pest become dominant among population.

First and third instar nymphs of *B. tabaci* (38 and 65% mortality) were significantly more susceptible to *L. muscarium* than the fourth instar (15%) in verbena plants. Similarly, first and second instars *B. tabaci* was more susceptible (50 and 55% mortality) than the third and fourth instar (25 and 20% mortality) on tomato foliage [36]. *L. lecanii* (ARSEF 7460) showed higher mortality against nymphs of *T. vaporariorum* followed by *L. longisporum* (ARSEF 7207) and *L. muscarium* (ARSEF 74601) @  $1 \times 10^7$  conidia  $\text{mL}^{-1}$  [16]. The pathogenicity of *L. lecanii* strains was more in pupae (59–72.5%) than adults (34–52.6%) after 6 days of inoculation [14]. *L. lecanii* ( $2.8 \times 10^7$  conidia/ml) isolated from *Scirtothrips bispinosus* (Bagnall) in tea showed higher mortality against larvae (60%) than adults (30%) of *S. bispinosus* under laboratory at same dose [21]. Mortality of nymphs of *Plannococcus citri* were more susceptible (84% mortality) after six days of treatment to *L. lecanii* @  $1 \times 10^7$  conidia  $\text{mL}^{-1}$  as compared to adults which showed 40% mortality [34]. *L. lecanii* hybrid strain 2aF43 @  $1 \times 10^7$  conidia  $\text{mL}^{-1}$  showed more efficacy against first instar nymphs of *T. vaporariorum* (68% mortality) as compared to 4th instar nymphs (30% mortality) and adults (60% mortality). Similarly, *L. lecanii* hybrid strain 2aF4 is more effective against first instar nymphs (46% mortality) as compared to 4th instar nymphs (30% mortality) [35].

## 7.3 Dose/inoculum level

Fungal inoculum level is the important factor which affects the performance. It is general trend that the higher fungal inoculum level gives higher insect mortality. However, sufficient inoculum level should be worked out for the particular pest to prevent the over inoculum wastage and to achieve higher mortality. Higher dose of *L. lecanii* ( $1.2 \times 10^9$  conidia  $\text{ha}^{-1}$ ) was caused 92.30 and 80.93% mortality of *Brevicoryne brassicae* Linnaeus and *Aleurodicus disperses* (Russell) respectively at 10 days after treatment in the laboratory, whereas in field conditions *L. lecanii* (V13) at  $2 \times 10^{12}$  conidia  $\text{ha}^{-1}$  causing 61.16% and 66.50% mortality of *B. brassicae* and *A. craccivora* respectively [2].

## 8. Efficacy of *Lecanicillium* spp. against sucking pests under laboratory/greenhouse/field

Efficacy of *Lecanicillium* spp., against aphids, whiteflies, thrips, scales and mealy bugs in the laboratory/greenhouse/field conditions w.r.to its mortality,  $\text{LC}_{50}$  and  $\text{LT}_{50}$  values were presented in the **Table 3**.

Strain/isolate	Conditions (Lab, GH, F)	Pest	Mortality/LC <sub>50</sub> /LT <sub>50</sub>	Temperature (°C)	Humidity (%)	References
<i>Lecanicillium lecanii</i>	Lab	<i>Bemisia argentifolii</i>	95–98%	20–25	100	[14]
<i>L. lecanii</i> (HRI 1.72)	Lab	<i>A. fabae</i>	LT <sub>50</sub> (2.79 d)	10–23		[23]
<i>L. lecanii</i> (HRI 1.72)	Lab	<i>M. persicae</i>	LT <sub>50</sub> (3.39 d)	23		[37]
<i>L. lecanii</i> (V16063)	Lab	<i>B. tabaci</i>	(94.9%) LC <sub>50</sub> = 2.57 x 10 <sup>5</sup> Conidia mL <sup>-1</sup>	25	95	[2]
<i>L. lecanii</i> (V3450)	Lab	<i>B. tabaci</i>	86.9 (LC <sub>50</sub> = 6.03 x 10 <sup>5</sup> conidia mL <sup>-1</sup> )	25	95	[2]
<i>L. longisporum</i>		<i>M. persicae</i> , <i>Macrosiphum euphorbiae</i> , <i>Aulacorthum solani</i>	(LT <sub>50</sub> = 2.4;1.8; 2.0 d) 100% mortality	25	95	[38]
<i>L. longisporum</i> (HRI 1.72)	Lab	<i>M. persicae</i> , <i>A. fabae</i> , <i>Acrithosiphon pisum</i> , <i>Sitobion avenae</i>	LT <sub>50</sub> = 74–78 h			[18]
<i>L. longisporum</i>	Cucumber	<i>A. gossypii</i>	100% (LT <sub>50</sub> = 6.9 d)	25.8	80.6	[33]
<i>L. longisporum</i> or <i>L. muscarium</i>	Lab	<i>Frankliniella occidentalis</i>	95%	20	70%	[39]
<i>L. lecanii</i>	Lab	<i>A. craccivora</i>	(LC <sub>50</sub> = 2.5 x 10 <sup>4</sup> spores mL <sup>-1</sup> ) (LT <sub>50</sub> = 3.9 x 10 <sup>8</sup> spores mL <sup>-1</sup> )			[40]
<i>L. muscarium</i> (1x10 <sup>7</sup> spores/ml)	Verbana, tomato (GH)	<i>B. tabaci</i>	65 and 55% mortality	20	95	[41]
<i>L. muscarium</i> (1x10 <sup>7</sup> conidia/ml)	Verbana, tomato (GH)	<i>B. tabaci</i>	85 and 80%	20	85	[36]
<i>L. longisporum</i>	Cucumber (GH)	<i>A. gossypii</i>	100%	19.0	80.2	[42]
<i>L. lecanii</i>	Tea (F)	<i>S. bispinosus</i>	30–60%	—	—	[21]
<i>L. attenuatum</i> ZJLSP07 and <i>L. psalliotae</i> ZJLA08	Lab	<i>Diaphorina citri</i>	100% (1x10 <sup>8</sup> conidia/ml)	25	90	[23]
<i>L. attenuatum</i> (SD-16, SDMP1 and 2)	Lab	<i>M. persicae</i>	100%	25	>90	[43]

Strain/isolate	Conditions (Lab, GH, F)	Pest	Mortality/LC <sub>50</sub> /LT <sub>50</sub>	Temperature (°C)	Humidity (%)	References
<i>L. lecanii</i>	Lab	<i>M. persicae</i> , <i>A. gossypii</i>	100% (1x10 <sup>8</sup> conidia/ml)	20	>90	[44]
<i>L. lecanii</i> (JMC-01)	Lab	<i>B. tabaci</i>	82.2% (1x10 <sup>8</sup> conidia/ml)	25	70	[45]
<i>L. lecanii</i> (FI 2482) and <i>L. muscarium</i> (FI 2481)	Lab	<i>Thaumastocoris peregrinus</i>	50 and 72%	25	65	[24]
<i>L. lecanii</i> 2aF4 3 and 2aF4	Lab	<i>T. vaporariorum</i>	83% (LC <sub>50</sub> = 5.3x10 <sup>4</sup> conidia/ml) and 84% (LC <sub>50</sub> = 7.8 x10 <sup>4</sup> conidia/ml)	23	99.6	[35]

GH; Green house, F; Field, LC<sub>50</sub>; Lethal concentration to kill 50% insects, LT<sub>50</sub>; Lethal time to kill 50% insects.

**Table 3.**  
Efficacy of *Lecanicillium* spp. against sucking pests under laboratory/greenhouse/field.

## 9. Substrates used for mass production of *Lecanicillium* spp.

*Lecanicillium* spp. can be mass multiplied by solid state fermentation (SSF) and liquid state fermentation (LSF) using different growth media. In SSF, different grains, agars and non-synthetic solid media were used for mass production of *Lecanicillium* spp. (Table 4).

Substrates	Conidia/Spores	References
<b>Media</b>		
Sabaroud dextrose agar	$2.87 \times 10^7$ conidia $\text{cm}^{-2}$	[46]
Malt extract agar	$5.23 \times 10^7$ conidia $\text{cm}^{-2}$	
Nutrient agar	$1.07 \times 10^7$ conidia $\text{cm}^{-2}$	
Corn meal agar	$0.09 \times 10^7$ conidia $\text{cm}^{-2}$	
Yeast peptone dextrose agar	$4.58 \times 10^7$ conidia $\text{cm}^{-2}$	
Potato dextrose agar	$2.91 \times 10^7$ conidia $\text{cm}^{-2}$	
<b>Grains</b>		
Rice	$8.43 \times 10^8$ spores $\text{g}^{-1}$	[47]
Wheat	$9.13 \times 10^8$ spores $\text{g}^{-1}$	
Sorghum	$11.31 \times 10^8$ spores $\text{g}^{-1}$	
Pearl millet	$10.17 \times 10^8$ spores $\text{g}^{-1}$	
Finger millet	$9.76 \times 10^8$ spores $\text{g}^{-1}$	
Maize	$7.54 \times 10^8$ spores $\text{g}^{-1}$	
Rice	$1.97 \times 10^9$ spores $\text{g}^{-1}$	[48]
Sorghum	$1.90 \times 10^9$ spores $\text{g}^{-1}$	
Finger millet	$1.66 \times 10^9$ spores $\text{g}^{-1}$	
Wheat	$1.65 \times 10^9$ spores $\text{g}^{-1}$	
Corn	$1.84 \times 10^9$ spores $\text{g}^{-1}$	
Polished rice	$5.7 \times 10^8$ conidia $\text{g}^{-1}$	[49]
Cooked rice	$1.5 \times 10^9$ conidia $\text{g}^{-1}$	
Rice bran	$1.4 \times 10^9$ conidia $\text{g}^{-1}$	
Crushed bajra +1% yeast extract (YE)	$17.49 \times 10^8$ conidia $\text{g}^{-1}$	[4]
Crushed sorghum +1% YE	$10.34 \times 10^8$ conidia $\text{g}^{-1}$	
Crushed navane +1% YE	$3.52 \times 10^8$ conidia $\text{g}^{-1}$	
Crushed maize +1% YE	$4.80 \times 10^8$ conidia $\text{g}^{-1}$	
Crushed rice +1% YE	$24.59 \times 10^8$ conidia $\text{g}^{-1}$	
Crushed wheat +1% YE	$3.54 \times 10^8$ conidia $\text{g}^{-1}$	
Rice bran	$24 \times 10^7$ conidia $\text{g}^{-1}$	[34]
<b>Agro wastes</b>		
Crushed maize cobs +10% molasses	$10.07 \times 10^4$ conidia $\text{g}^{-1}$	[4]
Wheat bran +10% molasses	$18.76 \times 10^4$ conidia $\text{g}^{-1}$	
Rice bran +10% molasses	$30.86 \times 10^4$ conidia $\text{g}^{-1}$	
Baggase +10% molasses	$10.88 \times 10^4$ conidia $\text{g}^{-1}$	

Substrates	Conidia/Spores	References
Press mud +10% molasses	$7.90 \times 10^4$ conidia $g^{-1}$	
Sugarcane molasses 3%	$8.35 \times 10^8$ spores $ml^{-1}$	[48]
Sugarcane molasses 4%	$8.56 \times 10^8$ spores $ml^{-1}$	
Sugarcane molasses 5%	$8.42 \times 10^8$ spores $ml^{-1}$	
<b>Non synthetic solid media</b>		
Carrot	$2.17 \times 10^8$ spores $g^{-1}$	[47]
Jack seeds	$4.11 \times 10^8$ spores $g^{-1}$	
Ladies finger	$3.12 \times 10^8$ spores $g^{-1}$	
Rice husk	$1.27 \times 10^8$ spores $g^{-1}$	
Saw dust	$0.69 \times 10^8$ spores $g^{-1}$	
Beet pulp	$23 \times 10^7$ conidia $g^{-1}$	[34]
<b>Non synthetic liquid media</b>		
Coconut water	$5.27 \times 10^8$ spores $g^{-1}$	[47]
Rice cooked water	$2.11 \times 10^8$ spores $g^{-1}$	
Rice wash water	$3.12 \times 10^8$ spores $g^{-1}$	
Wheat wash water	$1.21 \times 10^8$ spores $g^{-1}$	
<b>Liquid media</b>		
Potato carrot broth	$6.50 \times 10^7$ spores $mL^{-1}$	[48]
Potato dextrose broth	$3.95 \times 10^7$ spores $mL^{-1}$	
Potato sucrose broth	$6.30 \times 10^7$ spores $mL^{-1}$	
Jaggery yeast broth	$2.45 \times 10^7$ spores $mL^{-1}$	
Sucrose yeast broth	$2.50 \times 10^7$ spores $mL^{-1}$	
Molasses yeast broth	$8.33 \times 10^7$ spores $mL^{-1}$	

**Table 4.**  
*Substrates (media, grains, agro wastes) used for mass production of Lecanicillium spp.*

Among grains, rice is most suitable substrates for mass production ( $1.97 \times 10^9$  spores  $g^{-1}$ ) followed by sorghum ( $1.90 \times 10^9$  spores  $g^{-1}$ ) as compared to finger millet, wheat and corn ( $1.6\text{--}1.80 \times 10^9$  spores  $g^{-1}$ ) [48]. Similarly, crushed rice +1% yeast extract recorded higher spore yield ( $24.59 \times 10^8$  conidia  $g^{-1}$ ) followed by crushed bajra +1% yeast extract ( $17.49 \times 10^8$  spores  $g^{-1}$ ) as compared to crushed sorghum, maize and wheat [4].

Among different agro wastes used for multiplication of *L. lecanii*, the growth and sporulation were found to be better on rice bran +10% molasses ( $30.86 \times 10^4$  conidia  $g^{-1}$ ) followed by wheat bran +10% molasses ( $18.76 \times 10^4$  conidia  $g^{-1}$ ) and rice bran ( $15.98 \times 10^4$  conidia  $g^{-1}$ ). Complete inhibition of growth and reproduction of the fungus was noticed on bagasse and pressmud with 1 per cent yeast extract alone. However, growth was recorded when bagasse and press mud was supplemented with 10% molasses ( $10.88$  and  $7.90$  conidia  $g^{-1}$  respectively) [4]. Among agars, malt extract agar (MEA) yields high conidia production ( $5.23 \times 10^7$  conidia  $cm^{-2}$ ) followed yeast peptone dextrose agar ( $4.58 \times 10^7$  conidia  $cm^{-2}$ ) as compared to potato dextrose agar and sabaroud dextrose agar ( $2.91$  and  $2.87 \times 10^7$  conidia  $cm^{-2}$  respectively) [46]. In non-synthetic media, jack seeds produced high spore yield ( $4.11 \times 10^8$  spores  $g^{-1}$ ) followed by ladies' finger ( $3.12 \times 10^8$  spores  $g^{-1}$ ), carrot ( $2.17 \times 10^8$  spores  $g^{-1}$ ) and rice husk ( $1.27 \times 10^8$  spores  $g^{-1}$ ) [47].

In LSF, molasses yeast broth (MYB) supported maximum spore production of *L. lecanii* ( $8.33 \times 10^7$  spores  $\text{ml}^{-1}$ ) followed by potato carrot broth ( $6.5 \times 10^7$  spores/ml) and potato sucrose broth ( $6.3 \times 10^7$  spores  $\text{ml}^{-1}$ ) as compared to Sucrose yeast broth, Jaggery yeast broth and Potato dextrose broth ( $2.45\text{--}3.95 \times 10^7$  spores  $\text{mL}^{-1}$ ). Among non-synthetic liquid media, coconut water produced higher spores ( $5.27 \times 10^7$  spore's  $\text{g}^{-1}$ ) and biomass production than rice wash water ( $3.12 \times 10^7$  spore's  $\text{g}^{-1}$ ) as compared to rice cooked water and wheat wash water ( $1.21\text{--}2.11 \times 10^7$  spores  $\text{g}^{-1}$ ) [24]. The growth of *L. longisporum* conidial spores are higher in rice bran [ $24 \times 10^7$  conidia  $\text{g}^{-1}$ ] as compared to beet pulp [ $23 \times 10^7$  conidia  $\text{g}^{-1}$ ] [34].

## 10. Safety of *Lecanicillium* spp. to parasitoids/predators/pollinators

The safety of any bio control agent to parasitoids/predators/pollinators is the important aspect which should be studied thoroughly before its commercialization to avoid the hazards and disturbance of ecological balance. Effect of *L. lecanii* on aphid parasitoid *Aphidius colemani* (Viereck) which showed the normal development (approximately 90% adult emergence) when its cotton aphid, *A. gossypii* host was treated with *L. lecanii* conidia 5 or 7 days after parasitization. Fungus exposure 1 day before or up to 3 days after parasitization, however, reduced *Aphidius colemani* (Viereck) emergence from 0 to 10%. They suggested that the parasitoid and fungus may be used together for aphid bio control [50]. *L. lecanii* showed pathogenicity against predatory mite, *Phytoseiulus persimilis* Athias-Henriot but its effect was lower than that of spider mite, *Tetranychus urticae* (Koch) [51]. *L. lecanii* is safer to predatory coccinellid, *Coccinella septempunctata* Linnaeus and predatory mites, *Amblyseius ovalis* (Evans) and *Amblyseius longispinosus* (Evans) under field conditions [52]. The fungus *L. lecanii* was not pathogenic to *Chrysoperla carnea* (Stephens), *Coccinella septempunctata* (Linnaeus), *Episyrphus balteatus* (De Geer) and *Samia cynthia ricini* (Boisduval), but was found to be pathogenic to *Bombyx mori* (Linnaeus). Parasitism, adult emergence and adult longevity of *Trichogramma chilonis* (Ishii) were affected by fungal treatments. Aphid mummification and *Diaeretiella rapae* adult emergence were affected by the fungus. Results suggest that *L. lecanii* is compatible with natural enemies of cabbage aphid, *T. chilonis* and is harmless to silk worm [53]. *L. muscarium* at  $10^6$  and  $10^7$  spores  $\text{mL}^{-1}$  was safer to predatory mite *P. persimilis* [54]. Number of parasitized larvae of *Eretmocerus sp. nr. furuhashii* survival decreased with increasing concentrations of *L. muscarium* and only 29% emergence of pupae was observed at a conidial concentration of  $1 \times 10^8$  conidia  $\text{mL}^{-1}$ . Similarly, 67% emergence of adult *E. sp. nr. Furuhashii* was observed [55]. Parasitoid (*Diaeretiella rapae*) emergence was affected by application of *L. longisporum* before or after parasitism and longevity decreased in female F1 populations [56]. In the laboratory conditions, application of *L. muscarium* ( $1 \times 10^8$  conidia/ml) against *A. colemani* had not affected longevity and fertility of the female *A. colemani*. The combination of *Aphidius colemani* with *L. muscarium* reduced the aphid infestation in the semi field conditions as compared to *A. colemani* alone [30].

The *Lecanicillium* spp. is not harmful to humans during handling in the laboratory and field for the control of pests.

## 11. Compatibility of *Lecanicillium* spp. with agro chemicals

Chemical pesticides may have antagonistic or synergistic effect on the potentiality of *Lecanicillium* spp. and may disrupt natural epizootic. Under such epizootic

condition, it is expected to enhance effectiveness through joint action of pathogen and compatible insecticides, which would reduce not only the cost of protection but also reduce the contamination of the environment. The literature on compatibility of *Lecanicillium* spp. with agrochemicals is lacking.

Among different insecticides studied for their effect on *L. lecanii* under in-vitro, malathion was significantly detrimental (69.18% inhibition) than all other insecticides except quinalphos (66.76%). Conversely, endosulfon and chlorpyrifos were significantly safer (37.31 to 44.37%), followed by oxydemeton methyl and dimethoate (45.33 to 48.27% inhibition) [4]. Similarly, endosulfan completely inhibited the germination of conidia and hyphal growth. In contrast, diafenthiuron, thiamethoxam, imidacloprid, thiodicarb, primicarb, omethoate, acetamiprid, and pymetrozine were compatible with *L. lecanii* in planta [57]. Imidacloprid and cyromazine were compatible with *L. lecanii* in terms of vegetative growth, sporulation, conidial viability and pathogenicity against *T. urticae*. At the recommended concentration, the fungicides carbendazim, chlorothalonil, propiconazole, mancozeb and wettable sulphur completely inhibited the germination of candida (100%) except iprodione and triadimefom allowed 37.38 and 41.62% conidia to germinate respectively [4].

## 12. Commercial formulations

The commercial formulations based on *Lecanicillium* spp. are available in India and other countries are presented in **Table 5**. Number of manufacturers based on *Lecanicillium* spp. products is more in India however; the production is very low and not available to the farmers/stakeholders/growers on time as compared to synthetics due to dominant in pesticides market and lack of awareness to farmers/growers about biopesticides. In India, the efficacy of *Lecanicillium* spp., based products was less due to high temperature and low humidity as compared to temperate countries, even though in India, these products were used as one of the components in IPM and also used for the management of sucking pests of flowers and vegetables in greenhouse.

Country	Trade Name	Target pest	Country	Source
<i>Lecanicillium</i> spp.				
Honduras, El Salvador, Nicaragua, Jamaica	Verzam	Whiteflies, aphids, thrips, mites	Escuela Agrícola Panamericana Honduras	—
Colombia	Vercani WP	Whiteflies	Colombia	<a href="http://www.ica.gov.co">www.ica.gov.co</a>
Uruguay	Lecafol	Whiteflies	Lage y Cía. S.A., Uruguay	<a href="http://www.lageycia.com">www.lageycia.com</a>
<i>L. muscarium</i>				
Denmark, Finland, Italy, UK, Netherlands, Italy, Turkey, Switzerland, Japan, France, India	Mycotal	Whiteflies, thrips	Koppert Biological Systems, Netherlands	<a href="http://www.koppert.com">www.koppert.com</a>

Country	Trade Name	Target pest	Country	Source
<i>V. lecanii</i>				
India	Bio-Catch	Whitefly, Aphids, Thrips, Mealy bugs	M/s T. Stanes & Company, India	www.tstanes.com
	Multiplex Varsha	aphids, thrips, mealy bug, whitefly, scales mites	Multiplex Biotech Pvt. Ltd., Bengaluru, Karnataka, India	www.multiplexgroup.com
	Verti Guard	---Do--	Lokmangal Bio Tech Maharashtra, India	www.lokmangalbiotech.com
	Sun Bio Verti	---Do--	Sonkul Agro Industries Pvt. Ltd. Maharashtra, India	www.bioorganic.co.in
	Vertisterk	Scales, mealy bugs	Vijaya Agro Industries, Maharashtra, India	www.vijayaagro.com
	Green Basivert	Aphids, thrips, whitefly, mealy bug, scales	Greentech Biotech Laboratory, Tamil Nadu, India	www.agrizona.in
	Vertoco-P	whitefly, mealy bug	Utkarsh Agrochem Pvt. Ltd., Surat, India	www.utkarshagrochem.com

**Table 5.** Commercially available products based on *Lecanicillium* spp. [58, 59].

### 13. Conclusions

*Lecanicillium* spp. is promising biocontrol agent and can be used as one of the components of integrated pest management under green house and field conditions against sucking insect pests. *Lecanicillium* is multiplying on commercially available media (potato dextrose agar and broth etc.) till date but it can be mass multiplied at cheaper rate on solid grain media of sorghum and rice; liquid media of sugar cane molasses. It can be used effectively in conjunction with other natural enemies and compatible pesticides.

### Acknowledgements

Author is grateful to Director, Institute of Himalayan Bioresource Technology (Council of Scientific and Industrial Research, New Delhi), Ministry of Science and Technology, Government of India, Palampur, Himachal Pradesh, India for encouragement and support.

### Conflict of interest

Author declares that no conflict of interest is reported.

IntechOpen

IntechOpen

### **Author details**

Sajjalavarahalli Gangireddy Eswara Reddy  
Entomology Laboratory, CSIR-Institute of Himalayan Bioresource Technology,  
Palampur-176061, Himachal Pradesh, India

\*Address all correspondence to: [ereddy2001@yahoo.com](mailto:ereddy2001@yahoo.com)

### **IntechOpen**

---

© 2020 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

## References

- [1] Butt TM, Jackson CW, Murugan W. Fungi as Biocontrol Agents, Progress, Problems and Potentials. CBBS Publishing Co., UK; 2001.
- [2] Wang L, Huang J, You M, Liu B. Time dose mortality modelling and virulence indices for six strains of *Verticillium lecanii* against sweet potato whitefly, *Bemisia tabaci* (Gennadius). Journal of Applied Entomology. 2004; 128:494-500.
- [3] Gindin G, Barash I, Harari N, Raccach B. Effect of endotoxic compounds isolated from *Verticillium lecanii* on the sweet potato whitefly, *Bemisia tabaci*. Phytoparasitica. 1994; 22:189-196.
- [4] Bharathi HT. Isolation and characterization of entomopathogenic fungi and their effectiveness [thesis]. Dharwad: University of Agricultural Sciences (India); 2005.
- [5] Zare R, Gams W. A revision of *Verticillium* section Prostrata. IV. The genera *Lecanicillium* and *Simplicillium*. Nova Hedwigia. 2001; 73:1-50.
- [6] Zare R, Gams W, Culham A,. A revision of *Verticillium* sect. Prostrata. I. Phylogenetic studies using ITS sequences. Nova Hedwigia. 2000;71:465-480.
- [7] Gams W, Zare R. A revision of *Verticillium* Sect. Prostrata. III. Genetic classification. Nova Hedwigia. 2001; 72:329-337.
- [8] Goettel MS, Koike M, Kim JJ, Aiuchi D, Shinya R, Brodeur J. Potential of *Lecanicillium* spp. for management of insects, nematodes and plant diseases. Journal of Invertebrate Pathology. 2008; 98:256-61.
- [9] Shinde SV, Patel KG, Purohit MS, Pandya JR, Sabalpara AN. *Lecanicillium lecanii* (Zimm.) Zare and Games, an important biocontrol agent for the management of insect pests – a review. Agri Review. 2010; 31:235-252.
- [10] Kanaoka MA, Isogai S, Murakoshi M, Ichione A, Suzuki A, Tamura S. Bassionalide, a new insecticidal cyclodepsipeptide from *Beauveria bassiana* and *Verticillium lecanii*. Agricultural and Biological Chemistry. 1978; 42:629-640.
- [11] Suzuki A, Kanaoka MA, Isogai S, Murakoshi M, Ichione A, Tamura S. Bassionalid. A new insecticidal cyclodepsipeptide from *Beauveria bassiana* and *Verticillium lecanii*. Tetrahedron Letters. 1977;25, 2167-2170.
- [12] Soman AG, Gloer JB, Angawi RF, Wicklow, DT, Dowd PF. Vertilecanins: New Phenopicolinic Acid Analogues from *Verticillium lecanii*. Journal of Natural Products. 2001;64: 189-192.
- [13] Claydon N, Grove JF. Insecticidal secondary metabolic products from the entomogenous fungus *Verticillium lecanii*. Journal of Invertebrate Pathology. 1982; 40, 413-418.
- [14] Gindin G, Geschtovt NU, Raccach B, Barash I. Pathogenicity of *Verticillium lecanii* to different developmental stages of the silver leaf whitefly, *Bemisia argentifolii*. Phytoparasitica. 2000; 28:229-239.
- [15] Web of Science 2012 and 2020.
- [16] Scorsetti AC, Humber RA, Gregorio CD, Lastra CC. New records of entomopathogenic fungi infecting *Bemisia tabaci* and *Trialeurodes vaporariorum*, pests of horticultural crops, in Argentina. Biological Control. 2008;53:787-796.
- [17] Vu VH, Hong SI, Kim K. Selection of entomopathogenic fungi for aphid control. Journal of Bioscience and Bioengineering. 2007; 104:498-505.

- [18] Hesketh H, Aldersonb PG, Pyec BJ, Pella JK. The development and multiple uses of a standardized bioassay method to select hypocrealean fungi for biological control of aphids. *Biological Control*. 2008; 46:242-255.
- [19] Diaz BM, Oggerin M, Claudia, Lastra CCL, Rubio V, Fereres A. Characterization and virulence of *Lecanicillium lecanii* against different aphid species. *BioControl*. 2009; 54:825-835.
- [20] Ganassi S, Grazioso P, Moretti A, Sabatini MA. Effects of the fungus *Lecanicillium lecanii* on survival and reproduction of the aphid, *Schizaphis graminum*. *BioControl*. 2010; 55:299-312.
- [21] Subramaniam SR, Babu A, Pradeepa N, A new report of the Entomopathogen, *Lecanicillium lecanii* infecting larvae of tea thrips, *Scirtothrips bispinosus* (Bagnall). *Journal of Biosciences Research*. 2010; 1:146-148.
- [22] Chiriví-Salomón JS, Dannies G, RestrepoS, Sanjuan T. *Lecanicillium sabanense* sp. nov. (Cordycipitaceae) a new fungal entomopathogen of coccids. *Phytotaxa*. 2015; 234: 63-74.
- [23] Lu L, Cheng B, Du D, Hu X, Peng A, Pu Z, Zhang X, Huang Z, Chen Guoqing. Morphological, molecular and virulence characterization of three *Lecanicillium* species infecting Asian citrus psyllids in Huangyan citrus groves. *Journal of Invertebrate Pathology*. 2015; 125:45-55.
- [24] Corallo B, Simeto S, Martínez G, Gómez D, Abreo E, Altier N, Lupo S. Entomopathogenic fungi naturally infecting the eucalypt bronze bug, *Thaumastocoris peregrinus* (Heteroptera: Thaumastocoridae), in Uruguay. *Journal of Applied Entomology*. 2019; DOI: 10.1111/jen.12624.
- [25] Kope HH, Alfaro RI, Lavalley R. Effects of temperature and water activity on *Lecanicillium* spp. conidia germination and growth, and mycosis of *Pissodes strobe*. *Biological Control*. 2008;53:489-500.
- [26] Aiuchi D, Baba Y, Inami K, Shinya R, Tani M, Koike M. Variation in growth at different temperatures and production and size of conidia in hybrid strains of *Verticillium lecanii* (*Lecanicillium* spp.) (Deuteromycotina: Hyphomycetes). *Applied Entomology and Zoology*. 2008;43, 427-436.
- [27] Jaronski ST. Ecological factors in the inundative use of fungal entomopathogens. *BioControl*. 2010; 55:159-185.
- [28] Gao L, Liu XZ, Sun MH, Li SD, Wang JL. Use of a novel two-stage cultivation method to determine the effects of environmental factors on the growth and sporulation of several biocontrol fungi. *Mycoscience*. 2009; 50:317-321.
- [29] Rivas F, Nunez P, Jackson T, Altier N. Effect of temperature and water activity on mycelia radial growth, conidial production and germination of *Lecanicillium* spp. isolates and their virulence against *Trialeurodes vaporariorum* on tomato plants. *BioControl*. 2014; 59:99-109.
- [30] Mohammed AA, Hatcher PE. Combining entomopathogenic fungi and parasitoids to control the green peach aphid *Myzus persicae*. *Biological Control*. 2017; 110:44-55.
- [31] Ekbohm BS. Humidity requirements and storage of the entomopathogenic fungus (*Verticillium lecanii*) for use in greenhouses. *Annales Entomologicae Fennicae*.1981;47:61-62.
- [32] Milner RJ, Lutton JJ. Dependence of *Verticillium lecanii* (Fungi: Hyphomycetes) on high humidities for infection and sporulation using *Myzus persicae* (Homoptera: Aphididae) as host. *Environmental Entomology*. 1986; 15:380-382.

- [33] Kim JJ, Goettel M, Gillespie DR. Evaluation of *Lecanicillium longisporum*, (Vertalec) for simultaneous suppression of cotton aphid, *Aphis gossypii*, and cucumber powdery mildew, *Sphaerotheca fuliginea* on potted cucumbers. *Biological Control*. 2008; 45:404-409.
- [34] Ghaffari S, Karimi J, Kamali S, Mehdikhani E. Biocontrol of *Planococcus citri* (Hemiptera: Pseudococcidae) by *Lecanicillium longisporum* and *Lecanicillium lecanii* under laboratory and greenhouse conditions. *Journal of Asia-Pacific Entomology*. 2017; 20(2):605-612.
- [35] Aiuchi D, Moyo D, Ishikura S, Tani M, Kinoshita R, Rakotondrafara AM, Koike M. Virulence of *Lecanicillium* spp. (Hypocreales: Cordycipitaceae) hybrid strains against various biological stages of the greenhouse whitefly, *Trialeurodes vaporariorum* (Hemiptera: Aleyroidae). *Biocontrol Science and Technology*. 2020; DOI: 10.1080/09583157.2020.1771281.
- [36] Cuthbertson AGS, Walters KFA. Pathogenicity of the entomopathogenic fungus, *Lecanicillium muscarium*, against the sweet potato whitefly, *Bemisia tabaci* under laboratory and glasshouse conditions. *Mycopathologia*. 2005; 160:315-319.
- [37] Yeo H, Pell JK, Alderson PG, Clark SJ, Pye BJ. Laboratory evaluation of temperature effects on the germination and growth of entomopathogenic fungi and on their pathogenicity to two aphid species. *Pest Management Science*. 2003; 59:156-165.
- [38] Kim JJ, Goettel MS, Gillespie DR. Potential of *Lecanicillium* species for dual microbial control of aphids and the cucumber powdery mildew fungus, *Sphaerotheca fuliginea*. *Biological Control*. 2007; 40:327-332.
- [39] Down R, Cuthbertson AGS, Mathers JJ, Walters KFA. Dissemination of the entomopathogenic fungi, *Lecanicillium longisporum* and *L. muscarium*, by the predatory bug, *Orius laevigatus*, to provide concurrent control of *Myzus persicae*, *Frankliniella occidentalis* and *Bemisia tabaci*. *Biological Control*. 2009; 50:172-178.
- [40] Saranya S, Ushakumari R, Jacob S, Philip BM. Efficacy of different entomopathogenic fungi against cowpea aphid, *Aphis craccivora* (Koch). *Journal of Biopesticides*. 2010; 3:138-142.
- [41] Cuthbertson AGS, Walters KFA, Northing P. The susceptibility of immature stages of *Bemisia tabaci* to the entomopathogenic fungus *Lecanicillium muscarium* on tomato and verbena foliage. *Mycopathologia*. 2005;159,23-29.
- [42] Kim JJ, Goettel MS, Gillespie DR. Evaluation of *Lecanicillium longisporum*, (Vertalec) for simultaneous suppression of cotton aphid, *Aphis gossypii*, and cucumber powdery mildew, *Sphaerotheca fuliginea* in a greenhouse environment. *Crop Protection*. 2010;29: 540-544.
- [43] Lee WW, Shin TY, Bae SM, Woo SD. Screening and evaluation of entomopathogenic fungi against the green peach aphid, *Myzus persicae*, using multiple tools. *Journal of Asia-Pacific Entomology*. 2015; 18:607-615.
- [44] Mohammed AA, Kadhim JH, Kamaluddin ZNA. Selection of highly virulent entomopathogenic fungal isolates to control the greenhouse aphid species in Iraq. *Egyptian Journal of Biological Pest Control*. 2018; 28:71-78.
- [45] Xie T, Jiang L, Li J, Hong B, Wang X, Jia Y. Effects of *Lecanicillium lecanii* strain JMC-01 on the physiology, biochemistry, and mortality of *Bemisia tabaci* Q-biotype nymphs. 2019; PeerJ. 7: e7690 DOI
- [46] Kamp AM, Bidochka MJ. Conidium production by insect pathogenic fungi on commercially available agars. *Letters in Applied Microbiology*. 2002; 35:74-77.
- [47] Sahayaraj K, Namasivayam SKR. Mass production of entomopathogenic

fungi using agricultural products and by products. African Journal of Biotechnology. 2008; 7:1907-1910.

[48] Derakhshan A, Rabindra RJ, Ramanujam B, Rahimi M. Evaluation of different media and methods of cultivation on the production and viability of entomopathogenic fungi, *Verticillium lecanii* (Zimm.) Viegas. Pakistan Journal of Biological Science. 2008; 11:1506-1509.

[49] Vu VH, Hong SI, Kim K. Production of aerial conidia of *Lecanicillium lecanii* 41185 by solid state fermentation for use as a mycoinsecticide. Mycobiology. 2008; 36:183-189.

[50] Kim JJ, Kim KC, Roberts DW. Impact of the entomopathogenic fungus *Verticillium lecanii* on development of an aphid parasitoid, *Aphidius colemani*. Journal of Invertebrate Pathology. 2005; 88:254-256.

[51] Koike M, Kodama T, Kikuchi A, Okabe M, Kuramoti K, Saito Y. Effects of *Verticillium lecanii* (*Lecanicillium* spp.) against two spotted spider mite, *Tetranychus urticae* and its natural enemy *Phytoseiulus persimilis*. In: Abstracts of 38th annual meeting of society of invertebrate pathology, 7-11 August 2005; Anchorage, Alaska, USA.

[52] Smitha MS, Giraddi RS. Safety of pesticidal sprays to natural enemies in chilli (*Capsicum annuum* L.). J. Biological Control. 2006; 20:7-12.

[53] Derakhshan SA, Ramanujam B. Impact of entomopathogenic fungus, *Verticillium lecanii* (Zimmerman) Viegas on natural enemies of cabbage aphid, *Brevicoryne brassicae* (Linnaeus) and other beneficial insects. Proceedings of the National Symposium on Biological Control of Sucking Pests. 26-27 May, 2006; Bangalore, India.

[54] Donka A, Sermann H, Büttner C. Side-effect of the entomopathogenic fungus *Lecanicillium muscarium* on the predatory mite *Phytoseiulus persimilis*.

IOBC/WPRS Bulletin 45;267-269. In: Proceedings of the IOBC/WPRS Working Group "Insect Pathogens and Insect Parasitic Nematodes" and COST Action 862 "Bacterial Toxins for Insect Control, 22-25 June 2009; Pamplona, Spain. Ed. by Ehlers RU, Crickmore N, Enkerli J, Glazer I, Lopez-Ferber M, Tkaczuk C.

[55] Ren SX, Ali S, Huang Z, Wu JH, *Lecanicillium muscarium* as microbial insecticide against whitefly and its interaction with other natural enemies. Current Research, Technology and Education Topics in Applied Microbiology and Microbial Biotechnology. Ed. by Vilas AM, 340-348;2010.

[56] de Silva RJ, Martins ICF, de Alencar JDCC, Silva KP, Cividanés FJ, Duarte RT, Agostini LT, Polanczyk RA. The effect of the entomopathogenic fungus *Lecanicillium longisporum* (Petch) Zare & Gams (Hypocreales: Cordycipitacea) on the aphid parasitoid *Diaeretiella rapae* McIntoch (Hymenoptera: Braconidae, Aphidiinae). Entomologia Generalis. 2017; 36:219-229.

[57] Gurulingappa P, McGee P, Sword GA. In-vitro and in planta compatibility of insecticides and the endophytic entomopathogen, *Lecanicillium lecanii*. Mycopathologia. 2011; 172:161-168.

[58] Todd KJ, Svircev AM, Goettel MS, Woo SG. The use and regulation of microbial pesticides in representative jurisdictions worldwide [Internet]. 2010. IOBC Global, 99. [http:// www.iobc-lobal.org](http://www.iobc-lobal.org).

[59] Wraight SP, Lopes RB, Faria M. Microbial control of mite and insect pests of greenhouse crops. 2017; 237-252. In: Lawrence A. Lacey, editor. Microbial Control of Insect and Mite Pests From Theory to Practice. Elsevier; 2017. P. 237-252. Academic Press. DOI: 10.1016/C2015-0-00092-2.