

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,200

Open access books available

129,000

International authors and editors

150M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# Plant Uptake, Translocation and Metabolism of PBDEs in Plants

*Daniel Dobslaw, Bertram Kuch, Christine Woiski  
and Martina Kiel*

## Abstract

Polybrominated diphenyl ethers (PBDEs) have been widely used as flame retardants in concentrations up to 30 w% of the total mass of the products. Worldwide consumption of technically relevant PBDE mixtures was about 7500 tons (penta-BDEs), 3790 tons (octa-BDEs) and 56,100 tons (deca-BDE) in 2001 and about 50–60% of this total volume was discharged into environment only by agricultural use of sewage sludges. The use of PBDEs was strictly regulated from 2004 onwards due to their high emission load and their effect as endocrine disrupters, neurotoxins, and fertility reducing agents. Nevertheless, soils worldwide are contaminated by gaseous and particle-bound transport of PBDEs. Therefore, the uptake of PBDEs from contaminated agricultural land via crops and the food chain is a major human exposure pathway. However, uptake and intrinsic transport behavior strongly depend on crop specifics and various soil parameters. The relevant exposure and transformation pathways, transport-relevant soil and plant characteristics and both root concentration factor (RCF) and transfer factor (TF) as derivable parameters are addressed and quantified in this chapter. Finally, based on available crop specific data a general statement about the transport behavior of PBDEs in twelve different crops according to relevant PBDE congeners is given.

**Keywords:** Plant uptake, translocation, root concentration factor, PBDE, shoot concentration factor, food industry, crops

## 1. Introduction

Polybrominated diphenyl ethers (PBDEs) have been used for decades as flame retardants in a wide variety of products. Notable among these are building insulations, upholstered furniture, electrical goods, vehicles and aircrafts, foams, textiles, electrical insulations, and a variety of technical plastics such as acrylonitrile-butadiene-styrene copolymers (ABS), high impact polystyrene (HIPS), polybutylene terephthalate (PBT), or plain paper (PAP), where PBDEs were used in concentrations of 5–30 percent by weight. Despite a spectrum of 209 PBDE congeners, only three formulations were of technical relevance, namely pentabromodiphenyl ether (penta-BDEs), octabromodiphenyl ether (octa-BDEs), and perbrominated diphenyl ether (deca-BDE, BDE-209). The global demand (EU demand) of these mixtures in 2001 was about 7500 tons (EU:150 tons), 3790 tons (EU: 610 tons), and 56,100 tons (EU: 7600 tons) [1]. The use of BDE-209 reached its peak in 2003–2006 with 30,000 tons (China), 9600 tons (EU), 5000–10,000 tons (Northern

America), and 1600 tons (Japan) [2]. Because of their endocrine-disrupting properties, neurotoxicity, and negative impacts on fertility as well as their high environmental persistence, the use of these mixtures was strictly regulated by the Stockholm Convention of 2001 and finally banned in 2004 (penta-BDEs, octa-BDEs) and severely restricted for BDE-209 in 2019 according to the lower toxicity in case of higher degree of bromination [3, 4].

As former high-volume chemicals PBDEs are ubiquitous in environment today, but are mainly detectable in soil and dust samples. E-waste sites and waste water sludge were identified as main sources with BDE-209 as dominant congener [5, 6]. Hence, BDE-209 levels of 6.3–12,194.6 ng g DM<sup>-1</sup> (dry matter) at a ratio of 35–89.6% of the total PBDE where detected on e-waste recycling sites [7–9], while currently highest  $\sum$ PBDE levels of 8.70–18,451 ng g DM<sup>-1</sup> were reported for soil at an industrial production site of plastic parts in electrical industry in Changzhou [10]. PBDE levels of 7240–10,469 ng g DM<sup>-1</sup>, 180–370,000 ng g DM<sup>-1</sup>, and 270–110,000 ng g DM<sup>-1</sup> were also reported as currently highest levels in dust of industrial environment, house dust, and office dust samples in the UK, respectively [11]. In sludge samples of both municipal and industrial waste water treatment plants (WWTPs) in the USA, Turkey, and Hessian (Germany) levels of 85.5 ng g DM<sup>-1</sup> up to 2.5 w% of  $\sum$ PBDE were reported [12–14]. The annual input of PBDEs into the environment in the USA was quantified as 47.9–60.1 tons, where 24.0–36.0 tons were located in sewage sludge [6].

As WWTP sludge is commonly used as fertilizer in agriculture, PBDE contamination is not restricted to hotspots like e-waste sites, and ubiquitous spread is further increased by gaseous and particulate-based transport of PBDEs. Consequently, soil samples were positively tested towards PBDE contaminations in grassland and forest soils of UK and Norway ( $\sum$ PBDE: 65–12,000 pg. g DM<sup>-1</sup>) [15], Western Austria ( $\sum$ PBDE: 10.4–2744 pg. g DM<sup>-1</sup>) [16], Germany (BDE-47: <27–505 pg. g DM<sup>-1</sup>, BDE-209: <156–461 pg. g DM<sup>-1</sup>) [17], and the Arctic ( $\sum$ 12PBDEs ex BDE-209: 120 pg. g DM<sup>-1</sup>, and  $\sum$ PBDE: 1.7–416 pg. g DM<sup>-1</sup>) [17, 18].

Given the widespread distribution of PBDEs in soils, it can be assumed that they are absorbed by plants to a significant extent and are then introduced into humans via the food chain. The following sections are intended to show the factors influencing the plant uptake of PBDEs and their degradation products and to examine in detail the uptake and transport behavior of twelve selected crops. From these data, generally valid relationships for other crops will be derived using simple key parameters.

## 2. Physical characteristics and their effect on transport pathways and plant uptake

Because PBDEs show a wide range of molecular weight (328–959 g mol<sup>-1</sup>), lipophilicity (log K<sub>OW</sub> = 6–10), and volatility (log K<sub>OA</sub> = 9–16) [19, 20], BDE congener specific transport and plant uptake mechanisms (soil-air-plant vs. soil-soil moisture-root-plant) are highly different and dependent on specific substance parameters (vapor pressure, Henry coefficient, air-plant partition coefficient, K<sub>OW</sub> value, K<sub>OA</sub> value), meteorological parameters (temperature, wind rate, precipitation, temporal rainfall distribution, deposition kinetics of gaseous and particulate BDEs), long range transport, plant specific characteristics (species, lipid content, carbohydrate content, fiber content, foliage morphology, non-lipid plant parts, rind consistency), and rhizosphere factors [18, 21–23]. Under aspects of transport, low brominated BDEs (Br<sub>2</sub>-Br<sub>3</sub>) are mainly and medium brominated BDEs (Br<sub>4</sub>-Br<sub>5</sub>), depending on the study, are minorly to dominantly distributed as gaseous

compounds (BDE-15: 100%; BDE-28: 35–60%), while transmission and deposition of higher brominated congeners (Br<sub>6</sub>–Br<sub>10</sub>) are obligatorily characterized by adsorption of BDEs on a particulate phase [17, 22–25]. As a result of the particle-bound lower-range transport of the latter PBDEs, the PBDE pattern in soil and plant samples from densely populated regions and near hotspots shows high agreement with the present PBDE emission spectrum, while in sparsely populated regions the PBDE spectrum is dominated by low brominated congeners [7, 23]. Additionally, a significant concentration gradient of  $\sum$ PBDEs from both densely populated to sparsely populated regions and from emission sites to adjacent region can be observed [7, 26]. Hence, various studies around the world showed a wide range of PBDE concentrations in dust samples of 4.33–370,000 ng g DM<sup>-1</sup> [11, 20, 27–31] with a clear domination of BDE-209 in the range of 69.2–99.6% [11, 20, 25].

Due to the high molar mass and the lipophilicity of high brominated BDEs, plant uptake by the soil–soil moisture–root–plant pathway is of low relevance and restricted to low and medium brominated BDEs (Br<sub>2</sub>–Br<sub>5</sub>) like BDE-47, BDE-99 and BDE-100 [21, 32], even though intrinsic transport of BDE-209 was reported by single studies [33–35], but disproved by Wu et al. [36], where plant availability of BDE-209 was quantified as 0.3–0.5% of the initial soil concentration and 99.5–99.7% of BDE-209 are solely adsorbed on the soil matrix and the outer side of the roots. Hence, atmospheric uptake of high brominated BDEs is the dominant pathway, even though BDE-209 reveals a low ratio of 0.1% of the atmospheric PBDE pattern [24, 37].

### 3. Human exposure to PBDE contaminations and uptake

As a result of the presence of gaseous and particulate PBDEs in air, the human PBDE uptake is dominated by inhalation, but the relevance of this pathway is strongly affected by atmospheric PBDE levels. Hites and Sjödin et al. reported concentrations of 5.27–301 pg. m<sup>-3</sup> in ambient air and 0.06–67 ng m<sup>-3</sup> at indoor air, but increased levels up to 312.1 ng BDE-209 m<sup>-3</sup> at a Swedish e-waste recycling site [38, 39]. Average BDE-209 levels of 0.13 ng m<sup>-3</sup> (gaseous) and 140 ng m<sup>-3</sup> (particulate) were reported by Li et al. in 14 Chinese air samples and total BDE-209 uptake by inhalation was quantified as 3000 ng d<sup>-1</sup> (respiration) and 69 ng d<sup>-1</sup> (dust uptake), equivalent to 84% of the total daily uptake [40]. This finding was validated by multiple studies. As a consequence, 16% of daily PBDE uptake, and even higher ratios in case of lower ambient PBDE levels, are assigned to dietary uptake underlining the relevance as second dominant pathway.

### 4. Transformation and detoxification of PBDEs in plants

While PBDEs in the atmosphere are photolytically transformed by hydroxylation and subsequent transformation to lower brominated congeners or ring closure to the corresponding dibenzofurans [41, 42] and PBDEs in soil and sediments are mainly mineralized by stepwise debromination or detoxified by hydroxylation (OH-BDE) or methoxylation reactions (MeO-BDE) in the rhizosphere, strongly affected by the degree of bromination, concentration of oxygen, organic matter and microorganisms [43], intrinsic PBDEs in plants can be transformed by the same three transformation pathways. Exemplarily, transformation of BDE-28 and BDE-47 in maize was analyzed in detail by Wang et al. [44]. BDE-47 (Br<sub>4</sub>) was dominantly converted to 6-MeO-BDE-47 (275 ng•g<sup>-1</sup> DM) in the root phase, followed by 5-MeO-BDE-47 (40 ng•g<sup>-1</sup> DM),  $\sum$ Br<sub>2</sub>-BDEs (23 ng•g<sup>-1</sup> DM),  $\sum$ Br<sub>3</sub>-BDEs (20 ng•g<sup>-1</sup>

DM), and small quantities of two unidentified hydroxylated BDEs ( $8 \text{ ng}\cdot\text{g}^{-1}$  DM) with continued decrease over time, similar to the transformation behavior against BDE-28 ( $\text{Br}_3$ ). Similar results were observed in plants of pumpkins, rice, wheat and soybean for BDE-47 and BDE-99 with formation of 5-OH-BDE-47, 6-OH-BDE-47, 4'-OH-BDE-49, 4'-OH-BDE-42, 4-MeO-BDE-42, and BDE-28 in case of BDE-47 as parent congener, and with formation of 4-OH-BDE-99 and 4-MeO-BDE-99 in case of BDE-99 as soil contaminant [45, 46]. The total PBDE levels clearly dropped in all studies. In liver cells, transformation of both low and moderate brominated BDEs was shown by cytochrome P450 monooxygenases and glutathione-S-transferase, two enzyme complexes also found in plants, where they can potentially catalyze the same reactions. However, both enzyme sets were induced by BDE-209, but this congener was not converted [47]. In difference, various study showed comparable PBDE patterns both in soil and plant tissues at almost unchanged concentration levels over time, underlining negligible metabolism of PBDEs in plants [6]. In summary, PBDEs might be transformed in plants by debromination, hydroxylation and methoxylation reactions, but transformation behavior strongly depends on the plant species and the established microbial consortium in the rhizosphere [48].

## 5. Soil-root transport: RCF and TF value

### 5.1 RCF value of PBDEs

As a result of high bromination, high molecular weight, low mobility and high lipophilicity, BDE-209 as the dominant PBDE in soil is only marginally available for plants at levels of 0.3–0.5% of the initial concentration [36]. Despite the apparently low relevance, the soil – soil moisture – root uptake pathway is still of high relevance as tests with living and non-living roots of different plants showed 3.5–6 times higher BDE-209 levels in the living tissues [49]. Additional analysis of small-scale soil-based BDE gradients within the root plexus underlined the assumption of active BDE-209 uptake by plants [9] and was clearly proved by greenhouse experiments cultivating six different plant species in contaminated and non-contaminated soil in parallel [50]. In comparison to BDE-209 levels in contaminated soil, BDE-209 levels reached  $5.2\text{--}10.4 \text{ ng g DM}^{-1}$  and, therefore, less than 5%, i.e. more than 95% of BDE-209 contamination in plants could be attributed to root uptake and intrinsic plant transport. Both processes are coupled to plant transpiration elevating PBDE levels in shoots and leaves at dry weather conditions [51].

To increase comparability of PBDE uptake and intrinsic transport, both the root concentration factor (RCF) and the translocation factor (TF) were introduced in literature and correlated to the  $\log K_{\text{OW}}$  value of PBDEs [9]. Both parameters can be correlated in a clearly negative way, i.e. higher RCF values were detected in case of lower brominated PBDEs and, therefore, compounds with lower  $\log K_{\text{OW}}$  values than in case of higher  $\log K_{\text{OW}}$  values [9]. In detail, plant specific RCFs of BDE-209 were up to ten times lower than the RCFs of BDE-28 [7, 9, 19]. This observation may be explained both by the lack of water solubility and thereby restricted root uptake with the soil moisture phase, and the strong adsorptive behavior of higher brominated PBDEs in soil. Hence, with exception of some plants like radish, green squash, and soft-stem bulrush, which are well adapted for phytoremediation processes, RCFs are clearly less than 1 for all PBDE congeners [52]. However, PBDE uptake is significantly affected by the presence of organic solubilizers in the soil like extracts of wheat straw or pig manure, where BDE-47 uptake by wheat as an example was elevated by a factor of 3.1 (wheat manure) and 1.9 (pig manure), respectively [53]. Hence, PBDE uptake increases with increasing surfactant activity, whereas decreases

at weak surfactant activity and thereby increase of the organic content in soil [54]. Furthermore, PBDE uptake and RCFs as consequence are strongly affected by plant species specifics, physical and chemical soil properties, initial concentration levels of PBDEs, relevance of both gaseous and particulate atmospheric uptake, duration of the growth period, and organic soil content. Details will be described in Section 6.

In contrast, a positive correlation of both RCF and TF value was observed for maize in case of BDE-15, BDE-28, and BDE-47, which can be explained by an increased transpiration of the plants as these low-brominated congeners reveal higher water solubility [51].

## 5.2 TF value of PBDEs

The ratio of PBDE levels in shoots to the levels in roots is defined as translocation factor (TF). In contrast to the RCF, a general statement about the correlation of  $\log K_{OW}$  and TF is not appropriate, since no clear positive or negative correlation occurs as the TF value depends on numerous parameters like plant species specifics, initial PBDE levels in soil, the lipid content of the shoots, plant age, distance of the plant issue from the root plexus, and the hardly quantifiable effect of the soil-air-plant exposure pathway. In principle, a negative correlation may be assumed as lower  $\log K_{OW}$  values correlate with higher water solubility and, hence, higher intrinsic transport. As PBDE accumulate in the root area, stem and shoots show significantly lower contamination levels and relevance of atmospheric PBDE uptake significantly increases [51]. Nevertheless, the bioaccumulation and translocation behavior of PBDE in plants is not conclusively clarified and depends on numerous, partially insufficiently determined parameters.

## 6. Factors of PBDE plant uptake

Numerous studies focused on both physico-chemical and substance specific properties affecting plant uptake and biodegradation behavior of PBDE, where PBDE specifics (vapor pressure,  $K_{OW}$  value, air-water distribution  $K_{AW}$  value, air-plant distribution  $K_{AP}$  value), environmental parameters (temperature, wind rate, precipitation, temporal rain distribution, kinetics of both gaseous and particle-bound deposition), plant properties (species, lipid content, foliage morphology, ratio of non-lipid plant parts, rind thickness, contents of both sugar and fibers), as well as the presence of an microbial active rhizosphere were generally found to be highly germane. For bioavailability and thus biodegradability of PBDEs pH value and soil composition are of particular importance [18, 22, 23]. In detail, relevant parameters are:

### 6.1 Excretion of plant solubilizers

Easy metabolizable intermediates as amino acids, organic acids, sugars and exoenzymes are excreted by plants as detoxification strategy to improve microbial bioavailability and biodegradability of PBDE in the rhizosphere [33]. For example, hexose was excreted by *Kandelia obovate* to enhance microbial debromination of BDE-99 to  $Br_2$ -BDEs and  $Br_3$ -BDEs in soil [55].

### 6.2 Plant specifics

Plant specifics like plant morphology, wax layers of bay leaves and the lipid content of both leaf and roots strongly affect both atmospheric and soil-based uptake of PBDEs. Hence, accumulation of  $Br_3$ -BDEs to  $Br_{10}$ -BDEs in the wax layer

of wheat was determined as 29–93% of the total plant uptake [25], while BDE-209 accumulation in six different plant species at initial soil levels of  $4700 \text{ ng g DM}^{-1}$  ranged  $1822\text{--}10,933 \text{ ng plant}^{-1}$  with alfalfa showing lowest and maize showing highest levels [56].

### 6.3 Rhizosphere and mycorrhiza

The release of plant eluates into the mycorrhizal is part of the symbiosis between the plant and the mycorrhizal fungi promoting the plant's uptake of nutrients and the growth of microorganisms in the mycorrhizal area. Its secondary effect of enhanced microbial biodegradation and detoxification of PBDEs in the mycorrhizal area was shown several times in literature. As an example, increased depletion of  $470 \text{ ng g DM}^{-1}$  towards  $2250 \text{ ng g DM}^{-1}$  was observed for BDE-209 comparing rice plants with and without mycorrhizal fungi [57].

### 6.4 Specific root and leaf surface

A correlation between accumulation of PBDEs and high specific plant surface was clearly shown for BDE-209 in roots of radish, lettuce and taro [58, 59] and for  $\text{Br}_2\text{-BDE}$  to  $\text{Br}_{10}\text{-BDE}$  in both pine needles and eucalyptus leaves [60].

### 6.5 Lipid content

The lipid content of a plant, especially of the roots, shows strong effects on PBDE uptake and was successfully evaluated for various mosses, lichens, ryegrass, alfalfa, maize, radish, squash, and pumpkin. A positive correlation between lipid content and PBDE uptake (RCF), as well as a negative correlation in lipid content and intrinsic PBDE mobility (TF) was observed [19, 22, 61, 62].

### 6.6 Organic content of the soil

Similar to the lipid content an increase in organic content of soil evokes higher PBDE accumulation in the soil and, therefore, reduced PBDE plant uptake [20, 33, 63]. Exemplarily, PBDE uptake in carrots was reduced by 31.5–69.8% and soil-based biodegradation increased by 8.6–28.5% by addition of 1–4 w% of swine manure to the soil fraction [63]. Differentiation between TOC and DOC showed a clear improvement in adsorption of PBDE in the soil matrix at higher TOC levels, whereas no effect was observed in case of increased DOC levels [54].

### 6.7 Biochar

Since biochar is insoluble and thus is considered as a TOC increase of the soil, elevated accumulation of PBDE in the soil phase may occur [64].

### 6.8 Sewage sludge

As sewage sludge reveals a high TOC content, but also enhanced contamination levels with PBDE or their detoxification and degradation products, sewage sludge is a dominant PBDE exposure pathway. Hence, using contaminated sludges as agricultural fertilizers, increasing concentration levels and accumulation of PBDE in the soil phase and, finally, elevated PBDE plant uptake are observed. However, relevance of this effect strongly depends on the original PBDE contamination levels of the sludge. An overview of PBDE levels in different sewage sludges was

previously presented [52]. Similar to both lipid content and TOC content, a negative correlation between BDE-209 uptake and organic content, implemented by sewage sludge out, was observed [37].

## 6.9 Compost and digestate


In difference to sewage sludge, the PBDE load of compost and digestates is rather low as confirmed by various studies due to the low contamination levels of the plant educts (leaves, green waste, fruit and food residues). For instance, a broadly based study of biocompost, green waste compost and digestates in Baden-Wuerttemberg showed comparable median concentrations of  $13 \text{ ng g DM}^{-1}$ ,  $5.4 \text{ ng g DM}^{-1}$ , and  $13.7 \text{ ng g DM}^{-1}$  and confirmed to low relevance of these materials as PBDE emissions sources [65].

## 6.10 Soil humidity

As a result of the low water solubility of PBDE, high soil moisture effectively prevents evaporation of BDEs as well as plant uptake [64]. Correspondingly, a longer PBDE load may be expected at wet locations.

## 6.11 Plastic particles

Plastics may reveal partition coefficients of PBDEs of several orders of magnitude higher than those towards sewage sludge or soil at levels of 0.04–1.6 w% [66, 67]. In detail, accumulation of PBDEs in low density polyethylene films (LDPE), an adequate and commonly accepted model for diffusive transport phenomena across biomembranes, showed accumulation factors against the aqueous phase ( $K_{PEW}$ ) of  $10^5$ – $10^7$  in case of 23 BDE congeners covering the total spectrum of bromination [68]. Furthermore, a curvilinear correlation between the  $\log K_{PEW}$  and  $\log K_{OW}$  factor, very similar to the correlation of BCF and  $\log K_{OW}$  shown in **Figure 1** was observed and explained by energy barriers for diffusive transport into the LDPE structure [68]. The statement of rising  $K_{PEW}$  levels at higher lipophilicity of BDE congeners was further confirmed [67]. However,  $K_{PEW}$  levels are strongly affected



**Figure 1.** Correlation of  $\log K_{OW}$  and  $\log BCF$  of 25 BDEs of high environmental relevance (-3, -7, -17, -28, -30, -47, -49, -66, -85, -99, -100, -123, -153, -154, -155, -183, -184, -191, -197, -201, -202, -206, -207, -208, -209) applying simple mathematical models with/without correction.

by the type of plastic and the temperature level during test conditions as adsorption is an endothermic process. While enhanced accumulation of BDE congeners was observed for polyethylene, polypropylene, polystyrene, and polyamide, low accumulation levels were observed for polyvinylchloride [66–68]. Hence, the hypothetical potential of injection of plastic particles into the soil as sink for PBDE and thereby soil remediation was positively investigated.

### 6.12 Other additives

Additional additives like graphene,  $\text{TiO}_2$ ,  $\text{Al}_2\text{O}_3$ , Ag, and carbon nanotubes were considered as relevant for BDE-209 uptake in spinach, pumpkin, cucumber, corn and water spinach [36]. Indeed, an increased plant uptake was observed for all of these additives. Combinations of clay minerals like bentonite and oxidizing agents like sodium persulfate have also been positively tested for increased bioavailability of  $\text{Br}_3$ - $\text{Br}_{10}$  – BDEs [69].

### 6.13 Solubilizers

The addition of surfactant-active additives elevates both mobility of PBDE in the soil matrix and plant uptake as previously described.

### 6.14 Macro- and trace elements

Apart from undisputed relevance of macro- and trace elements both in the development of the microflora in the rhizosphere and in plant growth, positive effects on the degradation of BDEs were observed in individual cases. Exemplarily, nitrate as additive caused intensified desorption and biodegradation of BDE-99, but might cause inhibition in case of high levels [70, 71].

### 6.15 Heavy metals

Presence of elevated concentrations of heavy metals shows ambivalent effects. Metals like Ni or Fe cause an enhanced uptake of PBDE, which was justified by chemical debromination of BDE-209 and enhanced mobilization, uptake and transport of  $\text{Br}_8$ - to  $\text{Br}_{10}$ -BDEs in the roots and shoots of the plants [64]. Enhanced plant uptake of low brominated BDEs like BDE-47 was also observed (24.76% vs. < 1.5%) [72]. In contrast, reduced BDE-209 uptake of up to 50% by pumpkins was reported after addition of  $300 \text{ mg Cu kg DM}^{-1}$  and microbial mineralization was negatively affected at more elevated levels [70, 73]. Similar effects were shown for lead, where BDE-209 uptake was reduced by a factor of 2.9–3.7 by tall fescue at levels up to  $1950 \text{ mg Pb kg DM}^{-1}$  [74]. Heavy metal induced effects on PBDE plant uptake are also plant specific as shown for cadmium, where levels of up to  $14,800 \text{ ng g DM}^{-1}$  had no effect on BDE-209 uptake in case of black nightshade, but lifting effects in case of amaranth [75, 76].

In summary, presence of essential heavy metals like iron or copper at adequate concentrations might have a positive effect on PBDE degradation, while non-essential heavy metals at non-toxic levels reveal no effect.

## 7. Predictive mathematical models

Due to the broad spectrum of food plants, attempts were made to develop simple but sensitive models to predict PBDE plant uptake based on simple chemical

conditions and input variables as distribution equilibria, lipid content, organic matter, and initial PBDE soil-water concentration to achieve predictive statements about RCF, SCF (shoot concentration factor) or TF.

While these models provide comparatively good correlations for the RCF, they commonly fail in prediction of the TF, because this value is strongly influenced by intrinsic and atmospheric transport of BDEs in addition to the plant specific uptake of PBDE in the root plexus. Therefore, deviations in the range of two decades can be observed comparing model and real situation [77, 78]. Even after restriction of models to single pollutant situations instead of congener mixtures with variable concentration levels, and after focusing on single and simplified plants like lettuce, where differentiations between shoot and fruit or over the height of the shoot are not applicable, deviations of 25.3–58.2% of the model compared to real situation were reported for the insecticide chlorpyrifor [79].

Another highly relevant error is caused by incorrect analysis of intrinsic PBDE levels in roots in contrast to adsorptive fractions at the outer root surface affecting quality of environmental data. Hence, Briggs et al. [80] showed a significant decrease in BCF levels and thus RCF values of PBDE starting at a log  $K_{OW}$  value of approx. 6.5 (corresponds to a log BCF value of approx. 4.6) after elimination of externally adsorbed congeners (see **Figure 1**). This chart corresponds to Bintein's bilinear model [81], which was confirmed by Meylan et al. [82] for 610 non-ionic pollutants. This negative correlation at high log  $K_{OW}$  values and thus high lipophilicity bases on three restrictions of lipophilic compounds as follows:


- **Equilibrium kinetics:** The higher the lipophilicity of a pollutant, the longer it takes to achieve equilibrium state between two phases or compartments. The life span of annual crops might be too short to establish equilibria between soil and root or root and shoot [83].
- **Solubility:** Water solubility decreases with increasing lipophilicity and strongly lipophilic substances are primarily adsorb onto particles or surfaces. However, for absorptive root uptake of pollutants, a phase transition from soil to the liquid phase as well as from the liquid phase to the intrinsic root is required without adsorptive elimination at the tissue [80, 83]
- **Membrane permeability and cellular transport mechanisms:** The membrane-based cellular uptake of pollutants takes place by passive permeation [80]. The membrane permeability and thus bioavailability of contaminants is concisely described by Lipinski's 'Law of 5', stating out low absorption or membrane permeability at log  $K_{OW}$  values higher than 5, molecular weight higher than 500, more than 5 hydrogen bond donors and more than 10 (= 2 · 5) hydrogen bond acceptors. The former two requirements are fulfilled even in case of Br<sub>4</sub>- to Br<sub>5</sub> – BDEs. By means of known transport mechanisms into the cell, PBDE plant uptake may be affected by co-transport phenomena of biomolecules like amino acids.

## 8. RCF and TF values of specific crops


Following the extensive literature evaluation by Dobslaw et al. [52] twelve crops with the highest documented data density were selected and the occurring RCF and TF values for BDE-47 and BDE-209 were compared. The highest data density regarding the transition of PBDEs from soil to root or from root to plant was available for BDE-209 followed by BDE-47. In contrast to lower brominated

BDEs the concentrations of BDE-209 found in literature range over several decades, which facilitates an evaluation of the literature data concerning transition rates. In particular, the following species were selected: Rice (*Oryza sativa* L.), maize (*Zea mays* L.), prince's-feather (*Amaranthus hypochondriacus* L.), the group of bok choy, Chinese cabbage, broccoli, and filder cabbage (*Brassica* sp.), lettuce (*Lactuca sativa* sp.), spinach (*Spinacia oleracea* L.), sweet potato vine (*Ipomoea batatas* L.), radish (*Raphanus sativus* L.), carrot (*Daucus carota* L.), taro root (*Colocasia esculenta* L. Schott), pea (*Pisum sativum* L.), and pumpkin (*Cucurbita pepo* ssp. *pepo*).


**Figure 2** shows the concentrations of BDE-209 in the roots of these crops as a function of the corresponding soil concentrations (RCF), indicating the dependency of BDE-209 uptake on the soil concentration with a ratio of about 1:10 ( $c_{\text{root}}:c_{\text{soil}}$ ). Comparing shoots and corresponding soil a similar dependency on soil concentration with a ratio of about 1:18 ( $c_{\text{shoot}}:c_{\text{soil}}$ ) was found (**Figure 3**). This representation was chosen because of the higher data density being available. However,



**Figure 2.** BDE-209 – Root concentration factor (RCF), concentration in root as function of corresponding soil concentration.




**Figure 3.** BDE-209 – Concentration in shoot as function of corresponding soil concentration.




**Figure 4.**  
 BDE-209 – Translocation factor (TF), concentration in shoot as function of corresponding root concentration.

a TF could be derived from the comparison of the existing data pairs, which points to the dependence of the shoot concentrations on the concentrations in the root (Figure 4) with a ratio of about 1:9 ( $c_{\text{shoot}}:c_{\text{root}}$ ).


The extent of transition of the tetrabrominated BDE-47 from soil to root and shoot was also determined by the concentration of the flame retardant in the soil (Figures 5 and 6). For BDE-47, despite its lower  $\log K_{OW}$  in comparison to BDE-209, the transition rates are higher with about 1.1:1 ( $c_{\text{root}}:c_{\text{soil}}$ ) and 1:6 ( $c_{\text{shoot}}:c_{\text{soil}}$ ). Too few corresponding data pairs were available for the representation of the TF. A graphical summary of the results for the congeners 47 and 209 is given in Figure 7. The results are consistent with the prediction models described in Section 7 (see also Figure 1). Due to its high  $\log K_{OW}$ , BDE-209 is expected to have a lower uptake compared to BDE-47. With the models mentioned above, it is possible to adequately describe the uptake behavior of lipophilic PBDEs. Of particular interest is the trend that the uptake of the substances is primarily determined by the



**Figure 5.**  
 BDE-47 – Root concentration factor (RCF), concentration in root as function of corresponding soil concentration.



**Figure 6.**  
BDE-47 – Concentration in shoot as function of corresponding soil concentration.



**Figure 7.**  
BDE-47 and BDE-209 – Comparison of concentrations in soil, root and shoot (normalized,  $c(\text{soil}) = 1$ ).

concentration of the compounds in soil. A dependency of the uptake on the crop species might have been expected, but cannot be proven by the present data set.

## 9. Conclusion

PBDEs were widely used as flame retardants. Due to their effects as endocrine disruptors, neurotoxins, and on reproductive capacity as well, knowledge of terrestrial accumulation behavior and, in particular, their uptake by plants for food production is highly relevant. Uptake can occur via soil-air-plant pathway as well as soil-soil water-root-plant pathway. Transport and plant uptake behavior strongly depend on physical and chemical properties of the BDEs, environmental factors, large-scale atmospheric transport processes, plant properties as well as terrestrial rhizospheres. During both atmospheric and terrestrial transport PBDEs are subject to UV-induced (atmospheric) or microbial induced (terrestrial) transformation

and degradation processes like debromination, hydroxylation, methoxylation and ring closure to dibenzofurans. As PBDEs reveal high lipophilicity they tend to adsorption on lipophilic soil matrices and thereby show low uptake via soil–soil water-root-plant pathway and subsequent intrinsic transport. Hence, uptake and intrinsic transport are only expected for low brominated BDEs. Therefore, declining concentrations of PBDEs could be detected from soil via roots to shoots and final fruits, i.e. RCF and TF show negative correlation with their log  $K_{OW}$  values. Consistent with this statement, 84% of the human PBDE intake are attributed to respiration and inhalation of dust, while only 16% were correlated with dietary uptake. The actual exposure of vegetarian foods to PBDEs depends on the following parameters:

- Both microbial degradation and plant uptake of PBDE are elevated by release of easily biodegradable plant extracts like amino acids, organic acids, sugars and exoenzymes as commonly observed in symbiosis with rhizobia.
- The presence of rhizobia enhances microbial degradation and plant uptake of PBDE.
- Both atmospheric and terrestrial PBDE uptakes are strongly affected by the morphology of the plant (specific surface of leaves and roots) and the increasing lipid content of plant tissues accelerating PBDE uptake.
- PBDE immobilization and accumulation in soil is promoted by increasing TOC levels caused by implementation of compost, sewage sludge, digestates, or biochar.
- In contrast, increasing levels of DOC show no effect on plant uptake of PBDEs as long as there is no solubilizing effect by surfactants.
- As sewage sludge shows PBDE contamination up to 2.5 w%, it is an important emission source of PBDEs affecting soil contamination. In contrast, PBDE loads of compost and digestates are very low and commonly pollution effects are negligible.
- A decrease in evaporation losses and an enhanced immobilization tendency of PBDE can be observed in case of raising soil moisture.
- Solubilizers, ionic additives and nanoscale organic substances act as mobilizing agents increasing mobilization and plant uptake of PBDEs.
- Macroelements as nitrate favor terrestrial PBDE degradation through its function as alternative electron acceptor. Hence, plant's load is reduced.
- The presence of trace elements supports microbial transformation of PBDE in soil and therefore causes lower contamination of plants. In contrast, heavy metals seem to enforce PBDE uptake by plants by inhibition of terrestrial biodegradation processes.
- Current mathematical models allow high quality in prediction of the RCF value with a minimum of input parameters. In opposite, the prediction of SCF and TF values is not suitable due to the insufficient coverage of plant-specific parameters.

- According to Lipinski's 'Law of 5' Br<sub>4</sub>- and Br<sub>5</sub>-BDE congeners reveal the highest RCF levels. Higher polarity is required to gain high TF factors. Within the same isomers, even small differences in lipophilicity significantly change these values.
- Available PBDE data only show a linear correlation between soil concentration and plant uptake. Plant specific uptake effects cannot be observed.

## 10. Outlook

The poor biodegradability of PBDEs means that both the accumulation of PBDEs in the food chain and inhalation exposure will continue to be highly relevant issues in the next few decades. Problems comparable to those described for PBDEs are also expected for alternative brominated flame retardants such as hexabromobenzene, pentabromotoluene, 1,2-bis-(2,4,6-tribromophenoxy)ethane, or decabromodiphenylethane due to the likewise high degree of bromination, high persistence and thus a high bioaccumulation potential [20, 84]. A ubiquitous presence of these compounds was already proven [19].

Especially remarkable is the partial lack of metadata in existing publications, which makes an comprehensive evaluation, as it was tried in this publication, difficult. This concerns, for example, more detailed information about the soil composition, the content of organic carbon, lipid, and water of roots and shoots. Thinning of data, such as the aggregation of single congener data to sums (e.g.  $\sum$ PBDE) without further information about the composition makes a valuable data set useless.

## Acknowledgements

We would like to thank Lukas Lesmeister (DVGW, Karlsruhe), Melanie Mechler and Jörn Breuer (Center of Agricultural Technology Augustenberg, Karlsruhe) and Prof. Karl-Heinrich Engesser (ISWA, Stuttgart) for their continuous support. This work was supported by the German Federal Environment Agency at grant number FKZ 3718 74 210 0; AZ: 91 007 – 2/82.

## Conflict of interest

The authors declare that they have no conflict of interest.

## Author details

Daniel Dobslaw\*, Bertram Kuch, Christine Woiski and Martina Kiel  
Institute of Sanitary Engineering, Water Quality and Solid Waste Management,  
University of Stuttgart, Stuttgart, Germany

\*Address all correspondence to: [daniel.dobslaw@iswa.uni-stuttgart.de](mailto:daniel.dobslaw@iswa.uni-stuttgart.de)

## IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

## References

- [1] BSEF Bromine Science and Environmental Forum. Major Brominated Flame Retardants Volume Estimates - Total Market Demand by Region in 2001
- [2] ECHA European Chemicals Agency. Background document to the Opinion on the Annex XV dossier proposing restrictions on Bis(pentabromophenyl) ether: ECHA/RAC/RES-O-0000006155-77-01/D ECHA/SEAC/RES-O-0000006155-77-03/F. Helsinki; 2015
- [3] Sun Y, Wang C, Xu X, Ruan H. Responses of plants to polybrominated diphenyl ethers (PBDEs) induced phytotoxicity: A hierarchical meta-analysis. *Chemosphere*. 2020;**240**:124865. DOI: 10.1016/j.chemosphere.2019.124865
- [4] European Commission EC. EC Regulation 217/227 amending Annex XVII to Regulation (EC) No 1907/2006 of the European Parliament and of the Council concerning the Registration, Evaluation, Authorisation and Restriction of Chemicals (REACH) as regards bis(pentabromophenyl) ether.
- [5] Stone R. China. Confronting a toxic blowback from the electronics trade. *Science* (New York, N.Y.). 2009;**325**:1055. DOI: 10.1126/science.325\_1055
- [6] Venkatesan AK, Halden RU. Brominated flame retardants in U.S. biosolids from the EPA national sewage sludge survey and chemical persistence in outdoor soil mesocosms. *Water research*. 2014;**55**:133-142. DOI: 10.1016/j.watres.2014.02.021
- [7] Han Z-X, Wang N, Zhang H-L, Zhao Y-X. Bioaccumulation of PBDEs and PCBs in a small food chain at electronic waste recycling sites. *Environmental Forensics*. 2017;**18**:44-49. DOI: 10.1080/15275922.2016.1263900
- [8] Leung AOW, Luksemburg WJ, Wong AS, Wong MH. Spatial distribution of polybrominated diphenyl ethers and polychlorinated dibenzo-p-dioxins and dibenzofurans in soil and combusted residue at Guiyu, an electronic waste recycling site in southeast China. *Environmental Science & Technology*. 2007;**41**:2730-2737. DOI: 10.1021/es0625935
- [9] Zhang Y, Luo X-J, Mo L, Wu J-P, Mai B-X, Peng Y-H. Bioaccumulation and translocation of polyhalogenated compounds in rice (*Oryza sativa* L.) planted in paddy soil collected from an electronic waste recycling site, South China. *Chemosphere*. 2015;**137**:25-32. DOI: 10.1016/j.chemosphere.2015.04.029
- [10] Xu J, Qian W, Li J, Zhang X, He J, Kong D. Polybrominated diphenyl ethers (PBDEs) in soil and dust from plastic production and surrounding areas in eastern of China. *Environmental geochemistry and health*. 2019;**41**:2315-2327. DOI: 10.1007/s10653-019-00247-0
- [11] Tao F, Abdallah MA-E, Harrad S. Emerging and Legacy Flame Retardants in UK Indoor Air and Dust: Evidence for Replacement of PBDEs by Emerging Flame Retardants? *Environmental Science & Technology*. 2016;**50**:13052-13061. DOI: 10.1021/acs.est.6b02816
- [12] Hale RC, La Guardia MJ, Harvey E, Chen D, Mainor TM, Luellen DR, Hundal LS. Polybrominated diphenyl ethers in U.S. sewage sludges and biosolids: temporal and geographical trends and uptake by corn following land application. *Environmental Science & Technology*. 2012;**46**:2055-2063. DOI: 10.1021/es203149g

- [13] Leisewitz A, Fengler S, Seel P. Orientierende Messungen gefährlicher Stoffe: Landesweite Untersuchung auf organische Spurenverunreinigungen in hessischen Fließgewässern, Abwässern und Klärschlammern.: Zusammenfassender Abschlussbericht 1991 - 2003. Wiesbaden: Hessisches Landesamt für Umwelt und Geologie (HLUG); 2003
- [14] Demirtepe H, Imamoglu I. Levels of polybrominated diphenyl ethers and hexabromocyclododecane in treatment plant sludge: Implications on sludge management. *Chemosphere*. 2019;**221**:606-615. DOI: 10.1016/j.chemosphere.2019.01.060
- [15] Hassanin A, Breivik K, Meijer SN, Steinnes E, Thomas GO, Jones KC. PBDEs in European Background Soils: Levels and Factors Controlling Their Distribution. *Environmental Science & Technology*. 2004;**38**:738-745. DOI: 10.1021/es035008y
- [16] Freudenschuß A, Obersteiner Erik, Uhl Maria. Organische Schadstoffe in Grünlandböden: TEIL 3 – Endbericht. Wien: Umweltbundesamt GmbH; 2010
- [17] Dreyer A, Neugebauer F, Rüdell H, Klein R, Lohmann N, Rauert C, Koschorreck J. Halogenated flame retardants in tree samples applied as bioindicators for atmospheric pollution. *Chemosphere*. 2018;**208**:233-240. DOI: 10.1016/j.chemosphere.2018.05.033
- [18] Zhu C, Li Y, Wang P, Chen Z, Ren D, Ssebugere P, Zhang Q, Jiang G. Polychlorinated biphenyls (PCBs) and polybrominated biphenyl ethers (PBDEs) in environmental samples from Ny-Ålesund and London Island, Svalbard, the Arctic. *Chemosphere*. 2015;**126**:40-46. DOI: 10.1016/j.chemosphere.2015.01.043
- [19] She Y-Z, Wu J-P, Zhang Y, Peng Y, Mo L, Luo X-J, Mai B-X. Bioaccumulation of polybrominated diphenyl ethers and several alternative halogenated flame retardants in a small herbivorous food chain. *Environmental Pollution*. 2013;**174**:164-170. DOI: 10.1016/j.envpol.2012.11.024
- [20] Zhu H, Sun H, Yao Y, Gan Z, Wang Y, Kannan K. Legacy and alternative brominated flame retardants in outdoor dust and pine needles in mainland China: Spatial trends, dust-plant partitioning and human exposure. *Environmental pollution (Barking, Essex: 1987)*. 2018;**243**:758-765. DOI: 10.1016/j.envpol.2018.08.097
- [21] Klinčić D, Dvorščak M, Jagić K, Mendaš G, Herceg Romanić S. Levels and distribution of polybrominated diphenyl ethers in humans and environmental compartments: a comprehensive review of the last five years of research. *Environmental Science and Pollution Research*. 2020;**27**:5744-5758. DOI: 10.1007/s11356-020-07598-7
- [22] Yogui GT, Sericano JL, Montone RC. Accumulation of semivolatile organic compounds in Antarctic vegetation: a case study of polybrominated diphenyl ethers. *The Science of the total environment*. 2011;**409**:3902-3908. DOI: 10.1016/j.scitotenv.2011.06.010
- [23] Zhao Y-X, Qin X-F, Li Y, Liu P-Y, Tian M, Yan S-S, Qin Z-F, Xu X-B, Yang Y-J. Diffusion of polybrominated diphenyl ether (PBDE) from an e-waste recycling area to the surrounding regions in Southeast China. *Chemosphere*. 2009;**76**:1470-1476. DOI: 10.1016/j.chemosphere.2009.07.023
- [24] Gao M, Wang G, Lin B, Tariq M, Liu K, Zhang W. Study on arbor leaf and ring as a potential biological indicator for atmospheric polybrominated diphenyl ethers (PBDEs) distribution at e-wastes recycling sites. *International Journal of Environmental Science and*

Technology. 2019;**16**:8639-8652. DOI: 10.1007/s13762-019-02428-x

[25] Zhu H, Wang F, Li B, Yao Y, Wang L, Sun H. Accumulation and translocation of polybrominated diphenyl ethers into plant under multiple exposure scenarios. *Environment international*. 2020;**143**:105947. DOI: 10.1016/j.envint.2020.105947

[26] Graziani NS, Tames MF, Mateos AC, Silva JA, Ramos S, Homem V, Ratola N, Carreras H. Estimation of urban POP and emerging SVOC levels employing *Ligustrum lucidum* leaves. *Atmospheric Pollution Research*. 2019;**10**:1524-1530. DOI: 10.1016/j.apr.2019.04.010

[27] Wang J, Ma Y-J, Chen S-J, Tian M, Luo X-J, Mai B-X. Brominated flame retardants in house dust from e-waste recycling and urban areas in South China: implications on human exposure. *Environment international*. 2010;**36**:535-541. DOI: 10.1016/j.envint.2010.04.005

[28] Ma J, Addink R, Yun S, Cheng J, Wang W, Kannan K. Polybrominated dibenzo-p-dioxins/dibenzofurans and polybrominated diphenyl ethers in soil, vegetation, workshop-floor dust, and electronic shredder residue from an electronic waste recycling facility and in soils from a chemical industrial complex in eastern China. *Environmental Science & Technology*. 2009;**43**:7350-7356. DOI: 10.1021/es901713u

[29] Muenhor D, Harrad S, Ali N, Covaci A. Brominated flame retardants (BFRs) in air and dust from electronic waste storage facilities in Thailand. *Environment international*. 2010;**36**:690-698. DOI: 10.1016/j.envint.2010.05.002

[30] Schreder ED, La Guardia MJ. Flame retardant transfers from U.S. households (dust and laundry wastewater) to the aquatic environment. *Environmental*

*Science & Technology*. 2014;**48**:11575-11583. DOI: 10.1021/es502227h

[31] Harrad S, Goosey E, Desborough J, Abdallah MA-E, Roosens L, Covaci A. Dust from U.K. primary school classrooms and daycare centers: the significance of dust as a pathway of exposure of young U.K. children to brominated flame retardants and polychlorinated biphenyls. *Environmental Science & Technology*. 2010;**44**:4198-4202. DOI: 10.1021/es100750s

[32] Mueller KE, Mueller-Spitz SR, Henry HF, Vonderheide AP, Soman RS, Kinkle BK, Shann JR. Fate of Pentabrominated Diphenyl Ethers in Soil: Abiotic Sorption, Plant Uptake, and the Impact of Interspecific Plant Interactions. *Environmental Science & Technology*. 2006;**40**:6662-6667. DOI: 10.1021/es060776l

[33] Zhao L, Jiang J, Chen C, Zhan S, Yang J, Yang S. Efficiency and mechanism of the phytoremediation of decabromodiphenyl ether-contaminated sediments by aquatic macrophyte *Scirpus validus*. *Environmental science and pollution research international*. 2017;**24**:12949-12962. DOI: 10.1007/s11356-017-8900-1

[34] Vrkoslavová J, Demnerová K, Macková M, Zemanová T, Macek T, Hajslová J, Pulkrabová J, Hrádková P, Stiborová H. Absorption and translocation of polybrominated diphenyl ethers (PBDEs) by plants from contaminated sewage sludge. *Chemosphere*. 2010;**81**:381-386. DOI: 10.1016/j.chemosphere.2010.07.010

[35] Chow KL, Man YB, Tam NFY, Liang Y, Wong MH. Uptake and transport mechanisms of decabromodiphenyl ether (BDE-209) by rice (*Oryza sativa*). *Chemosphere*. 2015;**119**:1262-1267. DOI: 10.1016/j.chemosphere.2014.10.016

- [36] Wu X, Wang W, Zhu L. Enhanced organic contaminants accumulation in crops: Mechanisms, interactions with engineered nanomaterials in soil. *Environmental Pollution*. 2018;**240**:51-59. DOI: 10.1016/j.envpol.2018.04.072
- [37] Li H, Qu R, Yan L, Guo W, Ma Y. Field study on the uptake and translocation of PBDEs by wheat (*Triticum aestivum* L.) in soils amended with sewage sludge. *Chemosphere*. 2015;**123**:87-92. DOI: 10.1016/j.chemosphere.2014.12.045
- [38] Hites RA. Polybrominated diphenyl ethers in the environment and in people: a meta-analysis of concentrations. *Environmental Science & Technology*. 2004;**38**:945-956. DOI: 10.1021/es035082g
- [39] Sjödin A, Carlsson H, Thuresson K, Sjölin S, Bergman A, Ostman C. Flame retardants in indoor air at an electronics recycling plant and at other work environments. *Environmental Science & Technology*. 2001;**35**:448-454. DOI: 10.1021/es000077n
- [40] Li P, Wu H, Li Q, Jin J, Wang Y. Brominated flame retardants in food and environmental samples from a production area in China: concentrations and human exposure assessment. *Environmental monitoring and assessment*. 2015;**187**:719. DOI: 10.1007/s10661-015-4947-y
- [41] Ueno D, Darling C, Alae M, Pacepavicius G, Teixeira C, Campbell L, Letcher RJ, Bergman A, Marsh G, Muir D. Hydroxylated polybrominated diphenyl ethers (OH-PBDEs) in the abiotic environment: surface water and precipitation from Ontario, Canada. *Environmental Science & Technology*. 2008;**42**:1657-1664. DOI: 10.1021/es7021279
- [42] Kuch B, Schneider C, Rupp S, Recke Rvd, Bopp K, Metzger JW. Polybromierte Diphenylether und Tetrabrombisphenol A: Untersuchungen zum Abbau und Metabolismus, Bestimmung in Nahrungsmitteln: Abschlussbericht BWPLUS. Karlsruhe: Landesanstalt für Umwelt Baden-Württemberg (LUBW); 2005
- [43] Zhu H, Wang Y, Tam NFY. Microcosm study on fate of polybrominated diphenyl ethers (PBDEs) in contaminated mangrove sediment. *Journal of hazardous materials*. 2014;**265**:61-68. DOI: 10.1016/j.jhazmat.2013.11.046
- [44] Wang S, Zhang S, Huang H, Lu A, Ping H. Debrominated, hydroxylated and methoxylated metabolism in maize (*Zea mays* L.) exposed to lesser polybrominated diphenyl ethers (PBDEs). *Chemosphere*. 2012;**89**:1295-1301. DOI: 10.1016/j.chemosphere.2012.05.026
- [45] Pan L, Sun J, Wu X, Wei Z, Zhu L. Transformation of hydroxylated and methoxylated 2,2',4,4',5-brominated diphenyl ether (BDE-99) in plants. *Journal of Environmental Sciences*. 2016;**49**:197-202. DOI: 10.1016/j.jes.2016.06.017
- [46] Sun J, Liu J, Yu M, Wang C, Sun Y, Zhang A, Wang T, Lei Z, Jiang G. In vivo metabolism of 2,2',4,4'-tetrabromodiphenyl ether (BDE-47) in young whole pumpkin plant. *Environmental Science & Technology*. 2013;**47**:3701-3707. DOI: 10.1021/es4003263
- [47] Stapleton HM, Kelly SM, Pei R, Letcher RJ, Gunsch C. Metabolism of polybrominated diphenyl ethers (PBDEs) by human hepatocytes in vitro. *Environmental health perspectives*. 2009;**117**:197-202. DOI: 10.1289/ehp.11807
- [48] Chen J, Zhou HC, Wang C, Zhu CQ, Tam NF-Y. Short-term enhancement effect of nitrogen addition on microbial degradation and plant uptake of

polybrominated diphenyl ethers (PBDEs) in contaminated mangrove soil. *Journal of hazardous materials*. 2015;**300**:84-92. DOI: 10.1016/j.jhazmat.2015.06.053

[49] Chow KL, Man YB, Tam NFY, Liang Y, Wong MH. Removal of decabromodiphenyl ether (BDE-209) using a combined system involving TiO<sub>2</sub> photocatalysis and wetland plants. *Journal of hazardous materials*. 2017;**322**:263-269. DOI: 10.1016/j.jhazmat.2016.05.097

[50] Jiang J, Zhao H, Sun S, Wang Y, Liu S, Xie Q, Li X. Occurrence and profiles of halogenated phenols, polybrominated diphenyl ethers and hydroxylated polybrominated diphenyl ethers in the effluents of waste water treatment plants around Huang-Bo Sea, North China. *The Science of the total environment*. 2018;**622-623**:1-7. DOI: 10.1016/j.scitotenv.2017.11.323

[51] Zhao M, Zhang S, Wang S, Huang H. Uptake, translocation, and debromination of polybrominated diphenyl ethers in maize. *Journal of Environmental Sciences*. 2012;**24**:402-409. DOI: 10.1016/S1001-0742(11)60748-1

[52] Dobslaw D, Woiski C, Kiel M, Kuch B, Breuer J. Plant uptake, translocation and metabolism of PBDEs in plants of food and feed industry: A review. *Reviews in Environmental Science and Bio/Technology*. DOI: 10.1007/s11157-020-09557-7

[53] Li H, Shao F, Qiu Y, Ma Y. Solubility, uptake, and translocation of BDE 47 as affected by DOM extracted from agricultural wastes. *Environmental Science and Pollution Research*. 2019;**26**:19871-19878. DOI: 10.1007/s11356-019-05393-7

[54] Cheng Z, Wang Y, Wang S, Luo C, Li J, Chaemfa C, Jiang H, Zhang G. The influence of land use

on the concentration and vertical distribution of PBDEs in soils of an e-waste recycling region of South China. *Environmental Pollution*. 2014;**191**:126-131. DOI: 10.1016/j.envpol.2014.04.025

[55] Farzana S, Cheung SG, Tam NFY. Effects of aquaculture effluents on fate of 2,2',4,4',5-pentabromodiphenyl ether (BDE-99) in contaminated mangrove sediment planted with *Kandelia obovata*. *The Science of the total environment*. 2019;**691**:71-79. DOI: 10.1016/j.scitotenv.2019.07.057

[56] Wang Y, Luo C, Li J, Yin H, Zhang G. Influence of plants on the distribution and composition of PBDEs in soils of an e-waste dismantling area: Evidence of the effect of the rhizosphere and selective bioaccumulation. *Environmental Pollution*. 2014;**186**:104-109. DOI: 10.1016/j.envpol.2013.11.018

[57] Feng J, Shen X, Chen J, Shi J, Xu J, Tang C, Brookes PC, He Y. Improved rhizoremediation for decabromodiphenyl ether (BDE-209) in E-waste contaminated soils. *Soil Ecology Letters*. 2019;**1**:157-173. DOI: 10.1007/s42832-019-0007-9

[58] Wang S, Wang Y, Song M, Luo C, Li J, Zhang G. Distributions and compositions of old and emerging flame retardants in the rhizosphere and non-rhizosphere soil in an e-waste contaminated area of South China. *Environmental Pollution*. 2016;**208**:619-625. DOI: 10.1016/j.envpol.2015.10.038

[59] Wang S, Wang Y, Luo C, Li J, Yin H, Zhang G. Plant selective uptake of halogenated flame retardants at an e-waste recycling site in southern China. *Environmental Pollution*. 2016;**214**:705-712. DOI: 10.1016/j.envpol.2016.04.071

[60] Tian M, Chen S-J, Wang J, Luo Y, Luo X-J, Mai B-X. Plant uptake of atmospheric brominated flame retardants at an E-waste site in southern China. *Environmental Science &*

Technology. 2012;**46**:2708-2714. DOI: 10.1021/es203669n

[61] Huang H, Zhang S, Christie P, Wang S, Xie M. Behavior of decabromodiphenyl ether (BDE-209) in the soil-plant system: Uptake, translocation, and metabolism in plants and dissipation in soil. *Environmental Science & Technology*. 2010;**44**:663-667. DOI: 10.1021/es901860r

[62] Jian K, Zhao L, Ya M, Zhang Y, Su H, Meng W, Li J, Su G. Dietary intake of legacy and emerging halogenated flame retardants using food market basket estimations in Nanjing, eastern China. *Environmental pollution (Barking, Essex: 1987)*. 2020;**258**:113737. DOI: 10.1016/j.envpol.2019.113737

[63] Xiang L, Song Y, Bian Y, Liu G, Herzberger A, Gu C, Jiang X, Wang F. Manure amendment reduced plant uptake and enhanced rhizodegradation of 2,2',4, 4'-tetrabrominated diphenyl ether in soil. *Biology and Fertility of Soils*. 2018;**54**:807-817. DOI: 10.1007/s00374-018-1304-7

[64] Wu J, Yi Y, Fang Z, Tsang EP. Effects of biochar on phytotoxicity and translocation of polybrominated diphenyl ethers in Ni/Fe bimetallic nanoparticle-treated soil. *Environmental science and pollution research international*. 2018;**25**:2570-2579. DOI: 10.1007/s11356-017-0627-5

[65] Kuch B, Rupp S, Fischer K, Kranert M, Metzger JW. Untersuchungen von Komposten und Gärsubstraten auf organische Schadstoffe in Baden-Württemberg: Abschlussbericht. Karlsruhe: Landesanstalt für Umwelt Baden-Württemberg (LUBW); 2007

[66] Teuten EL, Rowland SJ, Galloway TS, Thompson RC. Potential for plastics to transport hydrophobic contaminants. *Environmental Science & Technology*. 2007;**41**:7759-7764. DOI: 10.1021/es071737s

[67] Xu P, Ge W, Chai C, Zhang Y, Jiang T, Xia B. Sorption of polybrominated diphenyl ethers by microplastics. *Marine Pollution Bulletin*. 2019;**145**:260-269. DOI: 10.1016/j.marpolbul.2019.05.050

[68] Bao L-J, You J, Zeng EY. Sorption of PBDE in low-density polyethylene film: implications for bioavailability of BDE-209. *Environmental Toxicology and Chemistry*. 2011;**30**:1731-1738. DOI: 10.1002/etc.564

[69] Ma J, Zhang Q, Chen F, Zhu Q, Wang Y, Liu G. Remediation of PBDEs-metal co-contaminated soil by the combination of metal stabilization, persulfate oxidation and bioremediation. *Chemosphere*. 2020;**252**:126538. DOI: 10.1016/j.chemosphere.2020.126538

[70] Yao Y, Wang B, He Y, Wang L, Corvini PF-X, Ji R. Fate of 4-bromodiphenyl ether (BDE3) in soil and the effects of co-existed copper. *Environmental pollution (Barking, Essex: 1987)*. 2020;**261**:114214. DOI: 10.1016/j.envpol.2020.114214

[71] Yan Y, Ma M, Zhang W, Ma W, Li M, Yan L. Effect of elevated nitrate on biodegradation of pentabromodiphenyl ether (BDE-99) and change in microbial communities during groundwater recharge with tertiary-treated municipal wastewater. *International Biodeterioration & Biodegradation*. 2017;**124**:128-137. DOI: 10.1016/j.ibiod.2017.03.022

[72] Pi N, Wu Y, Zhu HW, Wong YS, Tam NFY. The uptake of mixed PAHs and PBDEs in wastewater by mangrove plants under different tidal flushing regimes. *Environmental Pollution*. 2017;**231**:104-114. DOI: 10.1016/j.envpol.2017.07.085

[73] Lu M, Zhang Z-Z. Phytoremediation of soil co-contaminated with heavy metals and deca-BDE by co-planting of

*Sedum alfredii* with tall fescue associated with *Bacillus cereus* JP12. *Plant and Soil*. 2014;**382**:89-102. DOI: 10.1007/s11104-014-2147-0

[74] Chen F, Zeng S, Ma J, Li X, Zhang S, Zhu Q. Interactions between decabromodiphenyl ether and lead in soil-plant system. *Chemosphere*. 2019;**236**:124406. DOI: 10.1016/j.chemosphere.2019.124406

[75] Li H, Li X, Xiang L, Zhao HM, Li YW, Cai QY, Zhu L, Mo CH, Wong MH. Phytoremediation of soil co-contaminated with Cd and BDE-209 using hyperaccumulator enhanced by AM fungi and surfactant. *The Science of the total environment*. 2018;**613-614**:447-455. DOI: 10.1016/j.scitotenv.2017.09.066

[76] Li H, Huang WX, Gao MY, Li X, Xiang L, Mo CH, Li YW, Cai QY, Wong MH, Wu FY. AM fungi increase uptake of Cd and BDE-209 and activities of dismutase and catalase in amaranth (*Amaranthus hypochondriacus* L.) in two contaminants spiked soil. *Ecotoxicology and environmental safety*. 2020;**195**:110485. DOI: 10.1016/j.ecoenv.2020.110485

[77] Li Y, Chiou CT, Li H, Schnoor JL. Improved prediction of the bioconcentration factors of organic contaminants from soils into plant/crop roots by related physicochemical parameters. *Environment international*. 2019;**126**:46-53. DOI: 10.1016/j.envint.2019.02.020

[78] Collins CD, Finnegan E. Modeling the plant uptake of organic chemicals, including the soil-air-plant pathway. *Environmental Science & Technology*. 2010;**44**:998-1003. DOI: 10.1021/es901941z

[79] Hwang J-I, Lee S-E, Kim J-E. Comparison of theoretical and experimental values for plant uptake

of pesticide from soil. *PloS one*. 2017;**12**:e0172254. DOI: 10.1371/journal.pone.0172254

[80] Briggs GG, Bromilow RH, Evans AA. Relationships between lipophilicity and root uptake and translocation of non-ionised chemicals by barley. *Pesticide Science*. 1982;**13**:495-504. DOI: 10.1002/ps.2780130506

[81] Bintein S, Devillers J, Karcher W. Nonlinear dependence of fish bioconcentration on n-octanol/water partition coefficient. SAR and QSAR in environmental research. 1993;**1**:29-39. DOI: 10.1080/10629369308028814

[82] Meylan WM, Howard PH, Boethling RS, Aronson D, Printup H, Gouchie S. Improved method for estimating bioconcentration/bioaccumulation factor from octanol/water partition coefficient. *Environmental Toxicology and Chemistry*. 1999;**18**:664-672. DOI: 10.1002/etc.5620180412

[83] Nendza M. QSARs of bioconcentration - validity assessment of log P/log BCF correlations. Weinheim: VCH; 1991. DOI: 10.13140/2.1.1837.5364

[84] Liagkouridis I, Cousins AP, Cousins IT. Physical-chemical properties and evaluative fate modelling of 'emerging' and 'novel' brominated and organophosphorus flame retardants in the indoor and outdoor environment. *The Science of the total environment*. 2015;**524-525**:416-426. DOI: 10.1016/j.scitotenv.2015.02.106