

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,600

Open access books available

138,000

International authors and editors

170M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Citrus Fruits: Nutritive Value and Value-Added Products

Abu Saeid and Maruf Ahmed

Abstract

Citrus fruits are essential sources of food and energy and play a critical role in supplementing healthy diets. Citrus fruits contain mostly carbohydrates such as sucrose, glucose, and fructose and are good dietary fiber sources, which help prevent gastrointestinal disease and promote high circulating cholesterol. Besides, citrus fruits are also significant sources of vitamin C and various bioactive compounds. It is suggested that these components are of vital importance in improving human health due to their antioxidant properties and being converted to vitamin A. However, citrus fruit is still being used for different purposes like juice, jam, jelly, squash, pies, cake, candies, marmalades, etc. Most citrus waste materials are currently used as animal feed. Innovations are occurring in the conversion of citrus by-products into valuable commodities with the development of innovative technologies. This chapter has put up primary and secondary research findings of citrus fruits, especially lemon and pomelo, their chemical properties, composition, and their use in health and cosmetic needs.

Keywords: citrus, lemon, pomelo, nutritional properties, value-added products

1. Introduction

Citrus is an evergreen shrub that belongs to the Rutaceae family from South Asia, China, India and the Malay Archipelago, which is native to the subtropical and tropical regions of Asian regions [1]. The genus of citrus includes sweet orange (*C. sinensis*: 61.1 % of world citrus production), tangerine (*C. reticulata*: 19.9 %), limon and lime (*C. limon* and *C. aurantifolia*: 12.1 %) and grapefruit (*C. paradisi*: 5%). Minor types of citrus, which constitute much of the remaining 2.0%, include sour orange (*C. quarantium*), shaddocks (*C. grandis*), citrus (*C. medica*), which seem to be promising sources for many beneficial human nuts [2]. Citrus fruit is divided into two sections like peel and flesh (**Figure 1**). Peel is made from epicarp or flavedo (colored peripheral surface) and mesocarp or albedo (white soft middle layer). The peel (60–65%), internal tissues (30–35%), and seeds (0–10%) comprise citrus fruits [3]. Citrus fruits provide carbohydrates, such as sucrose, glucose, and fructose mostly. Fresh citrus fruits are also an immeasurable source of dietary fiber associated with gastrointestinal disease prevention and lowered circulating cholesterol. Citrus fruits also have a distinct aroma and delicious taste along with low protein and fat content.

Citrus fruits also provide the most potent source of vitamins C and B (thiamines, pyridoxines, niacins, riboflavin, pantothenic acids, and folate). The fruit also leads to the use of phytochemicals, such as carotenoids, flavonoids, and limonoids [1].




Figure 1.
Structure of the citrus fruit [2].

Citrus phytochemicals contain antibacterial, antiviral, antifungal, anti-carcinogenic, anti-thrombotic, or anti-inflammatory agents [4]. Several studies have proposed citrus fruit evaluation as a healthy and delicious diet [5]. Prior research suggested that citrus and citrus products are rich sources of vitamins, minerals, and dietary fibers [6]. However, the bioactive and non-nutrient compounds in citrus are appreciated to reduce the risk of various chronic diseases [7].

Citrus fruits are eaten as fresh goods and juice throughout the world. Peel is discarded as waste containing many secondary components with significant antioxidant activity related to other fruit portions [8]. In recent years, flavonoids such as polymethoxy flavones (PMFs), which are present in citrus fruits, have been attracted growing attention by their antioxidants [9] and anti-cancer properties [10]. Various bioactive compounds in citrus peel extract and powder may reduce overall cholesterol, triglycerides, LDL, and glucose levels [11]. Citrus by-products produce a range of value-added products, including essential oils, pectin, enzymes, single-cell collagen, natural antioxidants, ethanol, organic acids, and prebiotics. Orange, lemon, mandarin, and grapefruit contained essential oils show antifungal activity upon the fungi *A. niger*, *A. flavus*, *P. chrysogenum*, and *P. verrucosum*. The essential oil may be regarded as acceptable for the food industry as alternatives to chemicals [12]. Pectin extracted from *Citrus* peel is used in various industrial food processes as gelling agents, including jam, jellies, and as thickener, texturizer, emulsifier, and stabilizer in dairy products. Pectin is also used to jellify properties in the pharmaceutical, dental, and cosmetic industries [13]. Therefore, this chapter highlighted the nutritional values of major essential nutrients such as Vitamin C, carotenoids and vitamin A, Folate, Dietary fiber, flavonoids, and limonoids, as well as value-added products such as food ingredients, pectin, essential oil, enzymes, a natural antioxidant, and packaging film retrieved from citrus especially lemon and pomelo fruit.

2. Characteristics of citrus fruits (lemon and pomelo) and their chemical compositions

2.1 Lemon

Lemon (*Citrus limon* L. from Rutaceae) is one of the most common globally and ranks third among the Citrus species globally by 4,200,000 metric tons after orange and mandarin [14]. Lemon fruits typically consist of three parts: pulp, skins (albedo and flavedo), and seeds. It offers an extensive supply of natural compound

products such as citric acid, ascorbic acid, minerals, flavonoids and essential oils [15]. Lemon bioactive compounds like flavonoids, vitamins, minerals, dietary fiber (**Table 1**), and essential oils are used in the food, cosmetic, and pharmaceutical industries. Most by-products of the lemon juice industry can provide functional foods with nutritional substances such as non-digestible carbohydrates, dietary fiber and bioactive (flavonoids and ascorbic acid). Lemon fruits can function against photo-oxidation because carotenoids exist. Lemon fruit, rich in flavonoids, has a significant role in the healthy diet, particularly in preventing diseases such as obesity, diabetes, lowering blood lipids, cardiovascular disease, and some forms of cancer [18]. The citrus fruits used for direct consumption or converted into juices, jam, jelly, molasses, lemoncello beverage and more in addition to the lemon skin are added value products such as pectins, essential oil and functional ingredients [12, 18].

2.2 Pomelo

Pomelo is one of the most commonly grown and eaten citrus fruits and orange, mandarin, lemon, and grapefruit [19]. Pomelo (**Table 1**) is a promising source of carbohydrates, proteins, fiber, vitamins and minerals originating in warm tropical climates in south-eastern Asia [20]. The presence of bioactive (carotenoids, lycopene,

Component	Lemon (<i>Citrus limon</i>)	Pomelo (<i>Citrus maxima</i>)
Moisture (g/100g)	84.2	87.0
Fiber (g/100g)	1.6	1.60
Carbohydrate (g/100g)	10.8	11.5
Protein (g/100g)	0.9	0.6
Fat (g/100g)	0.8	0.20
Vitamin (mg/100g)		
β-carotene	50.0	120
Thiamine	0.02	0.03
Riboflavin	0.01	0.03
Vitamin C	37	20.0
Mineral (mg/100g)		
Ca	70.0	10.0
Mg	12.0	21.6
Na	1.50	2.70
P	10.0	20
K	148	106
Fe	0.23	0.40
Zinc	0.12	0.15
Cu	0.20	0.19
Total phenol (mg GAE/g)	204.40	70.56
Total flavonoids (mg QUE/g)	2750	13.06
Carotenoid (mg/100 mL)	0.31-0.35	0.72-0.73

Table 1.
 Chemical composition of Citrus fruits as [15–17].

polyphenols, flavonoids, limonoids, fiber and vitamin C) contributes to their protection against oxidative stress, hyperglycemia, and high blood pressure. Due to its essential health promotion properties, pomelo segments in food products are growing in importance in producing functional foods [21]. Pomelo is eaten fresh or made into juice [19], or pomelo fortified noodles help the diabetic population [21]. On the other hand, researchers have investigated alternative ways of restoring pomelo peels to the advantage of value-added products such as pectin, essential oils, polysaccharides, phytochemicals [19]. Production of juice and consumption of fresh fruit create large quantities of agricultural waste. The main components of wet Pomelo Peel waste, like other citrus fruits, include water, cellulose and hemicellulose, soluble sugars, lipids (mainly D-limonene), and bioactive compounds (i.e., polyphenols, mostly flavonoids).

3. Nutritional values of citrus fruits

Citrus has many natural plant compounds such as vitamin C, carotenoids (some can convert to vitamin A), folic acid, flavonoids, and fiber. **Table 2** shows the amount of vitamin and mineral consumption in lemon and pomelo fruits.

3.1 Vitamin C (ascorbic acid)

Citrus is a valuable source of vitamin C. By consuming a moderate amount of citrus fruits each day, an individual can achieve 100 percent Vitamin C level. Vitamin C is an essential water-soluble vitamin essential for the body's defense [22]. It is transmitted through muscle fibers, carnitine biosynthesis, neurotransmitters, collagen, and bones because these particles connect the fibers. The immune system

	Vitamin C	Vitamin A	Folate	Fiber
Oranges	53-88 mg	17 µg	30 µg	2.4 g
Children under 9 y (%)	213-589	3-6	15-20	10-13
Persons 9+ y	59-195	2-4	8-10	6-11
Pregnant/Lactating women	44-110	2-3	5-6	8-9
Grapes	31-61 mg	58 µg	13 µg	1.6 g
Children under 9 y	125-244	12-15	7-9	6-8
Persons 9+	35-135	6-10	3-4	4-8
Pregnant/lactating women	26-76	4-8	2-3	6
Tangerines	27-72 mg	46-144 µg	16 µg	1.8 g
Children under 9	107-480	9-36	8-11	7-9
Persons 9+	30-160	5-24	4-5	5-9
Pregnant/lactating women	21-90	4-19	3-4	6
Lemons/limes	29-61 mg	2-22 µg	11-16 µg	1.8-2.8 g
Children under 9	116-407	0.4-6	4-7	11-15
Persons 9+	32-135	0.2-4	2-4	9-13
Pregnant/lactating women	24-76	0.2-3	1-2	10

Table 2.

The number of nutrients and the percent of the recommended daily allowance or adequate intake met from the consumption of 100 g of selected citrus fruit [22].

can be effectively stimulated by consuming vitamin C, which boosts white blood cells [23]. When Vitamin C is taken for pregnancy, it can decrease pre-eclampsia risk [24]. Some studies indicate that vitamin C supplementation can reduce the severity of colds symptoms or duration [23]. Anti-oxidants such as Vitamin C could reduce the risk of artery stiffening and cardiovascular diseases [25]. Above 200 mg of vitamin C daily is a healthy intake, and citrus fruits are a huge source of this vitamin. Lemon provides 37 mg of ascorbic acid per 100 g of fruit [16]. Pomelos have 52.3 mg of ascorbic acid in 100 g of the flesh [26].

3.2 Carotenoids and Vitamin A

There are many types of carotenoids, including terpenes responsible for pigments commonly found in plants, and there are about 600 carotenoids in foods and 50 in human bodies [27, 28]. The highest carotenoid levels, such as lutein, zeaxanthin, lycopene, and vitamin A, are found in fruits and vegetables, including orange and carotene. Benefits of carotenoids in foods include improving immune function, promoting bone formation, promoting eye health, and maintaining visual quality [22]. There is a large amount of data supporting that carotenoids reduce the risk of cancer, macular degeneration, cataracts, skin damage to the sun, and cardiovascular diseases [29]. Higher consumption of β -carotene is linked to a lower breast cancer risk [30]. Beta carotene, lycopene, or lutein may decrease the rate of UV-induced lipid peroxidation in human skin fibroblast cells [30]. Lutein is inversely related to colorectal cancer in both men and women [31]. The levels of lutein, zeaxanthin, β -cryptoxanthin, and β -carotene in the lemon and pomelo, were around 2.95, 0.81, 0.81 and 10.3 ($\mu\text{g/g}$, db), respectively [32]. The content of carotenoids in pummelos' peel was 0.012-0.015 mg/gdb [33].

3.3 Folate (folic acid)

Folic acid, which is a water-soluble vitamin, and its derivatives are collectively called folate or folacin. The most notable folate compounds in Citrus are the reduced 5-methyl tetrahydrofolate (monoglutamate) and polyglutamate compounds [34]. Folate plays a vital role in DNA, which is involved in homocysteine regulation and protein production primarily through the methylation transfer reactions [22]. Because there is a high DNA production during pregnancy, a folate deficiency is significantly linked to birth defects such as neural tube defects [35]. Lack of folic acid caused higher levels of homocysteine, raising heart disease and atherosclerosis [22]. Previous studies show that citrus fruits' daily consumption can help improve folate levels, which will subsequently decrease blood homocysteine (tHcy), thus reducing cardiovascular disorder and neural tube defects [36]. Citrus is a parallel source of dietary folate that can help to cover up to 10% to 20% of the recommended daily allowance of adults, children, and infants with a consumption of 100 g of citrus fruits. The consumption of citrus fruit is an easy way to obtain vitamin C and dietary folate, which is vital for absorption in the body. Lemon, a citrus fruits representative, has eleven to sixteen micrograms of Folate in 100 grams [22]. According to El-Otmani and Ait-Oubahou [37] Citrus limon contained 11mg of folic acid per 100 g of citrus.

3.4 Dietary fibre

The fiber is found in vegetables and fruits cannot be digested and absorb in the small intestine. There are two kinds of dietary fiber; soluble and insoluble fiber. Insoluble fibers are highly fermentable and connected with carbohydrate and lipid

metabolism, while soluble fibers contribute to fecal bulk and reduce transit time [34]. Although pectin, cellulose, and hemicellulose comprise the most abundant dietary fiber on the plants, they also contain only trace amounts of lignin. Pectin is citrus' primary fiber, which occurs primarily in citrus peels and rinds. Consumption of citrus fruit can contribute significant quantities of pectin in a diet. Dietary incorporation of pectin appears to affect several metabolic and digestive processes; principal interest affects glucose absorption and cholesterol level [38, 39]. There is a significant benefit in consuming citrus fruit because of its pectin content. Dietary incorporation of pectin appears to have many implications for metabolic, digestive, and health affairs. One way fiber can reduce colon cancer is by diluting and trapping the harmful chemicals in the colon from bile-absorption and bile-excretion [34]. Scientific studies have proven that fiber can help promote laxation and satiety, the uptake and reabsorption of glucose, fat, cholesterol, and bile acids, thereby lessening heart disease risk and possibly enhancing healthy intestinal microbial fermentation [40, 41]. Citrus fruits significantly reduce cholesterol levels depending on the esterification degree of fiber consumption, viscosity, and molecular mass [22]. A fiber-rich diet has a low risk of deadly chronic diseases such as diabetes, heart disease, weight, and cancer and lowers cholesterol levels and blood sugar [42]. Several epidemiological studies reported that citrus peel support reducing plasma liver cholesterol, total serum cholesterol, serum triglyceride levels, and total liver lipids [43].

3.5 Flavonoids and limonoids

Citrus pulp, peel are rich sources of flavonoids. Toh et al. [44] found that pomelo peeled (356.95 mg/QE) had higher total flavonoid content than pomelo pulp (13.06 mg/QE). Makni et al. [15] found the amount of quercetin in lemon flesh (56.16 mg Eq Quercetin/g dry weight) was higher than in peel (27.50 mg Eq Quercetin/g dry weight). Citrus fruits are also rich in flavonoids such as hesperidin, hesperetin, naringin, naringenin, diosmin, quercetin, rutin, nobiletin, tangeretin, and others [45]. Citrus flavonoids have both antioxidant and anti-inflammatory properties, and because of that, it can increase the antioxidant capacity and effect reducing cholesterol and triglycerides levels and provide more excellent bone health [22]. Preclinical studies and clinical trials demonstrated that flavonoids' effects in the forms of hesperidin and its aglycone hesperetin prevent various types of diseases, including neurological, psychiatric, and cardiovascular disorders [46]. Over the years, naringin and hesperidin are gaining attention for their great antioxidant capacity, contributing sweet flavor to foods and beverages [47]. Some known naturally occurring flavonoids have potency in defending against certain types of RNA and DNA viruses [48].

Limonoids are also known as flavonoids, which are compounds found in citrus fruits. In citrus fruits, there are two groups of limonoids: aglycones and their corresponding glucosides. Bitter taste in citrus results from limonoids present. Limonin's most important constituents are glycosides called limonin and nomilin [22]. In animal and human cell lines, limonoids slow down the development of aggressive cancers like the pancreas, colon, stomach, and breasts. On the other hand, limonoids are also reported to reduce skin cancer in animal models. Limonoids are known for their medicinal or health beneficial effects like anti-cancer, anti-microbial and antimalarial activities [49]. Limonoids have antibacterial and antiviral effects. Some limonoids are known to stimulate the *in vivo* production of the detoxifying enzyme glutathione S-transferase in the liver and inhibit the formation of chemically induced tumor cells in the oral cavity, forestomach, small intestine, colon, lung, and skin of




Figure 2.
 Proposed model for signaling pathways leading to growth inhibition by obacunone in estrogen-responsive breast cancer (MCF-7) cells [53].

animals [50, 51]. Limonoids found in citrus fruits decreased the spread of cancer cells in animal studies [52]. Studies showed that lemon-lime oil in the form of obacunone appeared to prevent breast cancer by inhibiting aromatase enzyme and anti-inflammatory pathways [53] (**Figure 2**). Several studies have revealed that limonin and nomilin are found in fruits, pulp, and seeds. Pummelo juice contained 18 ppm of limonin and 29 ppm of total limonoid glucosides [54]. Wattanasiritham et al. [55] reported limonin content of 18 ppm in the juice of pomelo cultivars. Limonin levels in extracted juice from seven pummelo cultivars from Florida ranged from 10.07 to 29.62 ppm. As shown by analysis, mature lemon seeds contain 1300 µg/g of different limonoid glucosides on a fresh weight basis [56]. The mature seeds contain much higher amounts of glucosides than commercialized juice, therefore. Fong et al. [57] documented that the commercial lemon juice had only 82 µg/g of glucosides.

4. Value-added products

Citrus fruits are known for being highly fragrant, with a tart taste and higher vitamin C content. The world has a wider variety of citrus fruit because of the continued type changes, such as sour oranges, oranges, pummelos, lemons, and limes, among others [22]. Nowadays, citrus pulp/pomace, seed, and peel are used for various commercially valuable products such as food ingredients, pectin, essential oils, enzyme production, a natural antioxidant, and packaging film formation.

4.1 Food ingredient or food products

The nutritional supplement of pomelo fruit segments has been added to products to be developed like noodles prepared with 30% new segments and 5% dry. These noodles can satisfy those with regular diabetes and the general public [21]. A high dietary fiber food was created by reducing the dietary fiber-rich pomelo peel to a powder that contained nearly 50% of dietary fiber. Lemon fruit is usually eaten fresh, but it is also processed to make juices, jams, jellies, molasses, candies and much more [18]. Lemon juice has been used as a coagulant during the manufacture of wara cheese [58]. Another innovation implemented in the beef burger is to use lemons for “enhancing the cooking properties of the burger” [59]. Lario et al. [60] had reported that the high-fiber lemon powder extract from lemon peel debris by-products is an ideal additive for food products (as meat, dairy, and bakery products).

4.2 Pectin

Pectin is an agent of gelling, emulsifying, stabilizing, texturizing, which appears as a white to light brown powder broadly accepted as a functional ingredient [61]. Fruit peels are a highly desirable pectin source because they cover up to 20% of the fruit's total Pectin [62]. Pomelo is a highly valued source of natural Pectin. About 20.75% pectin is derived from lemon peel for jams. In the study, the high extraction (36.71%) of Pectin from lemon peel has something to contribute to this industry [63]. Moneim et al. [64] recommended utilizing 20.75% of the lemon peels' total weight in making pumpkin jam. The researchers were added 16.740% of Pectin from pomelo peel to the pressed carrots before storage [65]. Studies by Methacanon et al. [66] have shown that pectin yield was 23.19% for pomelo peel. On the other hand, Roy et al. [65] were found that pomelo peels are a good source of Pectin, and then carrot jam made by extracting Pectin from pomelo peel.

4.3 Essential oil

Essential oils (Eos) are volatile, complex, natural mixture of aromatic oils obtained from plants [67]. Citrus essential oil is commonly known to produce a good fragrance and has been officially approved for healthy public consumption. All over the world, EOs are used in cosmetics, perfumery, toiletries, flavoring, beverage, pharmaceuticals and other personal hygiene products [68–70]. Lemon oil is often used on the skin because of its antimicrobial and antifungal properties. Pomelo peel (PP) has approximately 299 recognized volatile compounds. A significant number of these volatile compounds is considered as terpenoids (189 volatiles, 63.2%). Various kinds of chemicals released are monoterpenoids, monocyclic monoterpenoids, bicyclic monoterpenoids, diterpenoids, acyclic sesquiterpenoids, monocyclic sesquiterpenoids, bicyclic sesquiterpenoids, and tricyclic sesquiterpenoids. Another major volatile present in PP EO is nonterpenoid alcohols (4.7%), nonterpenoid aldehydes (6.0%), nonterpenoid hydrocarbons (5.7%), and esters (8.7%). The unknown volatiles covered 11.7% (35 volatiles) of total volatile compounds. The structures of widely known terpenoids in Polypropylene EO has shown in **Figure 3a,b**. The most critical monoterpenoids (1 to 16) and sesquiterpenoids (17 to 29) are present in PP EO [19]. According to studies conducted, lemon essential oils retain aroma in foods because of their natural preservative and flavoring properties [71]. The effect of lemon essential oils on the cheesemaking process dramatically reduces microorganisms' population, especially those of the Enterobacteriaceae family [58]. In the food and pharmaceutical industries, citrus




Figure 3.
 (a) Major monoterpenoids in pomelo peel essential oils (1-16); (b) Major sesquiterpenoids in pomelo peel essential oils (17-29).

EOs can be employed to inhibit mold and fungal growth. Lemon EO (Citrus limon) was used as a possible fungicide to manage the pathogenic fungi attacking grapevines, namely *Eutypa* sp., *Botryosphaeria dothidea*, and *Fomitiporia mediterranea*. The antifungal activity was observed for EO against all the three fungi with the highest action against strain *Eutypa* sp. (82% inhibition) and the lowest tolerance (33.1%) towards *F. Mediterranean* [72]. These essential oils repress the growth of mold and yeast. The raspberries coated with alginate and lemon EO (0.2 percent) or orange EO (0.1 percent) halted bacteria, yeast, and mold and also reduced the quality deterioration right after harvest [73]. The lemon EOs mixed in the chitosan films can be used to control *L. Multicellular* pathogens in refrigerated foods. Researchers Rahmawati et al. [74] observed that an edible coating applied with lemon essential oil only delayed aging of tofu and fresh strawberry.

4.4 Enzymes

The most basic usage of citrus peels is to produce the pectinolytic enzyme for beneficial purposes. Larios et al. [75] studied endo-polygalacturonase production by *Aspergillus* sp. CH-Y-1043 using untreated lemon peel and citrus pectin as carbon sources. Lemon peel being employed as a substrate in submerged cultures to obtain high pectinase titers by *A. flavipes* FP-500 and *A. terreus* FP-370 [76].

A. niger produces approximately 2,181.23 U/L pectinases from lemon, peel pomace in a solid-state reactor [77]. Seyis and Aksoz [78] have shown that lemon pomace and peel are suitable substrates for heterotrophic xylanases enzyme production using fungus *Trichoderma harzianum*. *Aspergillus niger* LFP-1 was studied in solid-state fermentation (SSF) using pomelo (*Citrus grandis*) peels as a substrate [79]. Maller, et al. [80] determined that lemon peels are extremely capable of triggering the production of Polygalacturonase in the *aspergillus niveus*. Pectin lyase yield increased through fungal strain *Aspergillus oryzae* process derived from lemons peel and used in solid-state fermentations [81]. Studies said that Polemo pericarp powder utilized as a substrate for *Aspergillus oryzae* JMU316 has Naringinase enzyme [82]. Pectinase enzyme produced from pomelo peels by *Aspergillus niger* through Solid State Fermentation [83]. Lemon peels could be a good source of naringin, which could be used as a carbon source in submerged fermentation for naringinase production using *Aspergillus niger* [84]. Naringinase is essential for the production of sweetener precursors, preparation of prunin, aroma enhancement in winemaking, biotransformation of antibiotics, and rhamnose manufacturing [82].

4.5 Natural antioxidant

Antioxidants are chemical substances that can reduce or prevent the damage caused by free radicals in the body, thus reducing the risks of cardiovascular disease and cancer [85]. Results of studies showed that lemon peel contained almost 75.9% of antioxidant content. The unique ability of Paneer was derived from compounds found in the Peel of orange, lemon and pomegranate [85]. Peel taken from Tambun White pomelo type contains higher levels of antioxidants and is also a rich source of natural antioxidants [44]. Lemon peel and flesh had the highest antioxidant capacity, and they had a significant impact on the prevention of cardiovascular diseases and other diseases [86].

4.6 Packaging film formation

More sustainable, biodegradable plastic has gained popularity among environmental scientists. The researcher created plastic films from citrus peels. By applying the peels of citrus, biodegradable packaging material could be made [87]. Wu et al. [88] prepared fruit peel as the edible packaging film with high content biopolymer to form film for packaging. The film was designed to incorporate tea polyphenols, which causes interacting molecules to become more closely crowded. Soy protein with essential oil of lemon peel was used to create a degradable film and create cheese curdorsants for preservation [89]. Dias et al. [90] reported that the use of citrus essential oil and its aroma significantly improved consumers' health and significantly increased the acceptance of biscuits' packaging. Das et al. [91] demonstrated that chicken feather keratin combined with pomelo peel pectin to form biodegradable composite film and wrapping of fried fish fillets resulted in less weight loss, hardness value, and reduction in the surface microbial count.

5. Conclusion

Citrus has positive effects on human health, and it could be an essential raw material to the biotechnological industry. Citrus is a mighty source of vitamins, minerals, and dietary fibers. Bioactive and non-nutrient compounds in citrus are valuable for controlling chronic diseases such as diabetes, cholesterol, obesity, cardiovascular disease, and some forms of cancer. Besides citrus, vitamin C also

has other benefits, including fighting diseases such as cardiovascular disease, boosting white blood cells, immune function, and symptoms or duration of colds. Peel, flesh/pomace, and seed from the citrus fruit are employed in making different novel foods like noodles, extract pectin, enzyme extracts, and essential oil. Therefore this information might be necessary for the readers because it gives facts about the popular citrus fruit. Also, choosing the best citrus for an edible ingredient can be beneficial for citrus processors.

IntechOpen

Author details

Abu Saeid¹ and Maruf Ahmed^{2*}

¹ Department of Food Engineering, NPI University of Bangladesh, Manikganj, Bangladesh

² Department of Food Processing and Preservation, Hajee Mohammad Danesh Science and Technology University, Dinajpur, Bangladesh

*Address all correspondence to: maruf@hstu.ac.bd; maruffpp@gmail.com

IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Liu Y, Heying E, Tanumihardjo SA. History, Global Distribution, and Nutritional Importance of Citrus Fruits. *Comprehensive Reviews in Food Science and Food Safety*. 2012; **11**: 530-545. DOI: 10.1111/j.1541-4337.2012.00201.x
- [2] Mamma D, Christakopoulos P. Biotransformation of Citrus By-Products into Value Added Products. *Waste and Biomass Valorization*. 2014; **5**: 529-549. DOI: 10.1007/s12649-013-9250-y
- [3] Alnaimy A, Gad AE, Mustafa MM, Atta MAA, Basuony HAM. Using of Citrus By-products in Farm Animals Feeding. *Open Access Journal of Science*. 2017; **1**: 58-67. DOI: 10.15406/oajs.2017.01.00014
- [4] Abobatta FW. Nutritional Benefits of Citrus Fruits. *American Journal of Biomedical Science & Research*. 2019; **3**: 303-306. DOI: 10.34297/ajbsr.2019.03.000681
- [5] Economos C, Clay WD. Nutritional and health benefits of citrus fruits. *Food, Nutrition and Agriculture – FAO*. 2005; **24**: 11-18. <http://www.fao.org/3/a-x2650t/X2650t03.pdf>
- [6] Ghanem N, Mihoubi D, Kechaou N, Mihoubi NB. (2012). Microwave dehydration of three citrus peel cultivars: Effect on water and oil retention capacities, color, shrinkage and total phenols content. *Industrial Crops and Products*. 2012; **40**: 167-177. DOI: 10.1016/j.indcrop.2012.03.009
- [7] Bamise CT, Oziegbe EO. Laboratory Analysis of pH and Neutralizable Acidity of Commercial Citrus Fruits in Nigeria Department of Child Dental Health, Faculty of Dentistry. 2013; **7**: 72-76. DOI: 10.5829/idosi.abr.2013.7.4.73175
- [8] Manthey JA, Grohmann K. Phenols in Citrus Peel Byproducts. Concentrations of Hydroxycinnamates and Polymethoxylated Flavones in Citrus Peel Molasses. *Journal of Agricultural and Food Chemistry*. 2001; **49**:3268-3273. DOI: 10.1021/jf010011r
- [9] Chen XM, Tait AR, Kitts DD. Flavonoid composition of orange peel and its association with antioxidant and anti-inflammatory activities. *Food Chemistry*. 2017; **218**:15-21. DOI: 10.1016/j.foodchem.2016.09.016
- [10] Wang L, Wang J, Fang L, Zheng Z, Zhi D, Wang S, Li S, Ho CT, Zhao H. Anticancer activities of citrus peel polymethoxyflavones related to angiogenesis and others. *BioMed Research International*. 2014; **2014**:1-10. DOI: 10.1155/2014/453972
- [11] Ashraf H, Butt MS, Iqbal MJ, Suleria HAR. Citrus peel extract and powder attenuate hypercholesterolemia and hyperglycemia using rodent experimental modeling. *Asian Pacific Journal of Tropical Biomedicine*. 2017; **7**: 870-880. DOI: 10.1016/j.apjtb.2017.09.012
- [12] Viuda-Martos M, Ruiz-Navajas Y, Fernández-López J, Pérez-Álvarez J. Antifungal activity of lemon (*Citrus lemon L.*), mandarin (*Citrus reticulata L.*), grapefruit (*Citrus paradisi L.*) and orange (*Citrus sinensis L.*) essential oils. *Food Control*. 2008; **19**: 1130-1138. DOI: 10.1016/j.foodcont.2007.12.003
- [13] Pagán J, Ibarz A, Llorca M, Pagán A, Barbosa-Cánovas GV. Extraction and characterization of pectin from stored peach pomace. *Food Research International*. 2001; **34**: 605-612. DOI: 10.1016/S0963-9969(01)00078-3
- [14] Boluda-Aguilar M, López-Gómez A, Production of bioethanol by fermentation of lemon (*Citrus limon L.*) peel wastes pretreated with steam explosion. *Industrial Crops*

- and Products. 2013; **41**: 188-197. DOI: 10.1016/j.indcrop.2012.04.031
- [15] Makni M, Jemai R, Kriaa W, Chtourou Y, Fetoui H. Citrus limon from Tunisia: Phytochemical and Physicochemical Properties and Biological Activities. *BioMed Research International*. 2018; **2018**: 1-10. DOI: 10.1155/2018/6251546
- [16] Paul DK, Shaha RK. Nutrients, Vitamins and Minerals Content in Common Citrus Fruits in the Northern Region of Bangladesh. *Pakistan Journal of Biological Sciences*. 2004; **7**: 238-242. DOI: 10.3923/pjbs.2004.238.242
- [17] Kumar D, Ladaniya MS, Gurjar M. Underutilized Citrus sp. Pomelo (*Citrus grandis*) and Kachai lemon (*Citrus jambhiri*) exhale in phytochemicals and antioxidant potential. *Journal of Food Science and Technology*. 2019; **56**: 217-223. DOI: 10.1007/s13197-018-3477-3
- [18] González-Molina E, Domínguez-Perles R, Moreno DA, García-Viguera C. Natural bioactive compounds of Citrus limon for food and health. *Journal of Pharmaceutical and Biomedical Analysis*. 2010; **51**: 327-345. DOI: 10.1016/j.jpba.2009.07.027
- [19] Tocmo R, Pena-Fronteras J, Calumba KF, Mendoza M, Johnson JJ. Valorization of pomelo (*Citrus grandis* Osbeck) peel: A review of current utilization, phytochemistry, bioactivities, and mechanisms of action. *Comprehensive Reviews in Food Science and Food Safety*. 2020; **19**: 1969-2012. DOI: 10.1111/1541-4337.12561
- [20] Mäkynen K, Jitsaardkul S, Tachasamran P, Sakai N, Puranachoti S, Nirojsinlapachai N, Chattapat V, Caengprasath N, Ngamukote S, Adisakwattana S. Cultivar variations in antioxidant and antihyperlipidemic properties of pomelo pulp (*Citrus grandis* [L.] Osbeck) in Thailand. *Food Chemistry*. 2013; **139**: 735-743. DOI: 10.1016/j.foodchem.2013.02.017
- [21] Reshmi SK, Sudha ML, Shashirekha MN. Noodles fortified with Citrus maxima (pomelo) fruit segments suiting the diabetic population. *Bioactive Carbohydrates and Dietary Fibre*. 2020; **22**:100213. DOI: 10.1016/j.bcdf.2020.100213
- [22] Turner T, Burri B. Potential Nutritional Benefits of Current Citrus Consumption. *Agriculture*. 2013; **3**: 170-187. DOI:10.3390/agriculture3010170
- [23] Wintergerst ES, Maggini S, Hornig DH. Immune-enhancing role of vitamin C and zinc and effect on clinical conditions. *Annals of Nutrition and Metabolism*. 2006, **50**: 85-94. DOI: 10.1159/000090495
- [24] Chappell LC, Seed PT, Briley AL, Kelly FJ, Lee R, Hunt BJ, Parmar K, Bewley SJ, Shennan AH, Steer PJ, Poston L. Effect of antioxidants on the occurrence of pre-eclampsia in women at increased risk: A randomized trial. *Lancet* 1999, **354**: 810-816. DOI: 10.1016/S0140-6736(99)80010-5
- [25] Wang YC, Chuang YC, Ku YH. Quantitation of bioactive compounds in citrus fruits cultivated in Taiwan. *Food Chemistry*. 2007; **102**: 1163-1171. DOI: 10.1016/j.foodchem.2006.06.057
- [26] Haque MN, Saha BK, Karim MR, Bhuiyan MNH. Evaluation of Nutritional and Physico-Chemical Properties of Several Selected Fruits in Bangladesh. *Bangladesh Journal of Scientific and Industrial Research*. 2009; **44**: 353-358. DOI: 10.3329/bjsir.v44i3.4410
- [27] Britton G. Structure and properties of carotenoids in relation to function. *The FASEB Journal*. 1995; **9**: 1551-1558. DOI:10.1096/fasebj.9.15.8529834
- [28] Khachik F, Christopher JS; Smith JC, Louise MC, Steck A, Pfander H.

- Identification, quantification, and relative concentrations of carotenoids and their metabolites in human milk and serum. *Analytical Chemistry*. 1997; **69**: 1873-1881. DOI:10.1021/ac961085i
- [29] Aust O, Sies H, Stahl W, Polidori MC. Analysis of lipophilic antioxidants in human serum and tissues: tocopherols and carotenoids. *Journal of Chromatography A*. 2001; **936**: 83-93. DOI: 10.1016/s0021-9673(01)01269-9
- [30] Fraser PD, Bramley PM. The biosynthesis and nutritional uses of carotenoids. *Progress in Lipid Research*. 2004; **43**: 228-265. DOI: 10.1016/j.plipres.2003.10.002
- [31] Slattery ML, Benson J, Curtin K, Ma KN, Schaeffer D, Potter JD. Carotenoids and colon cancer. *The American Journal of Clinical Nutrition*. 2000; **71**: 575-582. DOI: 10.1093/ajcn/71.2.575
- [32] Wang YC, Chuang YC, Hsu HW. The flavonoid, carotenoid and pectin content in peels of citrus cultivated in Taiwan. *Food Chemistry*. 2008; **106**: 277-284. DOI: 10.1016/j.foodchem.2007.05.086
- [33] Xu CJ, Fraser PD, Wang WJ, Bramley PM. Differences in the carotenoid content of ordinary citrus and lycopene-accumulating mutants. *Journal of Agricultural and Food Chemistry*. 2006; **54**: 5474-5481. DOI: 10.1021/jf060702t
- [34] Silalahi J. Anticancer and health protective properties of citrus fruit components. *Asia Pacific Journal of Clinical Nutrition*. 2002; **11**:79-84. DOI: 10.1046/j.1440-6047.2002.00271.x
- [35] Green NS. Folic acid supplementation and prevention of birth defects. *The Journal of nutrition*. 2002; **132**: 2356S-2360S. DOI: 10.1093/jn/132.8.2356S
- [36] Brouwer IA, Van Dusseldorp M, West CE, Meyboom S, Thomas CMG, Duran M, Van Het Hof KH, Eskes TKAB, HautvastJGAJ, Steegers-TheunissenRPM. Dietary folate from vegetables and citrus fruit decreases plasma homocysteine concentrations in humans in a dietary controlled trial. *Journal of Nutrition*. 1999; **129**: 1135-1139. DOI: 10.1093/jn/129.6.1135
- [37] El-Otmani M, Ait-Oubahou A. Citrus spp.: orange, mandarin, tangerine, clementine, grapefruit, pomelo, lemon and lime. Elhadi MY, editor. In *Postharvest Biology and Technology of Tropical and Subtropical Fruits*. Woodhead Publishing Limited: 2011; p.437-514. DOI: 10.1533/9780857092762.437
- [38] Baker RA. Potential Dietary Benefits of Citrus Pectin and Fibre. *Food Technology*. 1994; **11**: 133-139.
- [39] Silalahi J. Hypocholesterolemic Factors in Foods. A Review. *Indonesian Food and Nutrition Progress*. 2000; **7**: 26-35.
- [40] Brown L, Rosner B, Willett WW, Sacks FM. Cholesterol-lowering effects of dietary fiber: A meta-analysis. *The American Journal of Clinical Nutrition*. 1999; **69**: 30-42. DOI: 10.1093/ajcn/69.1.30
- [41] Dikeman CL, Fahey GC. Viscosity as related to dietary fiber: A review. *Critical Reviews in Food Science and Nutrition*. 2006; **46**: 649-663. DOI: 10.1080/10408390500511862
- [42] Anderson JW, Baird P, Davis RH Jr, Ferreri S, Knudtson M, Koraym A, Waters V, Williams CL. Health benefits of dietary fiber. *Nutrition Reviews*. 2009; **67**:188-205. DOI: 10.1111/j.1753-4887.2009.00189.x

- [43] Terpstra AHM, Lapré JÁ, de Vries HT, Beynen AC. The hypocholesterolemic effect of lemon peels, lemon pectin, and the waste stream material of lemon peels in hybrid F1B hamsters. *European Journal of Nutrition*. 2002; **41**: 19-26. DOI:10.1007/s003940200002
- [44] Toh JJ, Khoo HE, Azrina A. Comparison of antioxidant properties of pomelo [*Citrus Grandis* (L) Osbeck] varieties. *International Food Research Journal*. 2013; **20**: 1661-1668.
- [45] Mahmoud AM, Hernández Bautista RJ, Sandhu MA, Hussein OE. Beneficial effects of citrus flavonoids on cardiovascular and metabolic health. *Oxidative Medicine and Cellular Longevity*. 2019; **2019**: 1-19. DOI: 10.1155/2019/5484138
- [46] Li C, Schluesener H. Health-promoting effects of the citrus flavanone hesperidin. *Critical Reviews in Food Science and Nutrition*. 2015; **57**: 613-631. DOI:10.1080/10408398.2014.906382
- [47] Hung PV, Nhi NHY, Ting LY, Phi NTL. *Chemical composition and biological activities of extracts from pomelo peel by-products under enzyme and ultrasound-assisted extractions*. *Journal of Chemistry*. 2020; **2020**: 1-7. DOI: 10.1155/2020/1043251
- [48] Puri M, Banerjee UC. Production, purification, and characterization of the debittering enzyme naringinase. *Biotechnology Advances*. 2000; **18**: 207-217. DOI: 10.1016/S0734-9750(00)00034-3
- [49] Roy A, Saraf S. Limonoids: Overview of Significant Bioactive Triterpenes Distributed in Plants Kingdom. *Biological and Pharmaceutical Bulletin*. 2006; **29**: 191-201. DOI:10.1248/bpb.29.191
- [50] Manners GD, Jacob RA, Breksa, Schoch TK, Hasegawa S. Bioavailability of Citrus Limonoids in Humans. *Journal of Agricultural and Food Chemistry*. 2003; **51**: 4156-4161. DOI: 10.1021/jf0300691
- [51] Yu J, Wang L, Walzem RL, Miller EG, Pike LM, Patil BS. Antioxidant Activity of Citrus Limonoids, Flavonoids, and Coumarins. *Journal of Agricultural and Food Chemistry*. 2005; **53**: 2009-2014. DOI:10.1021/jf0484632
- [52] Zunino SJ, Storms DH, Freytag TL, Adkins YC, Bonnel EL, Woodhouse LR, Breksa AP, Manners GD, Mackey BE, Kelley DS. Dietary supplementation with purified citrus limonin glucoside does not alter ex vivo functions of circulating T lymphocytes or monocytes in overweight/obese human adults. *Nutrition Research*. 2016; **36**: 24-30. DOI: 10.1016/j.nutres.2015.10.011
- [53] Kim J, Jayaprakasha GK, Patil BS. Obacunone exhibits anti-proliferative and anti-aromatase activity in vitro by inhibiting the p38 MAPK signaling pathway in MCF-7 human breast adenocarcinoma cells. *Biochimie*. 2014; **105**: 36-44. DOI:10.1016/j.biochi.2014.06.002
- [54] Ohta H, Hasegawa S. Limonoids in Pummelos [*Citrus grandis* (L.) Osbeck]. *Journal of Food Science*. 1995; **60**: 1284-1285. DOI:10.1111/j.1365-2621.1995.tb04574.x
- [55] Wattanasiritham L, Taweek K, Ratanachinakorn B. Limonin and Naringin in Pummelos (*Citrus grandis* (L.) Osbeck) In: 31st Congress on science and technology of Thailand, Suranaree University of Technology, Thailand. 2005
- [56] Fong CH, Hasegawa S, Herman Z, Ou P. Biosynthesis of limonoid glucosides in lemon (*Citrus limon*). *Journal of the Science of Food and Agriculture*.

1991; **54**: 393-398. DOI: 10.1002/jsfa.2740540310

[57] Fong CH, Hasegawa S, Herman Z, Ou P. Limonoid Glucosides in Commercial Citrus Juices. *Journal of Food Science*. 1989; **54**:1505-1506. DOI:10.1111/j.1365-2621.1989.tb05146.x

[58] Adetunji VO, Alonge DO, Singh RK, Chen J. Production of wara, a West African soft cheese using lemon juice as a coagulant. *LWT - Food Science and Technology*. 2008; **41**: 331-336. DOI:10.1016/j.lwt.2007.02.012

[59] Aleson-Carbonell L, Fernández-López J, Pérez-Alvarez JA, Kuri V. Characteristics of beef burger as influenced by various types of lemon albedo. *Innovative Food Science and Emerging Technologies*. 2005; **6**: 247-255. DOI: 10.1016/j.ifset.2005.01.002

[60] Lario Y, Sendra E, García-Pérez J, Fuentes C, Sayas-Barberá E, Fernández-López J, Pérez-Alvarez JA. Preparation of high dietary fiber powder from lemon juice by-products. *Innovative Food Science and Emerging Technologies*. 2004; **5**: 113-117. DOI: 10.1016/j.ifset.2003.08.001

[61] Mesbahi G, Jamalian J, Farahnaky A. A comparative study on functional properties of beet and citrus pectins in food systems. *Food Hydrocolloids*. 2005; **19**: 731-738. DOI: 10.1016/j.foodhyd.2004.08.002

[62] Chavan P, Singh AK, Kaur G. Recent progress in the utilization of industrial waste and by-products of citrus fruits: A review. *Journal of Food Process Engineering*. 2018; **41**:1-10. DOI: 10.1111/jfpe.12895

[63] Kanmani P, Dhivya E, Aravind J, Kumaresan K. Extraction and analysis of pectin from citrus peels: augmenting the yield from citrus limon using statistical experimental design. *Iranica Journal of Energy and Environment*.

2014; **5**: 303-312. DOI: 10.5829/idosi.ijee.2014.05.03.10

[64] Moneim A, Sulieman E, Khodari KMY, Salih ZA. Extraction of pectin from lemon and orange fruits peels and its utilization in jam making. *International Journal of Food Science and Nutrition Engineering*. 2013; **3**: 81-84. DOI: 10.5923/j.food.20130305.01

[65] Roy MC, Alam M, Saeid A, Das BC, Mia MB, Rahman MA, Eun JB, Ahmed M. Extraction and characterization of pectin from pomelo peel and its impact on nutritional properties of carrot jam during storage. *Journal of Food Processing and Preservation*. 2018; **42**: 1-9. DOI: 10.1111/jfpp.13411

[66] Methacanon P, Krongsin J, Gamonpilas C. Pomelo (*Citrus maxima*) pectin: Effects of extraction parameters and its properties. *Food Hydrocolloids*. 2014; **35**: 383-391. DOI: 10.1016/j.foodhyd.2013.06.018

[67] Figueiredo CA, Barroso JG, Pedro LG, Scheffer JJC. Factors affecting secondary metabolite production in plants: volatile components and essential oils. *Flavour and Fragrance Journal*. 2008; **23**:213-226. DOI: 10.1002/ffj.1875

[68] Palazzolo E, Laudicina VA, Germanà MA. Current and potential use of citrus essential oils. *Current Organic Chemistry*. 2013; **17**: 3042-3049. DOI: 10.2174/13852728113179990122

[69] Sarkic A, Stappen I. Essential oils and their single compounds in cosmetics—A critical review. *Cosmetics*. 2018; **5**:1-11. DOI: 10.3390/cosmetics5010011

[70] Mahato N, Sharma K, Koteswararao R, Sinha M, Baral ER, Cho MH. Citrus essential oils: Extraction, authentication and application in food preservation.

Critical Reviews in Food Science and Nutrition. 2019; **59**: 611-625. DOI: 10.1080/10408398.2017.1384716

[71] Klimek-Szczykutowicz M, Szopa A, Ekiert H. Citrus limon (lemon) phenomenon—a review of the chemistry, pharmacological properties, applications in the modern pharmaceutical, food, and cosmetics industries, and biotechnological studies. *Plants*. 2020; **9**: 119; DOI:10.3390/plants9010119

[72] Ammad F, Moumen O, Gasem A, Othmane S, Hisashi KN, Zebib B, Merah O. The potency of lemon (*Citrus limon L.*) essential oil to control some fungal diseases of grapevine wood. *Comptes Rendus Biologies*, 2018; **341**: 97-101. DOI: 10.1016/j.crv.2018.01.003

[73] Gomes MDS, Cardoso MDG, Guimarães ACG, Guerreiro AC, Gago CML, Vilas Boas EVDB, Dias CMB, Manhita ACC, Faleiro ML, Miguel MGC. Effect of edible coatings with essential oils on the quality of red raspberries over shelf-life. *Journal of the Science of Food and Agriculture*. 2017; **97**:929-938. DOI: 10.1002/jsfa.7817

[74] Rahmawati D, Chandra M, Santoso S, Puteri MG. Application of lemon peel essential oil with edible coating agent to prolong shelf life of tofu and strawberry. *AIP Conference Proceedings*. American Institute of Physics. 2017; 0200371-5 DOI: 10.1063/1.4973164

[75] Larios G, Garcia JM, Huitrón C. Endo-polygalacturonase production from untreated lemon peel by *Aspergillus sp.* CH-Y-1043. *Biotechnology Letters*. 1989; **11**:729-734. DOI: 10.1007/BF01044106

[76] Martínez-Trujillo A, Arreguín-Range L, García-Rivero M, Aguilar-Osorio G. Use of fruit residues for pectinase produc-

tion by *Aspergillus flavipes* FP-500 and *Aspergillus terreus* FP-370. *Letters in Applied Microbiology*. 2011; **53**: 202-209. DOI: 10.1111/j.1472-765X.2011.03096.x

[77] Ruiz HA, Rodríguez-Jasso RM, Rodríguez R, Contreras-Esquivel JC, Aguilar CN. Pectinase production from lemon peel pomace as support and carbon source in solid-state fermentation column-tray bioreactor. *Biochemical Engineering Journal*. 2012; **65**:90-95. DOI: 10.1016/j.bej.2012.03.007

[78] Seyis I, Aksoz N. Xylanase production from *Trichoderma harzianum* 1073 D3 with alternative carbon and nitrogen sources. *Food Technology and Biotechnology*. 2005; **43**:37-40.

[79] Darah I, Taufiq MMJ, Lim SH. Pomelo citrus grandis (L.) Osbeck peel as an economical alternative substrate for fungal pectinase production. *Food Science and Biotechnology*. 2013; **22**: 1683-1690. DOI: 10.1007/s10068-013-0267-6

[80] Maller A, Damásio AR, da Silva TM, Jorge JA, Terenzi HF, Polizeli M. Biotechnological potential of agro-industrial wastes as a carbon source to thermostable polygalacturonase production in *aspergillus niveus*. *Enzyme Research*. 2011; **2011**: 1-6. DOI: 10.4061/2011/289206

[81] Koser S, Anwar Z, Iqbal Z, Anjum A, Aqil T, Mehmood S, Irshad M. Utilization of *Aspergillus oryzae* to produce pectin lyase from various agro-industrial residues. *Journal of Radiation Research and Applied Sciences*. 2014; **7**: 327-332. DOI: 10.1016/j.jrras.2014.05.001

[82] Chen DX, Niu TG, Cai HN. Optimizing culture medium for debittering constitutive enzyme naringinase production by *Aspergillus oryzae* JMU316. *African Journal of Biotechnology*. 2010; **9**: 4970-4978.

- [83] Hussain K, Wajid A, Babar ME, Anwar Z, Farooqi S, Siddiqa A, Noreen S, Iqbal J. Production and optimization of pectinase from pomelo by *Aspergillus niger* through solid state fermentation. *Annals of Life Sciences*. 2019; **6**: 21-41.
- [84] Igbonekwu A, Omeje KO, Ezugwu AL, Eze SOO NO, Chilaka CF. Characterization of Nariginase obtained from *Aspergillus niger* by submerged fermentation using naringin extracted from lemon peels. *Research and Development in Material Science*. 2018; **4**: 420-424. DOI: 10.31031/rdms.2018.04.000599
- [85] Immanuel G, Singh S. Extraction of antioxidants from fruit peels and its utilization in paneer. *Journal of Food Processing and Technology*. 2014; **5**:1-5. DOI: 10.4172/2157-7110.1000349
- [86] Gorinstein S, Martín-Belloso O, Park YS, Haruenkit R, Lojek A, Íž M, Caspi A, Libman I, Trakhtenberg S. Comparison of some biochemical characteristics of different citrus fruits. *Food Chemistry*. 2001; **74**: 309-315. DOI: 10.1016/S0308-8146(01)00157-1
- [87] Shinde M, Sonawane SK, Patil S. Fruit peel utilization in food packaging. *Indian Food Industry Mag*. 2019; **1**:19-24
- [88] Wu H, Lei Y, Zhu R, Zhao M, Lu J, Xia D, Jiao C, Zhang Z, Shen G, Li S. Preparation and characterization of bioactive edible packaging films based on pomelo peel flours incorporating tea polyphenol. *Food Hydrocolloids*. 2019; **90**: 41-49. DOI: 10.1016/j.foodhyd.2018.12.016
- [89] Al-Sahlany STG. Production of biodegradable film from soy protein and essential oil of lemon peel and use it as cheese preservative. *Basrah Journal of Agricultural Sciences*. 2017; **30**: 27-35. DOI: 10.37077/25200860.2017.40
- [90] Dias MV, de Medeiros HS, Nilda de FFS, de Melo NR, Borges SV, João de DSC, Pereira JMT de AK. Development of low-density polyethylene films with lemon aroma. *LWT - Food Science and Technology*. 2013; **50**: 167-171. DOI: 10.1016/j.lwt.2012.06.005
- [91] Das P, Borah PP, Badwaik LS. Transformation of chicken feather keratin and pomelo peel pectin into biodegradable composite film. *Journal of Polymers and the Environment*. 2018; **26**:2120-2129. DOI: 10.1007/s10924-017-1109-z