

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,300

Open access books available

130,000

International authors and editors

155M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# Promising Detoxification Approaches to Mitigate Aflatoxins in Foods and Feeds

Vishakha Pandey

## Abstract

Aflatoxins are a group of naturally occurring carcinogenic mycotoxins produced by certain *Aspergillus* species in nuts, grains, oilseeds and vegetables. Ingestion of aflatoxin contaminated food and feed has extremely negative health implications in humans and livestock. Additionally, exporting countries face the trade barrier due to strict regulations in international market to maintain food quality. This led to huge economic losses to global exporters. Therefore, there is an urgent need for development of effective methods for detoxification of aflatoxins from food and feed to ensure food security. Till date, numerous methods for detoxification of aflatoxins from foods have been employed. Physical, chemical and biological treatment are the novel promising approaches for partial/complete detoxification of aflatoxins from the foodstuffs. In this chapter, we will address the efficacy and shortcomings of each methods in with respect to economic importance, human health and food security.

**Keywords:** Aflatoxin, mycotoxin, carcinogenic, detoxification, food security

## 1. Introduction

Aflatoxins are toxic secondary metabolites, mainly produced by many species of *Aspergillus*, namely *Aspergillus parasiticus*, *A. flavus*, *A. nomius*, *A. stollatus* [1, 2]. They contaminate various crops (wheat, maize, cotton), dried fruits, spices, meat and milk products [3, 4]. Aflatoxins were first discovered as a causative agent of Turkey X disease in early 1960s, led to the death of numerous turkey in England [5]. Generally, *A. flavus* specifically produces B-type aflatoxins while *A. parasiticus* produce B- as well as G-type aflatoxins [6]. Among these four aflatoxins, AFB1 being most dangerous, accounts for more than 75% of all food and feed related aflatoxin contamination [7]. Cytochrome P450 mediated metabolism of AFB1 in liver, resulting in its epoxidation to AFB1-exo-8, 9-epoxide and AFB1-endo-8, 9-epoxide, demethylation to aflatoxin P1 (AFP1) and hydroxylation to aflatoxin Q1 (AFQ1) and aflatoxin M1 (AFM1) [8]. The International Agency for Research on Cancer (IARC) has classified aflatoxin type B and G as Group 1 carcinogen and AF-M1 as Group 2B [9]. Derived from polyketides, consumption of aflatoxin contaminated food and feed cause a range of serious health complications in humans and animals, together named as aflatoxicosis [10, 11]. Short term exposure to high dose of aflatoxins results in jaundice, hemorrhage, liver damage and subsequent death

and long term exposure to sublethal levels of aflatoxins cause nutritional disorders, immunosuppression, cancer [12].

The challenges related to aflatoxins can be overcome by adopting innovative and novel strategies. The risk of aflatoxins can be reduced by preventing the contamination of foods from aflatoxins at the pre-harvest stage and removing/eliminating the aflatoxins from aflatoxin contaminated food at the post-harvest stage. Prevention strategies, including the use of pesticides, fertilizers, maintaining optimum temperature and moisture for storage, right harvesting time, are only partly responsible for achieving aflatoxin-free food and feed. In order to eliminate aflatoxin completely, post-harvest strategies are followed. The post-harvest strategies include cleaning, sorting, milling and dehulling [13], treatment at temperature between 237 and 306°C [14], mineral binders such as, montmorillonite, zeolite, aluminosilicate, bentonite that bind aflatoxins. Such binders may partially or completely counteract the toxicity of dietary aflatoxins [15]. Elliott et al. [16] have reported the cytotoxic effects induced by mineral binders like DNA damage, reduced cell viability, apoptosis, oxidative stress.

## **2. Novel strategies for aflatoxin degradation**

In the past decades, numerous promising novel strategies for aflatoxin mitigation have been developed. They are broadly categorized as physical, chemical and biological approaches. Physical strategies involve the utilization of radiations and cold plasma for the fast aflatoxin degradation [17]. Chemical methods include the treatment with electrolyzed oxidizing water, organic acids, ozone and natural plant extracts. These are methods have been widely used in several countries such as USA and China [18]. Microbial and enzymatic based conversion of highly toxic aflatoxins into less toxic or non-toxic metabolites are included in biological methods [19]. In this chapter, we will address each of these novel technologies for aflatoxin degradation in detail.

### **2.1 Physical treatment for aflatoxins degradation**

#### *2.1.1 Irradiation*

In the recent times, ionizing irradiation (viz. electron beam, gamma and ultraviolet rays) and nonionizing irradiation (viz. infrared waves, radio waves, visible light waves and microwaves) has been employed extensively for the degradation of aflatoxin present in the food and feed (**Table 1**).

Electron beam irradiation (EBI) technology has great potential for aflatoxin degradation. EBI technology offers the advantage of high effectiveness, low equipment cost, dosage control, short processing time, low heat generation, few variables and in-line processing [20]. EBI technology has been applied for degradation of aflatoxins in coconut agar [21]. Efficiency of EBI technology for degradation of aflatoxin is lesser than that of  $\gamma$  radiation. [22] found that EBI at doses of 10 and 5 kGy decreased the content of AFB1 in Brazilnuts by 65.7 and 53.3%, respectively, whereas  $\gamma$  irradiation at same doses led to reduction in AFB1 by 84.2 and 70.6%, respectively. Liu et al. [23] used EBI dose of 300 kGy for reduction of AFB1 by 70%. As the selected dose was ten times of the maximum permissible dosage allowed by FDA, so this method is not highly efficient and preferable in AFB1 degradation in peanut meal.

Gamma ( $\gamma$ ) rays has been the most preferred radiation source for the food owing to its high penetrability and reactivity. Treatment of food by gamma rays of

Physical Degradation method	Food product treated	Aflatoxin ( $\mu\text{g kg}^{-1}$ or $\mu\text{g L}^{-1}$ )	Degradation Percentage	Parameters for treatment	Reference
Electron beam	Brazil nut	AFB1 (4.8)	65.7	300 kGy	[22]
	Peanut	AFB1 (1000)	70.0	300 kGy	[23]
<sup>60</sup> Co gamma Irradiation	Red chili	AFB1 (11–35)	86–98	6 kGy	[28]
	Cattle feed	AFB1 (50)	85	10 kGy	[26]
	Corn	AFB1 (57–1210)	85.6–98.6	10 kGy	[26]
	Brazil nut	AFB1 (4.8)	84.2	10 kGy	[22]
	Peanut	AFB1 (300)	43	9 kGy	[114]
	White pepper	AFB1 (60), AFB2 (18), AFG1 (60) and AFG2 (18)	50.6, 35.2, 47.7 and 42.9	30 kGy	[31]
	Almond	AFB1 (20), AFB2 (20), AFG1 (20) and AFG2 (20)	19.3, 11.0, 21.1 and 16.6	15 kGy	[115]
UV irradiation	Peanut	AFB1 (2000)	100	220–400 nm at 0.8 mW cm <sup>-2</sup> for 80 min	[23]
	Peanut	AFB1 (350)	99.1	254 nm for 10 h	[116]
	Peanut oil	AFB1 (128)	96	365 nm at 55–60 mW cm <sup>-2</sup> for 20 min	[117]
	Peanut oil	AFB1 (52.0)	86.1	365 nm at 6.4 mW cm <sup>-2</sup> for 10 min	[118]
	Peanut oil	AFB1 (2000)	100	220–400 nm at 0.8 mW cm <sup>-2</sup> for 30 min	[119]
	Red chili powder	AFB1 (1872)	87.8	365 nm for 60 min	[120]
Pulsed light	Rice bran	AFB1 (36) and AFB2 (4.4)	90.3 and 86.7	0.52 J cm <sup>-1</sup> per pulse for 15 s	
	Rough rice	AFB1 (132) and AFB2 (45)	75.0 and 39.2	0.52 J cm <sup>-1</sup> per pulse for 80 s	[44]
Microwave heating	Peanut	AFB1 (5–183) and AFB2 (7–46.7)	50–60 and 100	Heating in microwave oven at 92°C for 5 min	[49]
	Corn flour	AFB1 (100)	67.7	Heating in microwave oven for 10 min	[50]
	Alkalized corn	AFB1 (22.5) and AFB2 (69.6)	36 and 58	1650 W	[48]

**Table 1.**  
 Physical methods for aflatoxin degradation in food and feed.

upto 10 kGy has no toxicological, or microbiological hazards [24]. Additionally,  $\gamma$  irradiation results in the interaction of high energy of  $\gamma$  rays with the water present in the food products. This produces highly reactive free radicals such as superoxide radical ( $\text{O}_2^{\bullet-}$ ), hydrogen ( $\text{H}^{\bullet}$ ) radical and hydroxyl ion ( $\text{OH}^-$ ) that in turn destroy

aflatoxins and also attack DNA of pathogenic microbes [14, 25]. Markov *et al.* [26] have used high-energy photons from cobalt-60 (gamma source) for destruction of pathogenic microbes by directly damaging DNA of microbial cells.

Many studies in the literature showed that treatment of food commodities with  $\gamma$  rays ranging from 5 to 10 kGy led to degradation of significant amount of aflatoxins. For instance, irradiation with  $\gamma$  rays at low dose of upto 6 kGy has reduced Aflatoxin B1 (AFB<sub>1</sub>) level in red chillies and fruits for around 90% [27, 28]. The AFB<sub>1</sub> level reduced for about 95% in maize seed samples with  $\gamma$  irradiation dose of 10 kGy [26]. AFB<sub>1</sub> in Brazil nuts can be eliminated upto 84.2% by irradiation of  $\gamma$  rays at 5 to 10 kGy [22]. Likewise,  $\gamma$  irradiation at dose of 10 kGy decreased ochratoxin A (OTA) in coffee beans and dry-cured meat for almost 100% and 22.5% respectively [29, 30]. Nevertheless, few studies in the literature concluded that  $\gamma$  irradiation could not effectively eliminate aflatoxins in food. For e.g. irradiation of black and white pepper with  $\gamma$  rays at 10 kGy dose has no significant effect on its aflatoxins content [31]. Gamma irradiation of poultry feed at 15 kGy dose resulted in 13.6, 11.0, 21.1, 18.2% decrease in AFG2, AFB2, AFG1 and AFB1, respectively [32]. However, the efficiency of  $\gamma$  rays - mediated aflatoxin degradation depends on numerous factors such as concentration of mycotoxin, dose of radiation, content of water, air humidity, composition of food and type and number of fungal strains [25, 31].

The advantage  $\gamma$  irradiation offers is high capacity for microbial inactivation that reduces the microbial load and increases shelf life of food. Gamma irradiation technology has been approved by more than 55 countries such as Japan, USA, European countries, China for food processing [33]. This technology is not preferable with high vitamin and lipid content because polyunsaturated fatty acids undergo extensive peroxidation in unsaturated bonds, leading to increased oxidative rancidity [34].

Apart from being an economical non-thermal technology for Aflatoxin decontamination, Ultraviolet (UV) irradiation is also highly cost effective and eco-friendly [35]. Treatment of food products with moderate doses of UV rays has no negative impact on its sensory and physicochemical properties [36]. Though UV rays can efficiently penetrate into transparent or clear liquids, its penetration efficiency through solids is limited. Therefore, granular or opaque foods has to be in the form of thin layer for decontamination by UV rays [37]. UV light effectively removed of Patulin (PAT) from apple cider and juice. Zhu *et al.* [38] have used different wavelengths of UVC for PAT reduction in apple juice. They found UVC of 222 nm wavelength was most effective. Exposure to UV affected the taste of apple cider and juice. Intensity and duration of UV irradiation are important factors that affect the elimination efficiency of aflatoxins elimination efficiency Irradiation with short wavelength (254 nm) and long wavelength (362 nm) UV rays for 30 minutes resulted in complete elimination of AF-B1 and AF-G1 in wheat grains, whereas exposure to same dose of short and long wavelength of UV rays for 2 hours reduced AF-B2 in wheat grains by 50 and 74% respectively [39]. Exposure of pistachio, almond and groundnut with UVC at 265 nm for 15 minutes led to 100% removal of AF-G2 from all the nut samples and complete degradation of AF-G1 in pistachio and almond. UV-C irradiation at 265 nm for 45 minutes showed 97% degradation of AF-B1 [40]. Treatment with UV-A and UV-B rays can also be used for reduction of mycotoxins produced from *A. parasiticus* and *A. carbonarius* in pistachio and grape media [41].

Non-thermal Pulsed light (PL) technology has been used for degradation of aflatoxins in food and feed. This FDA-approved technology generates short, high-intensity flashes of broad spectrum light (100–1100 nm) including UV, visible and IR radiation that destroy the nucleic acid and cell wall structure of microbes within

few seconds [42]. Eight PL flashes of  $1 \text{ J cm}^{-2}$  during 300 ms flash resulted in degradation of AFB1 in water by 92.7% [43]. In another study, PL at  $0.52 \text{ J cm}^{-1}$  per pulse was applied for 80 seconds and 15 seconds to treat rough rice and bran, respectively. It was observed that on 15 seconds of PL treatment AFB1 and AFB2 in rice bran reduced by 90.3 and 86.7% respectively, while on 80 seconds of treatment AFB1 and AFB2 in rough rice reduced by 75.0 and 39.2% respectively [44]. PL treatment also inactivated the mutagenicity and cytotoxicity of these aflatoxins. Abuagela et al. [45] treated dehulled peanuts with PL at  $0.4 \text{ J cm}^{-1}$  per pulse. No significant variation in the chemical properties including acidity value, fatty acid content and peroxide value of oil obtained from PL treated peanuts. Aflatoxins levels significantly decreased by 91% in PL treated dehulled peanuts. For large industrial scale application of PL technology would require the development of cost-effective equipment for PL treatment.

Microwave heating generates a high temperature ( $130^\circ\text{C}$  or higher) that is required for aflatoxin reduction in peanut and corn [46, 47]. Aflatoxin contaminated corn was microwave heated at 1650 W power for 5.5 minutes, resulted in reduction in AFB1, AFB2 by 36 and 58%, respectively [48]. Mobeen et al. [49] microwave cooked peanut and its products and reported reduction in AFB1 level by 50–60%, while level of AFB2 reduced to non-detectable limits. Treatment of corn flour by microwave heat for 10 minutes duration led to decrease in AFB1 content up to 67.7% [50]. Major drawback of aflatoxin decontamination by this method is the uneven distribution of temperature during microwave heating. This results in the generation of hot and cold spots in the treated food product [51]. Overheating in the hot spot may lead to loss of nutritional value and quality whereas lesser temperature in cold spot may not be sufficient for degradation of aflatoxins. In view of this, aflatoxin detoxification using microwave heating method has moderate success and limited application.

### 2.1.2 Cold plasma

The fourth state of matter, plasma predominantly consists of UV rays, ions, electrons, reactive nitrogen species (RNS), reactive oxygen species (ROS) [52]. Based on its temperature, plasma can be categorized into thermal and non-thermal (cold) plasma. Cold plasma is generated through electrical discharges in gases at temperature between  $30$  and  $60^\circ\text{C}$  and atmospheric pressure [53]. Cold plasma technology has been employed for aflatoxin degradation at ambient pressure and temperature [54, 55]. Ouf et al. [56] used argon cold plasma at atmospheric pressure for 9 minutes on *Aspergillus niger* spore and mycotoxin production in palm fruits. Authors found that mycotoxin fumonisin B2, OTA content reduced from 6 and  $25 \mu\text{g}/100 \text{ mm}^2$  respectively and all the spores were killed. Another study used dielectric barrier discharge nitrogen plasma (1150 W) for detoxification of aflatoxin inoculated dehulled hazelnuts. Authors demonstrated that after 12 minutes treatment around 70% of AFB1 was detoxified. They also showed that AFB1 and AFG1 were more sensitive whereas AFB2 and AFG2 was less sensitive to nitrogen plasma treatment [57]. Treatment with corona discharge plasma jet (CDPJ) for 30 minutes on AFB1 spiked rice and wheat samples and AFB1 on glass slides reduced the AFB1 concentration by 56.6, 45.7, 95% respectively [58]. Authors suggested the inconsistency in detoxification of AFB1 is due to possible chemical interaction of toxin with the food matrix. The effectiveness of cold plasma technology for aflatoxin degradation depends on the type of food commodities, kind of plasma system, operating parameters used (energy input, moisture, working gas) and the time of plasma exposure [53]. This novel technology has great potential for detoxification of aflatoxin in food and feed. However, it is still in its infancy and there is a need

to standardize conditions for plasma treatment suitable to decontaminate various foods. Further research is required to study the effect of plasma on the nutritional quality and organoleptic characteristics of food.

## **2.2 Chemical detoxification of aflatoxins**

### *2.2.1 Electrolyzed oxidizing water*

Electrolyzed oxidizing water (EOW) or electro-activated water is produced by the electrolysis of water containing 1% sodium chloride (NaCl). The resulting water is an electrolyzed one that can be used as a disinfectant. EOW can be categorized into two major types according to its pH and oxidation–reduction potential (ORP): a) neutral electrolyzed oxidizing water (NEOW) with pH of 5.0–6.5, ORP of 800–900 mV; b) acidic electrolyzed oxidizing water (AEOW) with pH of less than 3.0, ORP of more than 1000 mV [59]. Major advantage of EOW is that it turns to ordinary water after use that has no potential threat to animals and environment. Treatment of aflatoxin contaminated corn with NEOW at room temperature for 15 minutes, resulted in significant reduction in the genotoxicity and cytotoxicity of aflatoxins in HepG2 cells [60]. Gomez- Espinosa et al. [61] further confirmed that NEOW treatment of aflatoxin contaminated corn significantly reduces the aflatoxicosis in turkey.

### *2.2.2 Organic acids*

Organic acids have been widely used for aflatoxin degradation in food industry. AFB1 contaminated soyabean soaked in 1.0 N tartaric acid, lactic acid and citric acid, and at room temperature for 18 hours, resulted in reduction in AFB1 level by 95.1, 92.7 and 94.1%, respectively [62]. Acidulation with lemon juice for 60 minutes at 120°C for decontamination of AFB1 in roasted pistachio nuts, reduced the AFB1 content by 50.2% [63]. Acidulation can be combined with other decontamination technologies for better results. For instance, acidulation coupled with pulsed light technique led to aflatoxin degradation in peanuts up to 98.3%. On the other hand, pulsed light and citric acid treatment separately decreased aflatoxins by 78.1 and 20.2%, respectively [64]. Organic acid treatment results in leaching of nutrients such as water-soluble proteins, starch, minerals. However, it has several health benefits on livestock. The only drawback of this method is the high cost of organic acids.

### *2.2.3 Ozone*

Ozone (O<sub>3</sub>), a most powerful oxidizing, antimicrobial and disinfecting agent, can be used directly for decontaminating various food products [65]. FDA has granted ozone as Generally Recognized As Safe (GRAS) status for water and food industry [66]. Ozone cause the progressive oxidation of sulfhydryl group of amino acids of proteins, peptides and enzymes or polyunsaturated fatty acids into shorter molecular fragments. Ozone also result in degradation of unsaturated lipids in cell wall envelope, disruption and leakage of microbial cellular contents [67]. Ozone degrade aflatoxins AF-B1 and AF-G1 through an electrophilic attack on C8-C9 double bond of difuran ring, resulting in ozonide formation. This is followed by rearrangement into molozonide derivatives like organic acids, ketones and aldehydes [68]. On the contrary, AF-B2 and AF-G2 are more resistant to ozonisation as they lack C8-C9 double bond in their structure [69]. Ozone treatment of groundnut samples increased the rate of aflatoxin detoxification and has no effect

on the peroxide, resveratrol, acids and polyphenol content [70]. Treatment of AFB1- contaminated maize with ozone resulted in decreased toxicity of the treated samples [71].

Major advantages of using ozone as aflatoxin detoxification method are (i) ozone can be applied in gaseous as well as liquid forms (ii) no leftover residue after contact and no hazardous disposal (iii) easy on-site generation of ozone [65, 72]. Till date several studies have been conducted on ozonation at laboratory scale. There is a need to develop efficient equipments to scale up the process for successful commercial application of this technology for detoxification of aflatoxin contaminated food and feed.

#### 2.2.4 Plant extracts

Natural plant extracts are considered as food additives by food industry worldwide. They are well known for their anti-inflammatory and anti-microbial properties. Incubation of AFB1 with *Rosmarinus officinalis* aqueous leaf extracts at different time intervals showed 60.3% reduction in AFB1 after 48 hours of incubation [73]. Iram, *et al.* [74, 75] studied the effectiveness of aqueous extracts of *Corymbia citriodora* and *Trachyspermum ammi* for detoxification in AFB1 and AFB2 spiked corn. After treatment with *C. citriodora* leaf extract, AFB1 and AFB2 levels were reduced to 91.7 and 88.8%, respectively. AFB1 and AFB2 were degraded by 89.6 and 86.5%, respectively, following *T. ammi* seeds extract treatment. These results were in consistent with the findings of Velazhahan *et al.* [76]. In another study, authors tested the aqueous extracts from 31 medicinal plants for their AFB1 detoxifying ability. *Adhatoda vasica* Nees leaf extract showed the highest AFB1 degradation ability (98%) at 24 hours after incubation at 37°C [77]. The same group found that *Adhatoda vasica* Nees leaf extract contained the partially purified alkaloids that was responsible for its strong AFB1 detoxification ability. Brinda *et al.* [78] fed rats with spraydried formulation of *Adhatoda vasica* Nees leaf extract and exposed them to AFB1. They found that such pre-fed rats counteracted hepatotoxicity induced by AFB1 exposure. To date, there is a meager knowledge on the active compounds present in the plant extracts that is responsible for their aflatoxins detoxification ability. Further research is needed to gain insights into the aflatoxin detoxifying compounds, their mode of action and complex interaction with the food matrices.

### 2.3 Biological decontamination of aflatoxins

Biological degradation of aflatoxins involves microorganism or enzyme based detoxification of aflatoxins into less toxic or non-toxic metabolites. This method has emerged as an efficient and eco-friendly strategy for degradation of aflatoxins.

#### 2.3.1 Microbial degradation

Various microorganisms isolated from different sources can degrade aflatoxins present in food and feed. Risa *et al.* [79] investigated the effectiveness of 42 *Rhodococcus* strains for detoxification of AFB1 and zearalenone (ZON). 18 *Rhodococcus* strains were capable of degrading more than 90% of AFB1 and 15 strains could cease the genotoxicity in 72 hours. Only one of these strains, namely *R. percolatus* JCM 10087 T was capable of detoxifying ZON by more than 90% and decreasing the oestrogenicity by 70%. Another group studied the role of *R. pyridinivorans* K408 strain in detoxification of AFB1 in corn based stillage and revealed that the level of AFB1 in liquid and solid phases of whole stillage was

degraded by 75 and 63%, respectively [80]. A number of *Bacillus* strains such as *B. subtilis* ANSB060 [81], *B. subtilis* UTBSP1 [82], *B. subtilis* JSW-1 [83], *B. licheniformis* CFR1 [84], *B. licheniformis* BL010 [85], *B. velezensis* DY3108 [86] and *Bacillus* sp. TUBF1 [87] have been reported to degrade aflatoxins. *B. subtilis* has been approved as Generally Recognized As Safe (GRAS) strains of bacteria that is safe for commercial pharmaceutical and nutritional purpose.

Adebo *et al.* [88] used the lysates of bacterial strains (*Staphylococcus* sp. VGF2, *Pseudomonas fluorescens* and *Pseudomonas anguilliseptica* VGF1) isolated from gold-mine aquifer to study the degradation of AFB<sub>1</sub>. It was found that *Staphylococcus* sp. VGF2 lysate showed the highest AFB<sub>1</sub> degradation capacity of 100% while *P. fluorescens* and *P. anguilliseptica* VGF1 reduced the AFB<sub>1</sub> by 63 and 66.5%, respectively. About 124 *Streptomyces* strains were examined for AFB<sub>1</sub> degradation capability. It was found that 55% of the *Streptomyces* strains could degrade AFB<sub>1</sub> [89]. Eshelli *et al.*, [90] demonstrated that *S. lividans* TK 24 was capable of degrading AFB<sub>1</sub> by 88% in liquid culture upon 24 hour incubation. *E. coli* CG1061, AFB<sub>1</sub> detoxifying bacterium, was isolated from chicken cecum and found to degrade AFB<sub>1</sub> by 93.7% following 72 hours of incubation. It resulted in formation of degradation products of lower molecular weight [91]. The non-toxigenic strains of *Aspergillus niger* with aflatoxin detoxification capability have been used in the food industry. *A. niger* FS-UV1 strain, derived from *A. niger* FS-Z1 wild strain following UV irradiation showed superior AFB<sub>1</sub> detoxification capability up to 95.3% and significantly decreased mutagenic activity [92]. A novel thermophilic microbial consortium TADC7 was constructed for efficient, specific and stable AFB<sub>1</sub> degradation. Microbial consortium TADC7 showed the degradation of AFB<sub>1</sub> by more than 95% of the toxin after 72 hours of incubation in the temperature range between 50 and 60°C. 16S rRNA sequencing revealed that *Tepidimicrobium* and *Geobacillus* may play a major role in AFB<sub>1</sub> degradation [93].

### 2.3.2 Enzymatic degradation

Recently, the reports on isolation, identification and purification of aflatoxin-degrading enzymes from microorganisms have increased significantly. Various enzymes such as oxidases, peroxidases, reductases and laccases are capable of degrading aflatoxins. Laccases represent a class of multicopper oxidase enzymes widely present in fungi, plants, bacteria and insects. They catalyze oxidation of various phenolic and non-phenolic compounds coupled to reduction of molecular oxygen in water. Alberts *et al.* [94] first proposed the role of fungal laccases in decontamination of AFB<sub>1</sub>. Enzyme laccase (Lac2) produced and purified from *Pleurotus pulmonarius* showed AFB<sub>1</sub> degradation up to 90% with mediator acetoseringone present in the medium [95]. Verheecke *et al.* [96] documented purified enzymes effective for AFB<sub>1</sub> detoxification. Guo *et al.* [97] summarized the reaction mechanisms of aflatoxin-degrading enzymes responsible for AFB<sub>1</sub> detoxification in the review. Alberts *et al.* [94] purified laccase from *Trametes versicolor* and produced recombinant laccase using *A. niger*. The laccase from *T. versicolor* resulted in reduction in AFB<sub>1</sub> pro-oxidative properties by 59% and genotoxicity by 100% [98]. Compared to fungal laccases, bacterial laccases are excellent candidates for xenobiotics degradation because they are extremely thermostable, tolerant to alkaline conditions, wider range of pH and substrate spectrum [99]. Enzyme CotA laccase from *B. licheniformis* catalyze the C3-hydroxylation of AFB<sub>1</sub> and transformed toxic AFB<sub>1</sub> into products epi-aflatoxin Q<sub>1</sub> and aflatoxin Q<sub>1</sub> that are non-toxic human liver L-02 cells. Laccase CotA was found to be highly thermostable with the half life of 1 hour. It was capable of degrading AFB<sub>1</sub> in the temperature range between 60 and 80°C by more than 70% in a period of 30 minutes [100].

Peroxidases are a class of oxidoreductases that catalyze oxidation of substrates with the use of hydrogen peroxide or organic peroxide. They are mainly heme- proteins with contain iron (III) protoporphyrin IX as the prosthetic group. Peroxidases are widespread in nature, found in both prokaryotes and eukaryotes. Researchers have achieved 65 and 97% reduction in AFM1 and AFB1, respectively in milk with horse radish peroxidase (HRP) treatment at 30°C following 8 hours of incubation [101]. Another group used manganese peroxidase (MnP) isolated from *Phanerochaete sordida* YK-624 for AFB1 degradation. MnP reduced the AFB1 level by 86.0% and mutagenicity by 69.2% [102]. Type B dye decolorizing peroxidase (Rh\_DypB) was isolated and purified from *Rhodococcus jostii* and tested for AFB1 degradation. Authors achieved 96% degradation of AFB1 in the sodium malonate buffer [103].

Scientists are working constantly towards identification, isolation and purification of novel aflatoxin detoxifying enzymes from wide variety of organisms. For instance, an extracellular enzyme MADE was purified from *Myxococcus fulvus* ANSM068 from a culture supernatant. At the temperature of 35°C and pH 6.0, degraded AFB1 by 71.89% following 48 hours of incubation. The purified MADE also degraded AFM1 and AFG1 by 96 and 97%, respectively [104]. Xu *et al.* [105] purified enzyme MADE from *Bacillus shackletonii* that resulted in degradation of AFB1 by 48% with 72 hours of incubation at 70°C. BADE was found to be extremely thermostable, even at 100°C for 10 minutes. Nine F420H2- dependent reductases (FDRs) were identified and characterized from *Mycobacterium smegmatis*. FDRs catalyze the  $\alpha$ ,  $\beta$ -unsaturated ester reduction in aflatoxins, using deazaf-lavin as cofactor. They are categorized into two classes, namely FDR-A and FDR-B with FDR-A class of enzymes being 100 times more active against aflatoxins than FDA-B [106].

The main advantage of cell free enzyme based aflatoxin degradation is that it has no negative impact on the degradation ability and organoleptic properties of food products [88]. Although cell free aflatoxin degradation enzymes are extremely effective, however their application in food and feed industry is still limited due to certain shortcomings. Firstly, the low yield of aflatoxin-degrading enzymes in the native hosts. This can be solved by the intervention of recombinant DNA technology. Secondly, food processing generally requires the use of solvents, extreme temperature and pH conditions. This can effect the catalytic efficiency and stability of the wild type enzymes. The use of enzyme engineering technologies such as random or site directed mutagenesis could address these issues [107].

In comparison to natural enzymes, nanozymes are more robustness, cost effective, stabile and durable. Nanozymes are the nanomaterials with intrinsic enzyme like properties that catalyze the substrates of natural enzymes following the same catalytic mechanisms and kinetics under physiological conditions [108]. The laccase-mimicking nanozyme was prepared by coordinating guanosine monophosphate (GMP) with  $\text{Cu}^{2+}$  at room temperature. This led to formation of amorphous metal-organic framework (MOF) nanomaterial. Cu/GMP nanozyme has the same catalytic efficiency as the natural laccase but it is 2400-fold more cost-effective and more robust against extreme temperature, pH salt and storage conditions [109]. Nanozymes with peroxidase-like activity such as  $\text{CuMnO}_2$  nanoflakes [110],  $\text{FeMnO}_3$  nanoparticle-filled polypyrrole nanotubes [111], FePt nanoparticle-decorated graphene oxide nanosheets [112],  $\text{Pt}_{74}\text{Ag}_{26}$  nanoparticle-decorated ultrathin  $\text{MoS}_2$  nanosheets [113] have been prepared. The recent technological advancements will open the gate for the development and application of nanozymes to aflatoxin detoxification in food and feed industry.

### 3. Conclusion

Contamination caused by aflatoxins in food and feed poses a great threat to human and animal health and result in considerable economic loss to the agriculture production of the country. To produce healthy, high quality, aflatoxin free food products is the worldwide concern. Researchers have been working continuously for the development of effective aflatoxin decontamination strategies since decades. Despite all the efforts, there still a need to come up with an efficient decontamination technology that meets all these criteria for industrial commercialization: (i) Retain/enhance the nutritional quality of the food (ii) Efficiently reduce aflatoxins to the safe limits without leaving toxic residues (iii) cost effective and eco-friendly. So far, no aflatoxin decontamination strategy has been developed that meet all the mentioned criteria. For instance, physical and chemical methods may change the organoleptic properties and chemical composition of food and feed. Biological approaches are more specific, effective with more control over the generated bio-products of aflatoxin detoxification. Use of pure enzyme for aflatoxin degradation has no negative impact on the degradation efficiency, chemical composition and organoleptic properties of food products. Further, adoption of enzyme engineering technology would provide highly efficient and specific aflatoxin detoxifying enzymes in the near future.

### Acknowledgements

Author is grateful to University of Hyderabad for providing necessary facilities.

### Conflict of interest

Authors have no competing interests.

IntechOpen

### Author details

Vishakha Pandey  
University of Hyderabad, Hyderabad, India

\*Address all correspondence to: [vishakhapandey2@gmail.com](mailto:vishakhapandey2@gmail.com)

### IntechOpen

© 2021 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

## References

- [1] Reiter E, Zentek J, Razzazi E. Review on sample preparation strategies and methods used for the analysis of aflatoxins in food and feed. *Molecular Nutrition and Food Research*. 2009;53:508-524. DOI: 10.1002/mnfr.200800145.
- [2] Payne GA and Brown MP. Genetics and physiology of aflatoxin biosynthesis. *Annual Review of Phytopathology*. 1998;36:329-362. DOI: 10.1146/annurev.phyto.36.1.329.
- [3] Perrone G, Haidukowski M, Stea G, Epifani F, Bandyopadhyay R, Leslie JF, et al. Population structure and Aflatoxin production by *Aspergillus Sect. Flavi* from maize in Nigeria and Ghana. *Food Microbiology*. 2014. 41, 52e59. DOI: 10.1016/j.fm.2013.12.005.
- [4] Iqbal SZ, Jinap S, Pirouz AA and Ahmad Faizal AR . Aflatoxin M1 in milk and dairy products, occurrence and recent challenges: A review. *Trends in Food Science and Technology*. 2015;46:110e119. DOI:10.1016/J.TIFS.2015.08.005.
- [5] Blount WP. Turkey "X" disease. In the Proceeding of Turkeys. 1961;9:52-77.
- [6] Zinedine A and Manes J. Occurrence and legislation of mycotoxins in food and feed from Morocco. *Food Control*. 2009;20:334e344. DOI: 10.1016/j.foodcont.2008.07.002.
- [7] My A and Sachan D. Dietary factors affecting aflatoxin Bi carcinogenicity. *Malaysian Journal of Nutrition*. 1997;3:161-179. DOI: 10.1.1.581.8466.
- [8] Bbosa GS, Kitya D, Odda J and Ogwal-Okeng J. Aflatoxins metabolism, effects on epigenetic mechanisms and their role in carcinogenesis. *Health*. 2013; 05:14-34. DOI: 10.4236/health.2013.510A1003.
- [9] Baan R, Grosse Y, Straif K, Secretan B, El Ghissassi F, Bouvard V, Benbrahim-Tallaa L, Guha N, Freeman C, Galichet L, Cogliano V. WHO International Agency for Research on Cancer Monograph Working Group. A review of human carcinogens-Part F: chemical agents and related occupations. *Lancet Oncology*. 2009;12:1143-1144. DOI: 10.1016/S1470-2045(09)70358-4.
- [10] Roze LV, Hong SY and Linz JE. Aflatoxin biosynthesis: current frontiers. *Annual Review of Food Science and Technology*. 2013;4:293-311. DOI: 10.1146/annurev-food-083012-123702.
- [11] Kensler TW, Roebuck BD, Wogan GN and Groopman JD. Aflatoxin: A 50-year Odyssey of mechanistic and translational toxicology. *Toxicological Sciences*. 2011;120:28-48. DOI: 10.1093/toxsci/kfq283.
- [12] Marchese S, Polo A, Ariano A, Velotto S, Costantini S and Severino L. Aflatoxin B1 and M1: Biological properties and their involvement in cancer development. *Toxins*. 2018;10:214-233. DOI: 10.3390/toxins10060214.
- [13] Karlovsky P, Suman M, Berthiller F, De Meester J, Eisenbrand G, Perrin I, ... Dussort P. Impact of food processing and detoxification treatments on mycotoxin contamination. *Mycotoxin Research*. 2016;32: 179-205. DOI:10.1007/s12550-016-0257-7.
- [14] Rustom IYS. Aflatoxin in food and feed: Occurrence, legislation and inactivation by physical methods. *Food Chemistry*. 1997;59:57-67. DOI: 10.1016/S0308-8146(96)00096-9.
- [15] Vekiru E, Fruhauf S, Rodrigues I, Ottner F, Krska R, Schatzmayr G. ... Bermudez AJ. *In vitro* binding

- assessment and *in vivo* efficacy of several adsorbents against aflatoxin B1. *World Mycotoxin Journal*. 2015.;8:477-488. DOI: 10.3920/WMJ2014.1800.
- [16] Elliott CT, Connolly L and Kolawole O. Potential adverse effects on animal health and performance caused by the addition of mineral adsorbents to feeds to reduce mycotoxin exposure. *Mycotoxin Research*. 2019;36:115-126. DOI: 10.1007/s12550-019-00375-7.
- [17] Gavahian M and Cullen PJ. Cold plasma as an emerging technique for mycotoxin-free food: Efficacy, mechanisms and trends. *Food Reviews International*. 2019;36:1-22. DOI: 10.1080/87559129.2019.1630638.
- [18] Loi M, Paciolla C, Logrieco AF and Mule G. Plant bioactive compounds in pre- and postharvest management for aflatoxins reduction. *Frontiers in Microbiology*. 2020; 11:243-259. DOI:10.3389/fmicb.2020.00243.
- [19] Kumar P, Mahato DK, Kamle M, Mohanta TK, Kang SG. Aflatoxins: A Global Concern for Food Safety, Human Health and Their Management. *Frontiers in Microbiology*. 2017;7:2170-2180. DOI: 10.3389/fmicb.2016.02170.
- [20] Kim HJ, Jung S, Yong HI, Bae YS, Kang SN, Kim IS, Jo C. Improvement of Microbiological Safety and Sensorial Quality of Pork Jerky by Electron Beam Irradiation and by Addition of Onion Peel Extract and Barbecue Flavor. *Radiation Physics and Chemistry*. 2014;98:22-28. DOI: 10.1016/j.radphyschem.2014.01.003.
- [21] Rogovschi, V. D.; Aquino, S.; Nunes, T. C. F.; Gonçalves, E.; Corrêa, B. Villavicencio A L.C.H. Use Of Electron Beam On Aflatoxins Degradation In Coconut Agar. In: *Proceedings of the International Nuclear Atlantic Conference*; 2009.
- [22] Assuncao E, Reis TA, Baquiao AC, Correa B. Effects of Gamma and Electron Beam Radiation on Brazil Nuts Artificially Inoculated with *Aspergillus flavus*. *Journal of Food Protection*. 2015;78:1397-1401. DOI: 10.4315/0362-028X.JFP-14-595. PMID: 26197295.
- [23] Liu R, Lu M, Wang R, Wang S, Chang M, Jin Q and Wang X. Degradation of aflatoxin B1 in peanut meal by electron beam irradiation. *International Journal of Food Properties*. 2018;21:892-901. DOI: 10.1080/10942912.2018.1466321.
- [24] WHO. High-dose irradiation: wholesomeness of food irradiated with doses above 10 kGy. Report of a joint FAO/IAEA/WHO study group. *World Health Organization technical report series*. 1999;890:1-197.
- [25] Da Silva Aquino KA. Sterilization by gamma irradiation. In F. Adrovic (Ed.), *Gamma radiation* (pp. 171e206). Vienna, Austria: InTech. 2012. DOI: 10.5772/34901
- [26] Markov K, Mihaljevic, B, Domijan AM, Pleadin J, Delas F and Frece J. Inactivation of aflatoxigenic fungi and the reduction of aflatoxin B1 in vitro and in situ using gamma irradiation. *Food Control*. 2015;54:79-85. DOI: 10.1016/j.foodcont.2015.01.036.
- [27] Aziz NH, Moussa LA. A. Influence of gamma-radiation on mycotoxin producing moulds and mycotoxins in fruits. *Food Control*. 2002;13:281-288. DOI: 10.1016/S0956-7135(02)00028-2.
- [28] Iqbal SZ, Bhatti IA, Asi M R, Zuber M, Shahid M, Parveen I. Effect of  $\gamma$  irradiation on fungal load and aflatoxins reduction in red chillies. *Radiation Physics and Chemistry*. 2013;82:80-84. DOI: 10.1016/j.radphyschem.2012.09.015.
- [29] Kumar S, Kunwar, A, Gautam, S, Sharma, A. Inactivation of *A. ochraceus* spores and detoxification of ochratoxin A in coffee beans by gamma irradiation.

Journal of Food Sciences. 2012;77:44-51.  
DOI: 10.1111/j.1750-3841.2011.02572.x.

[30] Domijan AM, Pleadin J, Mihaljevic B, Vahcic N, Frece J, Markov K. Reduction of ochratoxin A in dry-cured meat products using gamma-irradiation. . Food Additives and Contaminants: Part A Chemistry Analysis Control Exposure and Risk Assess. 2015;32:1185-1191. DOI: 10.1080/19440049.2015.1049219.

[31] Jalili M, Jinap S and Noranizan A. Aflatoxins and ochratoxin a reduction in black and white pepper by gamma radiation. Radiation Physics and Chemistry. 2012;81:1786e1788. DOI: 10.1016/j.radphyschem.2012.06.001.

[32] Di Stefano V, Pitonzo R, Cicero N, D'Oca MC. Mycotoxin contamination of animal feedingstuff: detoxification by gamma-irradiation and reduction of aflatoxins and ochratoxin A concentrations. Food Additives and Contaminants Part A: Chemical Analysis Control Exposure and Risk Assessment. 2014;31:2034-2039. DOI: 10.1080/19440049.2014.968882.

[33] Priyadarshini A, Rajauria G, O'Donnell CP and Tiwari BK. Emerging food processing technologies and factors impacting their industrial adoption. Critical Reviews in Food Science and Nutrition. 2019; 59:3082-3101. DOI: 10.1080/10408398.2018.1483890.

[34] Caulfield CD, Cassidy JP and Kelly JP. Effects of gamma irradiation and pasteurization on the nutritive composition of commercially available animal diets. Journal of the American Association for Laboratory Animal Science. 2008;47:61-66. PMID: 19049256.

[35] Gayan E, Condon S and Alvarez I. Biological aspects in food preservation by ultraviolet light: A review. Food and Bioprocess Technology. 2014;7:1-20. DOI: 10.1007/s11947-013-1168-7.

[36] Delorme, M. M., Guimarães, J. T., Coutinho, N. M., Balthazar, C. F., Rocha, R. S., Silva, R., ... Cruz, A. G. Ultraviolet radiation: An interesting technology to preserve quality and safety of milk and dairy foods. Trends in Food Science and Technology. 2020;102:146-154. DOI: 10.1016/j.tifs.2020.06.001.

[37] Diao E, Li X, Zhang Z, Ma W, Ji N and Dong H. Ultraviolet irradiation detoxification of aflatoxins. Trends in Food Science and Technology. 2015;42:64-69. DOI: 10.1016/j.tifs.2014.12.001.

[38] Zhu Y, Koutchma T, Warrinner K, Zhou T. Reduction of patulin in apple juice products by UV light of different wavelengths in the UVC range. Journal of Food Protection. 2014;77:963-967. DOI: 10.4315/0362-028X.JFP-13-429.

[39] Atalla, MM, Hassanein NM, El-Beih AA and Youssef YA. Effect of fluorescent and UV light on mycotoxin production under different relative humidities in wheat grains. International Journal of Agriculture and Biology. 2004;6:1006-1012. DOI: 1006e1012

[40] Jubeen F, Bhatti IA, Khan MZ, Hassan ZU and Shahid M. Effect of UVC irradiation on aflatoxins in ground nut (*Arachis hypogea*) and tree nuts (*Juglans regia*, *Prunus dulcis* and *Pistachio vera*). Journal of the Chemical Society of Pakistan. 2012; 34:1366-1374. DOI: 1366e1374.

[41] Garcia-Cela E, Marin S, Sanchis V, Crespo-Sempere A and Ramos AJ. Effect of ultraviolet radiation A and B on growth and mycotoxin production by *Aspergillus carbonarius* and *Aspergillus parviticus* in grape and pistachio media. Fungal Biology. 2015;119:67e78. DOI: 10.1016/j.funbio.2014.11.004.

[42] Oms-Oliu G, Martin-Belloso O and Soliva-Fortuny R. Pulsed light

- treatments for food preservation. A review. *Food and Bioprocess Technology*. 2010;3:13-23. DOI: doi.org/10.1007/s11947-008-0147-x.
- [43] Moreau M, Lescure G, Agoulon A, Svinareff P, Orange N, Feuilloley M. Application of the pulsed light technology to mycotoxin degradation and inactivation. *J Appl Toxicol*. 2013;33:357-363. DOI: 10.1002/jat.1749.
- [44] Wang B, Mahoney NE, Pan Z, Khir R, Wu B, Ma H and Zhao L. Effectiveness of pulsed light treatment for degradation and detoxification of aflatoxin B1 and B2 in rough rice and rice bran. *Food Control*. 2016;59:461-467. DOI: 10.1016/j.foodcont.2015.06.030.
- [45] Abuagela MO, Iqdam BM, Mostafa H, Gu L, Smith ME and Sarnoski PJ. Assessing pulsed light treatment on the reduction of aflatoxins in peanuts with and without skin. *International Journal of Food Science and Technology*. 2018;53:2567-2575. DOI: 10.1111/ijfs.13851.
- [46] Pluyer HR, Ahmed EM, Wei CI. Destruction of Aflatoxins on Peanuts by Oven- and Microwave-Roasting. *Journal of Food Protection*. 1987;50:504-508. DOI: 10.4315/0362-028X-50.6.504. PMID: 30965437.
- [47] Farag RS, Rashed MM, Abo Hgger AA. Aflatoxin destruction by microwave heating. *International Journal of Food Science and Nutrition*. 1996;47:197-208. DOI: 10.3109/09637489609012581.
- [48] Perez-Flores GC, Moreno-Martinez E, Mendez-Albores A. Effect of microwave heating during alkaline-cooking of aflatoxin contaminated maize. *Journal of Food Sciences*. 2011;76:T48-T52. DOI: 10.1111/j.1750-3841.2010.01980.x.
- [49] Mobeen, AK, Aftab A, Asif A and Zuzzer AS. Aflatoxins B1 and B2 contamination of peanut and peanut products and subsequent microwave detoxification. *Journal of Pharmacy and Nutrition Sciences*. 2011;1:1-3. DOI: 10.6000/1927-5951.2011.01.01.01.
- [50] Alkadi, H and Altal J. Effect of microwave oven processing treatments on reduction of aflatoxin B1 and ochratoxin A in maize flour. *European Journal of Chemistry*. 2019;10:224-227. DOI: 10.5155/eurjchem.10.3.224-227.1840.
- [51] Menon A, Stojceska V and Tassou SA. A systematic review on the recent advances of the energy efficiency improvements in non-conventional food drying technologies. *Trends in Food Science and Technology*. 2020;100:67-76. DOI: 10.1016/j.tifs.2020.03.014.
- [52] Kogelschatz U. Atmospheric-pressure plasma technology. *Plasma Physics and Controlled Fusion*. 2004;46:B63-B75. DOI: 10.1088/0741-3335/46/12B/006.
- [53] Hertwig C, Meneses N and Mathys A. Cold atmospheric pressure plasma and low energy electron beam as alternative nonthermal decontamination technologies for dry food surfaces: A review. *Trends in Food Science and Technology*. 2018;77:131-142. DOI: 10.1016/j.tifs.2018.05.011
- [54] Gavahian M and Cullen PJ. Cold plasma as an emerging technique for mycotoxin-free food: Efficacy, mechanisms, and trends. *Food Reviews International*. 2019;36:1-22. DOI: 10.1080/87559129.2019.1630638.
- [55] Misra NN, Yadav B, Roopesh MS and Jo C. Cold plasma for effective fungal and mycotoxin control in foods: Mechanisms, inactivation effects, and applications. *Comprehensive Reviews in Food Science and Food Safety*. 2019;18:106-120. DOI: 10.1111/1541-4337.12398.

- [56] Ouf SA, Basher AH, Mohamed AA. Inhibitory effect of double atmospheric pressure argon cold plasma on spores and mycotoxin production of *Aspergillus niger* contaminating date palm fruits. *Journal of Science and Food Agriculture*. 2015;95:3204-3210. DOI: 10.1002/jsfa.7060.
- [57] Siciliano I, Spadaro D, Prella A, Vallauri D, Cavallero MC, Garibaldi A. and Gullino ML. Use of cold atmospheric plasma to detoxify hazelnuts from aflatoxins. *Toxins*. 2016;8:125. DOI: 10.3390/toxins8050125.
- [58] Puligundla P, Lee T and Mok C. Effect of corona discharge plasma jet treatment on the degradation of aflatoxin B1 on glass slides and in spiked food commodities. *Lwt-Food Science and Technology*. 2019;124:108333. DOI: 10.1016/j.lwt.2019.108333.
- [59] Huang YR., Hung YC, Hsu SY, Huang YW and Hwang DF. Application of electrolyzed water in the food industry. *Food Control*. 2008;19:329-345. DOI: 10.1016/j.foodcont.2007.08.012.
- [60] Jardon-Xicotencatl S, Diaz-Torres R, Marroquin-Cardona A, Villarreal-Barajas T, Mendez-Albores A. Detoxification of Aflatoxin-Contaminated Maize by Neutral Electrolyzed Oxidizing Water. *Toxins (Basel)*. 2015;23;7:4294-4314. DOI: 10.3390/toxins7104294.
- [61] Gomez-Espinosa D, Cervantes-Aguilar FJ, Del Río-García JC, Villarreal-Barajas T, Vazquez-Duran A, Mendez-Albores A. Ameliorative Effects of Neutral Electrolyzed Water on Growth Performance, Biochemical Constituents, and Histopathological Changes in Turkey Poults during Aflatoxicosis. *Toxins (Basel)*. 2017;9:104-119. DOI:10.3390/toxins903010.
- [62] Lee J, Her JY, Lee KG. Reduction of aflatoxins (B<sub>1</sub>, B<sub>2</sub>, G<sub>1</sub>, and G<sub>2</sub>) in soybean-based model systems. *Food Chemistry*. 2015;189:45-51. DOI: 10.1016/j.foodchem.2015.02.013.
- [63] Rastegar H, Shoeibi S, Yazdanpanah H, Amirahmadi M, Khaneghah AM, Campagnollo FB and S Sant Ana, A. Removal of aflatoxin B1 by roasting with lemon juice and/or citric acid in contaminated pistachio nuts. *Food Control*. 2017; 71:279-284. DOI: 10.1016/j.foodcont.2016.06.045.
- [64] Abuagela MO, Iqdam BM, Mostafa H, Marshall SM, Yagiz Y, Marshall MR, ... Sarnoski P. Combined effects of citric acid and pulsed light treatments to degrade B-aflatoxins in peanut. *Food and Bioprocess Processing*. 2019; 117:396-403. ISSN : 0960-3085.
- [65] Pandiselvam R, Subhashini, S, Banuu Priya EP, Kothakota A, Ramesh SV and Shahir S. Ozone based food preservation: A promising green technology for enhanced food safety. *Ozone: Science & Engineering*. 2019;41:17-34. DOI: 10.1080/01919512.2018.1490636.
- [66] Food and Drug Administration (FDA). Secondary direct food additives permitted in food for human consumption. *Federal Register*. 2001;66:33829-33830.
- [67] Das E, Gurakan GC and Bayindirli A. Effect of controlled atmosphere storage, modified atmosphere packaging and gaseous ozone treatment on survival of *Salmonella Enteridis* on cherry tomatoes. *Food Microbiology*. 2006;23:430e438. DOI: 10.1016/j.fm.2005.08.002.
- [68] Jalili, M. A review on aflatoxins reduction in food. *Iranian Journal of Health, Safety and Environment*. 2016;3:445-459.
- [69] Agriopoulou S, Koliadima A, Karaiskakis G and Kapolos J. Kinetic

- study of aflatoxins' degradation in the presence of ozone. *Food Control*. 2016;61:221e226. DOI: 10.1016/j.foodcont.2015.09.013.
- [70] Chen R, Ma F, Li PW, Zhang W, Ding XX, Zhang Q, Li M, Wang YR, Xu BC. Effect of ozone on aflatoxins detoxification and nutritional quality of peanuts. *Food Chemistry*. 2014;146:284-288. DOI: 10.1016/j.foodchem.2013.09.059.
- [71] Luo X, Wang R, Wang L, Li Y, Bian Y and Chen Z. Effect of ozone treatment on aflatoxin B1 and safety evaluation of ozonized corn. *Food Control*. 2014;37:171e176. DOI: 10.1016/j.foodcont.2013.09.043.
- [72] Torres AM, Palacios SA, Yerkovich N, Palazzini JM, Battilani P, Leslie JF, ... Chulze SN. *Fusarium* head blight and mycotoxins in wheat: Prevention and control strategies across the food chain. *World Mycotoxin Journal*. 2019; 12:333-355. DOI: 10.3920/WMJ2019.2438.
- [73] Ponzilacqua B, Rottinghaus GE, Landers BR and Oliveira CAF. Effects of medicinal herb and Brazilian traditional plant extracts on in vitro mycotoxin decontamination. *Food Control*. 2019;100:24-27. DOI: 10.1016/j.foodcont.2019.01.009.
- [74] Iram W, Anjum T, Iqbal M, Ghaffar A and Abbas M. Mass spectrometric identification and toxicity assessment of degraded products of aflatoxin B1 and B2 by *Corymbia citriodora* aqueous extracts. *Scientific Reports*. 2015, 5:14672-14687. DOI: doi.org/10.1038/srep14672.
- [75] Iram W, Anjum T, Iqbal M, Ghaffar A, Abbas M. Structural Elucidation and Toxicity Assessment of Degraded Products of Aflatoxin B1 and B2 by Aqueous Extracts of *Trachyspermum ammi*. *Frontiers in Microbiology*. 2016;7:346-362. DOI: 10.3389/fmicb.2016.00346.
- [76] Velazhahan R, Vijayanandraj S, Vijayasamundeeswari A, Paranidharan V, Samiyappan R, Iwamoto T, ... Muthukrishnan, S. Detoxification of aflatoxins by seed extracts of the medicinal plant, *Trachyspermum ammi* (L.) Sprague ex Turrill - Structural analysis and biological toxicity of degradation product of aflatoxin G1. *Food Control*. 2010;21:719-725. DOI: 10.1016/j.foodcont.2009.10.014.
- [77] Vijayanandraj S, Brinda R, Kannan K, Adhithya R, Vinothini S, Senthil K, Chinta RR, Paranidharan V, Velazhahan R. Detoxification of aflatoxin B1 by an aqueous extract from leaves of *Adhatoda vasica* Nees. *Microbiological Research*. 2014;169:294-300. DOI: 10.1016/j.micres.2013.07.008.
- [78] Brinda R, Vijayanandraj S, Uma D, Malathi D, Paranidharan V, Velazhahan R. Role of *Adhatoda vasica* (L.) Nees leaf extract in the prevention of aflatoxin-induced toxicity in Wistar rats. *Journal of the Sciences of Food and Agriculture*. 2013;93:2743-2748. DOI: 10.1002/jsfa.6093.
- [79] Risa A, Krifaton C, Kukolya J, Kriszt B, Cserhati M, Tancsics A. Aflatoxin B1 and Zearalenone-Detoxifying Profile of *Rhodococcus* Type Strains. *Current Microbiology*. 2018;75:907-917. DOI: 10.1007/s00284-018-1465-5.
- [80] Prettl Z, Desi E, Lepossa A, Kriszt B, Kukolya J and Nagy E. Biological degradation of aflatoxin B1 by a *Rhodococcus pyridinivorans* strain in by-product of bioethanol. *Animal Feed Science and Technology*. 2017;224:104-114. DOI: 10.1016/j.anifeedsci.2016.12.011.
- [81] Gao X, Ma Q, Zhao L, Lei Y, Shan Y and Ji C. Isolation of *Bacillus subtilis*: Screening for aflatoxins B1, M1, and G1 detoxification. *European Food Research and Technology*. 2011;232:957-962. DOI: 10.1007/s00217-011-1463-3

- [82] Farzaneh M, Shi Z-Q, Ghassempour A, Sedaghat N, Ahmadzadeh M, Mirabolfathy M and Javan-Nikkhah M. Aflatoxin B<sub>1</sub> degradation by *Bacillus subtilis* UTBSP1 isolated from pistachio nuts of Iran. *Food Control*. 2012;23:100-106 DOI: 10.1016/j.foodcont.2011.06.018.
- [83] Xia X, Zhang Y, Li M, Garba B, Zhang Q, Wang Y, ... Li P. Isolation and characterization of a *Bacillus subtilis* strain with aflatoxin B<sub>1</sub> biodegradation capability. *Food Control*. 2017;75:92-98. DOI: 10.1016/j.foodcont.2016.12.036.
- [84] Raksha Rao K, Vipin AV, Hariprasad P, Anu Appaiah KA and Venkateswaran G. Biological detoxification of aflatoxin B<sub>1</sub> by *Bacillus licheniformis* CFR1. *Food Control*. 2017;71:234-241. DOI: 10.1016/j.foodcont.2016.06.040
- [85] Wang Y, Zhang H, Yan H, ... Zhang Z. Effective Biodegradation of Aflatoxin B<sub>1</sub> Using the *Bacillus licheniformis* (BL010) Strain. *Toxins (Basel)*. 2018;10:497-513. DOI:10.3390/toxins10120497.
- [86] Shu X, Wang Y, Zhou Q, ... Wu L.. Biological Degradation of Aflatoxin B<sub>1</sub> by Cell-Free Extracts of *Bacillus velezensis* DY3108 with Broad PH Stability and Excellent Thermostability. *Toxins (Basel)*. 2018;10:330-345. DOI:10.3390/toxins10080330
- [87] El-Deeb B, Altalhi A, Khiralla G, Hassan S and Gherbawy Y. Isolation and characterization of endophytic *Bacilli* bacterium from maize grains able to detoxify aflatoxin B<sub>1</sub>. *Food Biotechnology*. 2013;27:199-212. DOI: 10.1080/08905436.2013.811083.
- [88] Adebo OA, Njobeh PB, Sidu S, Tlou MG, Mavumengwana V. Aflatoxin B<sub>1</sub> degradation by liquid cultures and lysates of three bacterial strains. *International Journal of Food Microbiology*. 2016;233:11-19. DOI: 10.1016/j.ijfoodmicro.2016.06.007.
- [89] Harkai P, Szabo I, Cserhati M, Krifaton C, Risa A, Rado J, ... Kriszt B. Biodegradation of aflatoxin-B<sub>1</sub> and zearalenone by *Streptomyces* sp. collection. *International Biodeterioration and Biodegradation*. 2016;108:48-56. DOI: 10.1016/j.ibiod.2015.12.007.
- [90] Eshelli M, Harvey L, Edrada-Ebel R, McNeil B. Metabolomics of the biodegradation process of aflatoxin B<sub>1</sub> by actinomycetes at an initial pH of 6.0. *Toxins (Basel)*. 2015;7:439-456. DOI: 10.3390/toxins7020439
- [91] Wang L, Wu J, Liu Z, Shi Y, Liu J, Xu X, Hao S, Mu P, Deng F, Deng Y. Aflatoxin B<sub>1</sub> Degradation and Detoxification by *Escherichia coli* CG1061 Isolated From Chicken Cecum. *Frontiers in Pharmacology*. 2019;9:1548-1557. doi: 10.3389/fphar.2018.01548.
- [92] Sun X, Sun C, Zhang X, Zhang H, Ji J, Liu Y and Tang L. Aflatoxin B<sub>1</sub> decontamination by UV-mutated live and immobilized *Aspergillus niger*. *Food Control*. 2016;61:235-242. DOI: 10.1016/j.foodcont.2015.09.017.
- [93] Wang Y, Zhao C, Zhang D, Zhao M, Zheng D, Lyu Y, Cheng W, Guo P, Cui Z. Effective degradation of aflatoxin B<sub>1</sub> using a novel thermophilic microbial consortium TADC7. *Bioresource Technology*. 2017;224:166-173. DOI: 10.1016/j.biortech.2016.11.033.
- [94] Alberts JF, Gelderblom WC, Botha A and Van Zyl WH. Degradation of aflatoxin B<sub>1</sub> by fungal laccase enzymes. *International Journal of Food Microbiology*. 2009; 135:47-52. DOI: 10.1016/j.ijfoodmicro.2009.07.022.
- [95] Loi M, Fanelli F, Zucca P, ... Mule G. Aflatoxin B<sub>1</sub> and M<sub>1</sub> Degradation by Lac2 from *Pleurotus pulmonarius* and Redox Mediators. *Toxins (Basel)*. 2016;8:245. DOI:10.3390/toxins8090245
- [96] Verheecke C, Liboz T, Mathieu F. Microbial degradation of aflatoxin B<sub>1</sub>:

Current status and future advances. International Journal of Food Microbiology. 2016;237:1-9. DOI: 10.1016/j.ijfoodmicro.2016.07.028.

[97] Guo Y, Zhao L, Ma Q, Ji C. Novel strategies for degradation of aflatoxins in food and feed: A review. Food Research International. 2020;109878: ISSN 0963-9969. DOI: 10.1016/j.foodres.2020.109878.

[98] Zeinvand-Lorestani H, Sabzevari O, Setayesh N, Amini M, Nili-Ahmadabadi A, Faramarzi MA. Comparative study of in vitro prooxidative properties and genotoxicity induced by aflatoxin B1 and its laccase-mediated detoxification products. Chemosphere. 2015;135:1-6. DOI: 10.1016/j.chemosphere.2015.03.036.

[99] Guan ZB, Luo Q, Wang HR, Chen Y, Liao XR. Bacterial laccases: Promising biological green tools for industrial applications. Cellular and Molecular Life Sciences. 2018;75:3569-3592. DOI: 10.1007/s00018-018-2883-z.

[100] Guo Y, Qin X, Tang Y, Ma Q, Zhang J and Zhao, L. *cota* laccase, a novel aflatoxin oxidase from *Bacillus licheniformis*, transforms aflatoxin B1 to aflatoxin Q1 and epi-aflatoxin Q1. Food Chemistry. 2020;325:Article 126877. DOI: 10.1016/j.foodchem.2020.126877.

[101] Marimon Sibaja, KV, de Oliveira Garcia, S, Feltrin ACP, Diaz Remedi R, Cerqueira MBR, Badiale-Furlong E and Garda-Buffon, J. Aflatoxin biotransformation by commercial peroxidase and its application in contaminated food. Journal of Chemical Technology and Biotechnology. 2018;94:1187-1194. DOI: 10.1002/jctb.5865.

[102] Wang J, Ogata M, Hirai H, Kawagishi H. Detoxification of aflatoxin B1 by manganese peroxidase from the white-rot fungus *Phanerochaete sordida*

YK-624. FEMS Microbiology Letters. 2011;314:164-169. doi: 10.1111/j.1574-6968.2010.02158.x.

[103] Loi M, Renaud JB, Rosini E, Pollegioni L, Vignali E, Haidukowski M, Sumarah MW, Logrieco AF, Mulè G. Enzymatic transformation of aflatoxin B<sub>1</sub> by Rh\_DypB peroxidase and characterization of the reaction products. Chemosphere. 2020;250:126296. DOI: 10.1016/j.chemosphere.2020.126296.

[104] Zhao LH, Guan S, Gao X, Ma QG, Lei YP, Bai XM, Ji C. Preparation, purification and characteristics of an aflatoxin degradation enzyme from *Myxococcus fulvus* ANSM068. Journal of Applied Microbiology. 2011;110:147-155. doi: 10.1111/j.1365-2672.2010.04867.x.

[105] Xu L, Eisa Ahmed MF, Sangare L, Zhao Y, Selvaraj JN, Xing F, Wang Y, Yang H, Liu Y. Novel Aflatoxin-Degrading Enzyme from *Bacillus shackletonii* L7. Toxins (Basel). 2017;9:36. DOI: 10.3390/toxins9010036.

[106] Lapalikar GV, Taylor MC, Warden AC, Scott C, Russell RJ, Oakeshott JG. F420H2-dependent degradation of aflatoxin and other furanocoumarins is widespread throughout the actinomycetales. PLoS One. 2012;7:e30114. DOI: 10.1371/journal.pone.0030114.

[107] Zhang Y, Geary T and Simpson BK. Genetically modified food enzymes: A review. Current Opinion in Food Science. 2019;25:14-18. DOI: 10.1016/j.cofs.2019.01.002.

[108] Wei H, Wang E. Nanomaterials with enzyme-like characteristics (nanozymes): next-generation artificial enzymes. Chemical Society Review. 2013;42:6060-6093. DOI: 10.1039/c3cs35486e.

[109] Liang H, Lin F, Zhang Z, Liu B, Jiang S, Yuan Q, Liu J. Multicopper

Laccase Mimicking Nanozymes with Nucleotides as Ligands. *ACS Applied Materials and Interfaces*. 2017;9:1352-1360. DOI: 10.1021/acsami.6b15124

[110] Chen M, Yang B, Zhu J, Liu H, Zhang X, Zheng X, Liu Q. FePt nanoparticles-decorated graphene oxide nanosheets as enhanced peroxidase mimics for sensitive response to H<sub>2</sub>O<sub>2</sub>. *Material Science and Engineering C Materials for Biological Applications*. 2018;90:610-620. DOI: 10.1016/j.msec.2018.05.004.

[111] Chi MQ, Chen SH, Zhong MX, Wang C and Lu XF. Self-templated fabrication of FeMnO<sub>3</sub> nanoparticle-filled polypyrrole nanotubes for peroxidase mimicking with a synergistic effect and their sensitive colorimetric detection of glutathione. *Chemical Communications*. 2018;54:5827-5830. DOI:10.1039/C8CC01574K.

[112] Chen M, Yang B, Zhu J, Liu H, Zhang X, Zheng X, Liu Q. FePt nanoparticles-decorated graphene oxide nanosheets as enhanced peroxidase mimics for sensitive response to H<sub>2</sub>O<sub>2</sub>. *Materials Science & Engineering C-Materials for Biological Applications*. 2018;90:610-620. DOI: 10.1016/j.msec.2018.05.004

[113] Cai S, Han Q, Qi C, Lian Z, Jia X, Yang R, Wang C. Pt<sub>74</sub>Ag<sub>26</sub> nanoparticle-decorated ultrathin MoS<sub>2</sub> nanosheets as novel peroxidase mimics for highly selective colorimetric detection of H<sub>2</sub>O<sub>2</sub> and glucose. *Nanoscale*. 2016;8:3685-3693. DOI: 10.1039/c5nr08038j.

[114] Patil H, Shah NG, Hajare SN, Gautam S and Kumar G. Combination of microwave and gamma irradiation for reduction of aflatoxin B<sub>1</sub> and microbiological contamination in peanuts (*Arachis hypogaea L.*). *World Mycotoxin Journal*. 2019;12:269-280. DOI: 10.3920/WMJ2018.2384.

[115] Vita DS, Rosa P and Giuseppe A. Effect of gamma irradiation on aflatoxins and ochratoxin A reduction in almond samples. *Journal of Food Research*. 2014;3:113-118. DOI: 10.5539/jfr.v3n4p113.

[116] Garg N, Aggarwal M, Javed S and Khandal RK. Studies for optimization of conditions for reducing aflatoxin contamination in peanuts using ultraviolet radiations. *International Journal of Drug Development and Research*. 2013; 5: 408-424.

[117] Mao J, He B, Zhang L,... Zhang W. A Structure Identification and Toxicity Assessment of the Degradation Products of Aflatoxin B<sub>1</sub> in Peanut Oil under UV Irradiation. *Toxins (Basel)*. 2016;8:332. DOI: 10.3390/toxins8110332.

[118] Diao E, Shen X, Zhang Z, Ji N, Ma W and Dong H. Safety evaluation of aflatoxin B<sub>1</sub> in peanut oil after ultraviolet irradiation detoxification in a photodegradation reactor. *International Journal of Food Science & Technology*. 2015;50:41-47. DOI: 10.1111/ijfs.12648.

[119] Liu R, Jin Q, Huang J, Liu Y, Wang X, Mao W and Wang S. Photodegradation of aflatoxin B<sub>1</sub> in peanut oil. *European Food Research and Technology*. 2011;232: 843-849. DOI: 10.1007/s00217-011-1452-6.

[120] Tripathi S, Mishra HN. Enzymatic coupled with UV degradation of aflatoxin B<sub>1</sub> in red chili powder *Journal of Food Quality*. 2010;33:186-203. DOI: 10.1111/j.1745-4557.2010.00334.x.