

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,900

Open access books available

124,000

International authors and editors

140M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



## Simulating peatland methane dynamics coupled to a mechanistic model of biogeochemistry, hydrology, and energy: Implications to climate change

Takeshi Ise<sup>1,5</sup>, Allison L. Dunn<sup>2,6</sup>,

Steven C. Wofsy<sup>3,7</sup> and Paul R. Moorcroft<sup>4,8</sup>

<sup>1</sup> Japan Agency for Marine-Earth Science and Technology, 3173-25 Showa-machi, Kanazawa-ku, Yokohama, Kanagawa, 236-0001, Japan

<sup>2</sup> Department of Physical and Earth Sciences, Worcester State College, 486 Chandler Street, Worcester, MA 01602, U.S.A.

<sup>3</sup> Department of Earth and Planetary Sciences, Harvard University, 20 Oxford Street, Cambridge, MA 02138, U.S.A.

<sup>4</sup> Department of Organismic and Evolutionary Biology, Harvard University, 22 Divinity Avenue, Cambridge, MA 02138 U.S.A.

<sup>5</sup> Correspondence author. [ise@jamstec.go.jp](mailto:ise@jamstec.go.jp), phone: +81-45-778-5595, fax: +81-45-778-5706, Mailing address: 3173-25 Showa-machi, Kanazawa-ku, Yokohama, Kanagawa, 236-0001, Japan

### Abstract

Northern peat lands have a strong potential to modify climate through changes in soil organic carbon (SOC) and methane (CH<sub>4</sub>). A dynamic interaction among climate, soil physical properties (e.g., temperature and moisture), and biogeochemistry (e.g., quantity and quality of SOC) determines the peat land system. Due to this interaction, CH<sub>4</sub> production, oxidation, and transport dynamics changes dramatically under climate change. To appropriately study the future CH<sub>4</sub> in a predictive manner, a simulation model must be able to reproduce the inherent dynamic interaction in the peat land system.

Here, these complex interactions were simulated simultaneously in a biogeochemical peat land model coupled with mechanistic soil hydrology and thermal dynamics (ED2.0-peat). The model successfully reproduced soil physical profiles and the resultant SOC and CH<sub>4</sub> observed in a poor fen of northern Manitoba. With an experimental simulation of 4°C warming, a significant long-term decline in CH<sub>4</sub> emission was found, caused by a loss in substrate and prevalence of aerobic conditions. However, there was a transient increase in CH<sub>4</sub> emission shortly after warming because of time lag between the temperature dependence of microbial activity (a fast response to climate change) and the loss in peat depth (a slow response).

## 1. Introduction

High latitudinal regions are predicted to undergo particularly intensive warming (Holland and Bitz, 2003), with SOC in northern peat lands likely to be strongly affected (Ise et al., 2008). Peat decomposition has a potential to induce a strong biogeochemical feedback to the global climate system by adding excess CO<sub>2</sub> to the atmosphere (Friedlingstein et al., 2006). In addition, northern peat lands are a large source of CH<sub>4</sub>, (~10%; Schlesinger, 1997; Frohking et al., 2006), another major greenhouse gas. Developing a process-based understanding of peat land dynamics therefore is critical to predicting peat land responses to climate and feedbacks onto global climate.

Soil thermal and moisture conditions strongly controls wetland CH<sub>4</sub> dynamics, in addition to SOC accumulation and decomposition (Pelletier et al., 2007). The position of water table is the essential determinant of peat land CH<sub>4</sub> dynamics because CH<sub>4</sub> production by methanogens is a strictly anaerobic process, and CH<sub>4</sub> oxidation by methanotrophs consumes CH<sub>4</sub> in aerobic layers (Fig. 1). In addition, those biological processes are dependent on temperature of each layer. To accurately simulate CH<sub>4</sub> dynamics of a peat land, appropriate reproduction of short-term (sub-daily), depth-structured soil physics (i.e., hydrology and energy) is needed.

Moreover, in long-term, changes in peat quantity and quality that alter substrate conditions should be appropriately represented. Biogeochemical and physical dynamics in organic soils are markedly different from those in mineral soils. In particular, the amount of SOC stored in peat affects its physical properties, with peat surfaces rising and falling as SOC changes. During periods of SOC accumulation, the area becomes moister as humified peat generally has a higher water holding capacity and a lower hydraulic conductivity than mineral soils (Clymo, 1984). This causes the water table to rise, and the resultant anaerobic condition slows SOC decomposition, leading to further accumulation (positive feedback). The rate of water table rise does not keep up with the peat growth rate indefinitely, however, and thus this eventually acts as a negative feedback on further SOC accumulation (the proportion of SOC below water table becomes smaller through time, and the resultant higher decomposition rate restricts peat growth; Ise et al., 2008). Water table depth, which is a critical factor regulating decomposition rates, should be simulated from hydrological changes in the physical characteristics of peat, thereby capturing those feedbacks on SOC accumulation and decomposition. To our knowledge, only ED2.0-peat (Ise et al., 2008) has examined those processes simultaneously and mechanistically. Thus the mechanistic calculation of soil energy and moisture by ED2.0-peat is advantageous to study the future trend of CH<sub>4</sub> in a predictive manner, under various environmental conditions.

In this paper, we introduce CH<sub>4</sub> production, oxidation, and transportation sub models into ED2.0-peat—a peat carbon model with dynamic hydrological and thermal processes—to simulate peat land CH<sub>4</sub> dynamics in a predictive manner, together with dynamically-changing soil conditions. The peat biogeochemical model is coupled with a soil physics model, allowing peat depth to change with SOC, and simulate the resultant changes in soil properties. As a continuation from Ise et al. (2008), we parameterize the model with field observations from a poor fen in Manitoba, Canada and evaluate the model with the observed SOC, CH<sub>4</sub> emission rate, soil temperature, and water table depth. Using an experimental warming, responses of peat SOC to a warming climate are then studied. It should be noted that the explicit reproduction of the dynamic peat land system (i.e., vertical

profiles of temperature, moisture, and SOC) is the prerequisite for the predictive study of CH<sub>4</sub> dynamics, and we describe the model structure and outputs of ED2.0-peat accordingly.

## 2. Methods

ED2.0-peat was applied to the Northern Study Area (NSA) Fen site of the Boreal Ecosystem and Atmosphere Study (BOREAS; Sellers et al., 1995; Trumbore and Harden, 1997) in northern Manitoba, Canada. This study area is close to the northern limit of boreal forest, and in the discontinuous permafrost zone (Veldhuis et al., 2002). The area is periodically covered by standing water especially in spring.

The site-specific parameters were obtained and summarized from the BOREAS dataset. The average bulk density of fibrous peat is 0.092 gSoil/cm<sup>3</sup>. The average bulk density of humic peat (0.38 gSoil/cm<sup>3</sup>) is taken from the modal estimate for Oh horizons in BOREAS Northern Study Area, among the range of 0.13 gSoil/cm<sup>3</sup>. The carbon fraction of fibrous peat is 0.4 gC/gSoil, and that of humic peat is 0.3 gC/gSoil (Apps and Halliwell, 1999). Litter input rate to fibrous layer was estimated from the net primary productivity (NPP) reported by Bond-Lamberty et al. (2004; 0.26 kgC m<sup>-2</sup> y<sup>-1</sup>) and partitioned into labile (0.17 kgC m<sup>-2</sup> y<sup>-1</sup>) and recalcitrant (0.09 kgC m<sup>-2</sup> y<sup>-1</sup>) fractions according to a relationship between nitrogen density and lignin fraction of plant biomass described by Parton et al. (1987), using a generalized lignin fraction of boreal forest species including bryophytes calculated by Ise and Moorcroft (2006). Meteorological data (1994-2005) from the eddy covariance tower at the Old Black Spruce site (OBS; 7 km southwest of the Fen site; Dunn et al., 2007) were used to drive the simulation with a 30-minute time step. Missing meteorological data (<10%) were filled with data from the closest weather station at Thompson Airport, Manitoba (39 km east of the site).

ED2.0-peat model structure and data acquisition are described in detail by Ise et al. (2008). To evaluate the model performance, ED2.0-peat was first driven by the meteorological data of 1994, the year of the BOREAS field campaign for soil thermal and hydrological observation at the Fen site. Mean water table depth, soil temperature, and CH<sub>4</sub> emission data of 4 observation subsides at Fen site for year 1994 were calculated to evaluate the model. 12 years of meteorological data (1994-2005) from the OBS flux tower were then used repeatedly to drive long-term simulations (see Supplementary Information in Ise et al. (2008) for the 12-year variability in simulation). To obtain insights about possible effects of future climate change, transient responses of the peat land system under the climate change and SOC sensitivities under a new climate regime were studied. We simulated CH<sub>4</sub> production, oxidation, transport, and emission to the atmosphere at NSA Fen site based on Walter and Heimann (2000) and parameterization was identical to their application to a poor fen in Michigan unless otherwise noted below.

The dynamic peat depth is characteristic to ED2.0-peat, and the CH<sub>4</sub> dynamics simulation is applied to each peat layer separately (Fig. 1). The position of water table determines the ratio between CH<sub>4</sub> production and oxidation. CH<sub>4</sub> production rates are different among substrates (fibrous and humid peat). The CH<sub>4</sub> production rate of the fibrous layer  $R_{prod.f}$  [ $\mu\text{mol m}^{-2} \text{h}^{-1}$ ] is:

$$R_{prod.f} = R_0 \cdot f_{labile.f} \cdot f_{in} \cdot TD_f \cdot 1000 \cdot z_f \cdot f_{anox.f} \quad (1)$$

where  $TD_f$  is the temperature dependence factor of the fibrous layer calculated as equation 5 in Walter and Heimann (2000), with mean soil temperature of that layer simulated by ED2.0-peat.  $f_{labile-f}$  is the relative liability of fibrous layer (1.0) based on Ise and Moorcroft (2006).  $R_0$  is the  $CH_4$  production rate [ $\mu M h^{-1}$ ] at the reference temperature, as described in Walter and Heimann (2000). The depth of the fibrous layer  $z_f$  and the anoxic fraction of that layer  $f_{anox-f}$  are obtained from ED2.0-peat (Ise et al., 2008). Similarly,  $CH_4$  production from humid layer is:

$$R_{prod-h} = R_0 \cdot f_{labile-h} \cdot f_{in} \cdot TD_h \cdot 1000 \cdot z_h \cdot f_{anox-h} \quad (2)$$

$TD_h$ ,  $z_h$ , and  $f_{anox-h}$  are based on ED2.0-peat variables.  $f_{labile-f}$  is the relative liability of humid layer (0.044) based on Ise and Moorcroft (2006). Then,  $CH_4$  oxidation rates of the fibrous layer  $R_{oxid-f}$  [ $\mu mol m^{-2} h^{-1}$ ] and the humid layer  $R_{oxid-h}$  [ $\mu mol m^{-2} h^{-1}$ ] is Michaelis-Menten type functions of  $CH_4$  concentration of each layer  $[CH_4]_f$  and  $[CH_4]_h$ , respectively:

$$R_{oxid-f} = V_{max} \cdot \frac{[CH_4]_f}{k_m + [CH_4]_f} \cdot TD_f \cdot 1000 \cdot z_f \cdot (1 - f_{anox-f}) \quad (3)$$

$$R_{oxid-h} = V_{max} \cdot \frac{[CH_4]_h}{k_m + [CH_4]_h} \cdot TD_h \cdot 1000 \cdot z_h \cdot (1 - f_{anox-h}) \quad (4)$$

where  $V_{max}$  and  $K_m$  are oxidation parameters described by Walter and Heimann (2000). Three modes of  $CH_4$  transport (diffusion, bubbling, and vascular transport) were explicitly modeled. We assume that the plant rooting depth is at the boundary between fibrous and humid peat and root distribution is uniform. Thus, when the water table is above this boundary, the fast  $CH_4$  transportation through ebullition (bubbling) occurs as described in Walter and Heimann (2000).




Fig. 1. Schematic diagram of ED2.0-peat, when water table is in fibrous layer (a) or in humid layer (b). Position of water table determines proportions of aerobic vs. anaerobic peat depth in each layer.  $CH_4$  production occurs in anaerobic layers, and  $CH_4$  oxidation occurs in aerobic layers. Three modes of  $CH_4$  transportation are also shown. Diffusion is a slow, continuous process driven by  $CH_4$  concentration gradient. Ebullition occurs when  $CH_4$  concentration of an anaerobic layer exceeds a threshold (see section 2). When water table is above rooting depth,  $CH_4$  is also transported through plant vascular system (see section 2).

To evaluate the model performance, we compared the simulation results by ED2.0-peat against field-observed data of water table depth, soil temperature at 10 cm below peat surface, and CH<sub>4</sub> emission rate from BOREAS NSA Fen site for 1994. Observed variables are gap-filled composites from 4 observation subsides and 32 measurement collars located in hummock, hollow, lawn, and moat. Gaps in the data were filled by linear interpolation. Each measurement time series are weighted equally to estimate mean and variance. Residuals were assumed to be normally distributed to generate 95-percent confidence intervals.

### 3. Results

The model reasonably reproduced water table and soil temperature dynamics observed at the site for 1994 (Fig. 2). ED2.0-peat generally captured seasonal patterns of soil thermal and moisture conditions, and the simulated trends were generally within the range of intra-site variations. There were some notable, systematic underestimate in the simulation of soil temperature in early summer because ED2.0-peat lacked treatment of lateral water flow that can bring a large amount of warm surface water into the site, which is typical for fens especially after soil melt. CH<sub>4</sub> emission rates were influenced by soil temperature and water table position and showed a similar seasonal pattern with these soil physical properties. The simulated CH<sub>4</sub> dynamics was generally consistent with the field observation. The SOC accumulation under the current climate was 116 kgC/m<sup>2</sup>, (Ise et al., 2008), within the estimate range of 79-120 kgC/m<sup>2</sup> based on field studies (Trumbore and Harden, 1997).

Fen SOC was very sensitive to climate change. An experimental warming of 4°C caused an 86% loss of SOC (Fig. 3), because of the higher sensitivity caused by a positive feedback process between water loss and decomposition; an acceleration in SOC decomposition caused by the warming lowered the water table, resulting in further decomposition due to enhanced aerobic conditions (Fig. 3a and b). A reduction in summer-time soil insulation also accelerated decomposition (see Ise et al., 2008). Because of the radical changes in soil depth and hydrological and thermal environments, CH<sub>4</sub> emission responded to the climate change sensitively and showed a complex trajectory (Fig. 4). Due to the drier conditions caused by the climate change and loss in substrate, the equilibrium CH<sub>4</sub> emission rate under the warmer climate regime was significantly smaller than that under the current climate. However, the warming caused a transient increase in CH<sub>4</sub> emission due to the temperature dependency of methanogenic microorganisms (fast response: dominating in simulation years 2000-2100). Then, however, the SOC exposed to warmer and drier climate started to decompose and the reduction in soil depth and water holding capacity eventually decreased CH<sub>4</sub> emission (slow response). During the transition period after the climate change (simulation years 2200-2700), the water table was periodically in humid layer (Fig. 3b), and this caused dramatic reduction of CH<sub>4</sub> emission. After simulation year 2700, the new equilibrium of peat system was established and the mean position of water table is again in the fibrous layer, and thus the CH<sub>4</sub> emission rates were partially recovered.




Fig. 2. (a) Water table depth, (b) soil temperature [ $^{\circ}\text{C}$ ] at 10 cm below peat surface, and (c)  $\text{CH}_4$  emission dynamics for BOREAS Fen site in 1994. Observed variables are gap-filled composite datasets from 4 observation subsides and 32 measurement collars located in hummock, hollow, lawn, and moat. Each measurement time series are weighted equally to estimate mean and variance. Residuals were assumed to be normally distributed to generate 95-percent confidence intervals (grey shades).




Fig. 3. 4000-year simulation of 12-year means of fraction below water table, (a) fibrous and (b) humid layers and (c) changes in peat depth. This simulation is applied to BOREAS Northern Study Area Fen site. 1994-2005 meteorological data is used repeatedly for this 4000-year simulation. A uniform rise of temperature by 4°C is applied after simulation year 2000.




Fig. 4. 4000-year simulation of CH<sub>4</sub> emission from BOREAS Northern Study Area Fen site. 1994-2005 meteorological data is used repeatedly for this 4000-year simulation. A uniform rise of temperature by 4°C is applied after simulation year 2000.

#### 4. Discussion

To our knowledge, this study is the first attempt to simulate  $\text{CH}_4$  dynamics with a physics-based, fast-timescale model of soil temperature and moisture coupled to a long-term, biogeochemical model. ED2.0-peat captured the essential physical dynamics of BOREAS NSA Fen site. The short-term  $\text{CH}_4$  dynamics with environmental dependency was reasonably reproduced. In contrast to the preexisting  $\text{CH}_4$  models (e.g., Walter and Heimann, 2000), ED2.0-peat does not need soil moisture and temperature data as forcing variables. Thus, this model can be used in a predictive manner, under the future environmental changes. Moreover, ED2.0-peat explicitly considers long-term changes in soil properties such as peat depth. Thus, the  $\text{CH}_4$  dynamics under a novel environment is estimated from an integrated system of soil temperature, moisture, and biogeochemistry. In addition, due to the characteristics of a mechanistic physics model, the model can reproduce a transient behavior in  $\text{CH}_4$  dynamics under climate change. Together with a study for the shallow peat site (Ise et al., 2008) ED2.0-peat correctly reproduced soil biophysics of various peat lands in Northern Manitoba.

To appropriately study the future  $\text{CH}_4$  in a predictive manner, ED2.0-peat explicitly simulated the dynamic interaction of soil temperature, moisture, and SOC in the peat land system. Our modeling scheme with mechanistically simulated soil hydrological and thermal processes has implications to future climate-soil interactions. An increase in temperature due to anthropogenic climate change will affect SOC decomposition rates because of their sensitivity to temperature. The Fen site lost 86% of SOC when subjected to a warming of  $4^\circ\text{C}$ , a rather conservative estimate of projected long-term climate change in boreal regions (IPCC, 2007). This significant collapse of peat land SOC is plausible when the global distribution of SOC is considered (Fig. 5). The transition from boreal to temperate biomes is currently characterized by a significant loss in stored SOC. If a temperature rise of  $4^\circ\text{C}$  at this site allows it to cross this threshold, a considerable shift in equilibrium SOC may occur. However, it should be noted that the Fen site also showed transient resistance against the collapse (Fig. 3c, simulation years 2000-2200); during the first few hundred years after the climate change, the decrease in SOC is rather slow, due to recalcitrance of slowly decomposing SOC.




Fig. 5. Global pattern of terrestrial SOC and mean annual temperature. Data from  $1^\circ \times 1^\circ$  mean annual temperature of 1961-1990 and SOC up to 1 m (Global Soil Data Task 2000). The curve is a piecewise 90-percentile quintile regression.

For this study, as noted in Ise et al. (2008), we intentionally turned off the dynamic vegetation processes of ED2.0 (Medvigy et al., 2009) to construct the current version of ED2.0-peat. Responses of vegetation processes against climate change such as litter production may have a strong effect on CH<sub>4</sub> dynamics through substrate quantity and quality. However, as shown here, this study is the first attempt to explicitly simulate a long-term physical interplay between soil properties and biogeochemistry, and the resultant transient dynamics was complex and nonlinear. Our aim here was to show the inherent behavior in soil. Considering dynamic vegetation will increase the complexity of the system further, and a systematic analysis will be very difficult. Thus, we kept the model as simple as possible for this study, and an explicit treatment of dynamic vegetation will be our next project. In summary, this study has shown how the mechanistic linkages that exist between the physical and biogeochemical dynamics of peat lands have strong implications for the response of northern peat lands to climate change.


### Acknowledgements

We appreciate helpful discussion with Profs. Steve Frohling, David Foster, and James McCarthy. The study of TI is supported by the James Mills Peirce Fellowship provided by the Department of Organismic and Evolutionary Biology at Harvard University and the Innovative Program of Climate Change Projection for the 21st Century (KAKUSHIN Program) of the Ministry of Education, Culture, Sports, Science and Technology, Japan.

### 5. References

- Apps, M.J., Halliwell, D., 1999. BOREAS TE-13 biometry reports, data set. Available online at [<http://www.daac.ornl.gov>] from Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, TN, U.S.A.
- Bond-Lamberty, B., Wang, C., Gower, S.T., 2004. Net primary production and net ecosystem production of a boreal black spruce wildfire chronosequence. *Global Change Biology* 10, 473-487.
- Clymo, R.S., 1984. The limits to peat bog growth. *Philosophical Transactions of the Royal Society of London, Series B* 303, 605-654.
- Dunn, A.L., Barford, C.C., Wofsy, S.C., Goulden, M.L., Daube, B.C., 2007. A long-term record of carbon exchange in a boreal black spruce forest: means, responses to interannual variability, and decadal trends. *Global Change Biology* 13, 577-590.
- Friedlingstein, P., Cox, P., Betts, R., Bopp, L., Von Bloh, W., Brovkin, V., Cadule, P., Doney, S., Eby, M., Fung, I., Bala, G., John, J., Jones, C., Joos, F., Kato, T., Kawamiya, M., Knorr, W., Lindsay, K., Matthews, H.D., Raddatz, T., Rayner, P., Reick, C., Roeckner, E., Schnitzler, K.G., Schnur, R., Strassmann, K., Weaver, A.J., Yoshikawa, C., Zeng, N., 2006. Climate-carbon cycle feedback analysis: results from the (CMIP)-M-4 model intercomparison. *Journal of Climate* 19, 3337-3353.
- Frohling, S., Roulet, N., Fuglestedt, J., 2006. How northern peatlands influence the earth's radiative budget: sustained methane emission versus sustained carbon sequestration. *Journal of Geophysical Research* 111, G01008, doi:10.1029/2005JG000091.

- Global Soil Data Task, 2000. Global soil data products CD-ROM (IGBP-DIS). International Geosphere-Biosphere Programme - data and information services. Available online at [<http://www.daac.ornl.gov/>] from Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, TN, U.S.A.
- Holland, M.M., Bitz, C.M., 2003. Polar amplification of climate change in coupled models. *Climate Dynamics* 21, 221-232.
- Intergovernmental Panel on Climate Change, 2007. Climate change 2007: the scientific basis. Available online at [[http://www.grida.no/climate/ipcc\\_tar/](http://www.grida.no/climate/ipcc_tar/)].
- Ise, T., Moorcroft, P.R., 2006. The global-scale temperature and moisture dependencies of soil organic carbon decomposition: an analysis using a mechanistic decomposition model. *Biogeochemistry* 80, 217-231.
- Ise, T., Dunn, A.L., Wofsy, S.C., Moorcroft, P.R., 2008. High sensitivity of peat decomposition to climate change through water-table feedback. *Nature Geoscience* 1, 763-766.
- Medvigy, D., Wofsy, S.C., Munger, J.W., Hollinger, D.Y., Moorcroft, P.R., 2009. Mechanistic scaling of ecosystem function and dynamics in space and time: Ecosystem Demography model version 2. *Journal of Geophysical Research—Biogeosciences*, 114, G01002, 10.1029/2008JG000812
- Parton, W.J., Schimel, D.S., Cole, C.V., Ojima, D.S., 1987. Analysis of factors controlling soil organic matter levels in Great Plains Grasslands. *Soil Science Society of America Journal* 51, 1173-1179.
- Pelletier, L., Moore, T.R., Roulet, N.T., Garneau, M., Beaulieu-Audy, V., 2007. Methane fluxes from three peatlands in the La Grande Riviere watershed, James Bay lowland, Canada. *Journal of Geophysical Research—Biogeosciences* 112, G01018, doi:10.1029/2006JG000216.
- Schlesinger, W.H., 1997. *Biogeochemistry: an Analysis of Global Change*. Academic Press, San Diego, CA, 588pp.
- Sellers, P., Hall, F., Margolis, H., Kelly, B., Baldocchi, D., Denhartog, G., Cihlar, J., Ryan, M.G., Goodison, B., Crill, P., Ranson, K.J., Lettenmaier, D., Wickland, D.E., 1995. The boreal ecosystem-atmosphere study (BOREAS): an overview and early results from the 1994 field year. *Bulletin of American Meteorological Society* 76, 1549-1577.
- Trumbore, S.E., Harden, J.W., 1997. Accumulation and turnover of carbon in organic and mineral soils of the BOREAS northern study area. *Journal of Geophysical Research—Atmospheres* 102, 28817-28830.
- Veldhuis, H., Eilers, R.G., Mills, G.F., 2002. Permafrost distribution and soil climate in the glacial Lake Agassiz basin in northcentral Manitoba, Canada. In: 17th World Conference on Soil Science, Bangkok, Thailand, 13-20 August.
- Walter, B.P., Heimann, M. 2000. A process-based, climate-sensitive model to derive methane emissions from natural wetlands: application to five wetland sites, sensitivity to model parameters, and climate. *Global Biogeochemical Cycles*, 14, 745-765.



## **Climate Change and Variability**

Edited by Suzanne Simard

ISBN 978-953-307-144-2

Hard cover, 486 pages

**Publisher** Sciyo

**Published online** 17, August, 2010

**Published in print edition** August, 2010

Climate change is emerging as one of the most important issues of our time, with the potential to cause profound cascading effects on ecosystems and society. However, these effects are poorly understood and our projections for climate change trends and effects have thus far proven to be inaccurate. In this collection of 24 chapters, we present a cross-section of some of the most challenging issues related to oceans, lakes, forests, and agricultural systems under a changing climate. The authors present evidence for changes and variability in climatic and atmospheric conditions, investigate some of the impacts that climate change is having on the Earth's ecological and social systems, and provide novel ideas, advances and applications for mitigation and adaptation of our socio-ecological systems to climate change. Difficult questions are asked. What have been some of the impacts of climate change on our natural and managed ecosystems? How do we manage for resilient socio-ecological systems? How do we predict the future? What are relevant climatic change and management scenarios? How can we shape management regimes to increase our adaptive capacity to climate change? These themes are visited across broad spatial and temporal scales, touch on important and relevant ecological patterns and processes, and represent broad geographic regions, from the tropics, to temperate and boreal regions, to the Arctic.

### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Takeshi Ise, Allison Dunn, Steven Wofsy and Paul Moorcroft (2010). Simulating Peatland Methane Dynamics Coupled to a Mechanistic Model of Biogeochemistry, Hydrology, and Energy: Implications to Climate Change, *Climate Change and Variability*, Suzanne Simard (Ed.), ISBN: 978-953-307-144-2, InTech, Available from: <http://www.intechopen.com/books/climate-change-and-variability/simulating-peatland-methane-dynamics-coupled-to-a-mechanistic-model-of-biogeochemistry-hydrology-and>

**INTECH**  
open science | open minds

### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166

### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821

[www.intechopen.com](http://www.intechopen.com)

[www.intechopen.com](http://www.intechopen.com)

IntechOpen

IntechOpen

© 2010 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](#), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen