

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,700

Open access books available

121,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# Assessment of Environmental Exposure to Benzene: Traditional and New Biomarkers of Internal Dose

Piero Lovreglio<sup>1</sup> et. al.\*

<sup>1</sup>*Dipartimento di Medicina Interna e Medicina Pubblica,  
Sezione di Medicina del Lavoro "E.C. Vigliani"  
University of Bari, Bari,  
Italy*

## 1. Introduction

In view of the widespread ubiquity of benzene in the environment and of its carcinogenic effects on man, this toxicant poses a public health problem that has prompted nations to undertake active measures to contain environmental concentrations below the limit judged to be an acceptable risk threshold for the general population (World Health Organization [WHO], 2000).

In the occupational field, until the 1950s benzene was the solvent most commonly employed in some industrial processes, and especially in rubber, printing and shoemaking industries, because of its chemico-physical properties and low cost. This caused exposure of these workers to high benzene concentrations that induced toxic effects and acute non lymphocytic leukemia (Agency for Toxic Substances and Disease Registry [ATSDR], 2007; WHO, 1993). Due to these adverse effects, its use in industrial processes was then abandoned, replacing benzene firstly by hexane, but this proved to provoke peripheral neuropathies, and then by less toxic solvents such as heptane. In Italy the use of benzene as a solvent is banned by Law 245/1963, although traces below 2% are permitted in solvents of a different chemical nature (Italian Parliament, 1963).

Benzene is still used as a raw material or intermediate product in the chemical industry, mainly to synthesize ethylbenzene, cumene and cyclohexane, and to a limited extent as a

---

\* Maria Nicolà D'Errico<sup>1</sup>, Silvia Fustinoni<sup>2</sup>, Ignazio Drago<sup>1</sup>, Anna Barbieri<sup>3</sup>, Laura Sabatini<sup>3</sup>, Mariella Carrieri<sup>4</sup>, Pietro Apostoli<sup>5</sup>, Leonardo Soleo<sup>1</sup>

<sup>1</sup> *Dipartimento di Medicina Interna e Medicina Pubblica, Sezione di Medicina del Lavoro "E.C. Vigliani".  
University of Bari, Bari, Italy*

<sup>2</sup> *Dipartimento di Medicina del Lavoro, University of Milan and Fondazione IRCCS Ca' Granda Ospedale  
Maggiore Policlinico, Milano, Italy.*

<sup>3</sup> *Dipartimento di Medicina Interna, dell'Invecchiamento e Malattie Nefrologiche,  
Unità Operativa di Medicina del Lavoro, University of Bologna, Bologna, Italy*

<sup>4</sup> *Dipartimento di Medicina Ambientale e Sanità Pubblica, University of Padova, Padova, Italy*

<sup>5</sup> *Dipartimento di Medicina Sperimentale ed Applicata, Sezione di Medicina del Lavoro ed Igiene Industriale,  
University of Brescia, Brescia, Italy*

chemical reagent in the laboratory. It is contained in crude oil, that is its main source of production nowadays, and is formed as a result of the incomplete combustion of fossil fuels such as coal and, to a lesser extent, wood.

Moreover, fuels derived from crude oil contain benzene not only because it is already present in the raw material but also because it is formed during the refining process (Brief et al. 1980; Holmberg & Lundberg 1985). For this reason, occupational exposure to benzene, albeit at concentrations of 2-3 orders of magnitude below the occupational limit values, such as the threshold limit value-time weighted average (TLV-TWA) of 1600  $\mu\text{g}/\text{m}^3$  proposed by the American Conference of Governmental Industrial Hygienists (ACGIH), and the European limit value of 3250  $\mu\text{g}/\text{m}^3$  issued by the European Directive 1999/38/CE, is still present in oil refining and petrochemical industries, as well as in fuel tanker drivers and filling station attendants, workers in cokeries and in chemical laboratories (ACGIH, 2010; European Parliament, 1999).

The natural sources of emission into the atmosphere, originally volcanoes and fires, play a negligible role while human activities are the main source of benzene released in the environment (ATSDR, 2007). Among human activities one of the principal forms of emissions of benzene in the environment is in automobile exhaust. Evidence in literature reports higher concentrations of this toxicant in urban and dense traffic areas than in less busy traffic and rural areas.

Another source of environmental pollution by benzene is evaporation during filling operations at gasoline stations or during loading and unloading of fuel tanker lorries, even if the use of aspiration systems to retrieve the vapors during such operations has significantly reduced these emissions, by up to 75% (Duarte-Davidson et al., 2001). In addition, higher concentrations of benzene than the background outdoor levels are present inside vehicles, including buses and private cars, due not only to the penetration inside the body of the vehicle of exhaust fumes from other vehicles, but also to leaks from gasoline tanks and gasoline leads and circuits (Geiss et al., 2009; Li et al., 2009). The levels of benzene emitted by vehicles depend on the type and age of the vehicle, the type of traffic and the ventilation within the body of the car (Duarte-Davidson et al., 2001). Finally, in some geographic areas, apart from road traffic, other sources of benzene emission into the atmosphere, such as factories, hazardous waste dumps and domestic wood fires, can play a significant role in causing outdoor pollution (Barrefors & Petersson, 1995; Edgerton & Shah, 1992).

In western nations, the pollution of urban areas by benzene has gradually declined in recent years, thanks to the legislation introduced to reduce the content of benzene in fuels, currently limited in Italy to concentrations below 1% in volume (law n. 413/97), and to the application in the EU of the norms targeting an annual average exposure for benzene of 5  $\mu\text{g}/\text{m}^3$  as from 2010 (European Commission, 2000; Italian Parliament, 1997). Airborne benzene concentrations are proportionally lower than the true quantity of emissions thanks to the rapid chemical degradation of this toxicant, largely as a reaction to hydroxyl radicals. This limits the persistence of volatile benzene in the atmosphere to a few days or hours.

Active cigarette smoking is another important source of exposure to benzene in the general population, as mainstream smoke contains quantities of 28.0-105.9  $\mu\text{g}/\text{cigarette}$ , and sidestream smoke no less than 70.7-134.3  $\mu\text{g}/\text{cigarette}$  (International Agency for Research on Cancer [IARC], 2004). Wallace (1989) estimated a benzene intake of 1.8 mg/day with an average consumption of 32 cigarettes/day, equal to at least 10-fold the intake of a non-smoker, while Duarte-Davidson et al. (2001) estimated a dose of 400  $\mu\text{g}/\text{day}$  of benzene

retained within the organism by a smoker of 20 cigarettes/day. Thus, cigarette smoking can induce a similar or higher benzene intake than that occurring in most occupationally exposed workers in western nations, and certainly higher than the intake caused by environmental exposure to airborne benzene levels near the upper limits for air quality. Moreover, the role of smoking as a source of benzene has become progressively more important as urban pollution levels have declined (Duarte-Davidson et al. 2001; Hattemer-Frey et al. 1990). In fact, in the period 1989-1997 Fruin et al. (2001) observed a marked reduction in the benzene quota derived from the environment (12%) and thus a proportional increase in the quota due to cigarette smoking (78%). Passive smoking can also induce significant exposure to benzene, equal to about 10% of the total intake in non smokers, and higher than the quota contributed by the entire set of US industrial emissions in the atmosphere (Duarte-Davidson et al., 2001; Wallace, 1995). Confirming this, higher benzene concentrations have been reported in homes with one or more smokers among the inhabitants (median  $10.6 \mu\text{g}/\text{m}^3$ ) than in non smoker homes ( $7.0 \mu\text{g}/\text{m}^3$ ) (Wallace 1989).

In man, benzene has myelotoxic and carcinogenic effects. As regards the former environmental exposure to benzene is unlikely to cause the myelosuppression effects previously observed in occupational contexts, since a NOAEL of  $1790 \mu\text{g}/\text{m}^3$  has been individuated for these effects, that will probably never again be reached in the human living environment (Collins et al., 1997). The carcinogenic effect manifests in the form of a greater prevalence of acute non lymphocytic leukemia, as stated above. Benzene has been classified carcinogenic to humans by the IARC, ever since the first classification published in the 1980s (IARC, 1982). A review conducted in 2009 confirmed benzene in the same group, associated with acute non lymphocytic leukemia. Instead, there is still only limited evidence of a causal relationship with the onset of acute lymphocytic leukemia, chronic lymphocytic leukemia, multiple myeloma and non-Hodgkin lymphoma (Baan et al., 2009).

Exposure to benzene is considered to induce the onset of leukemias because benzene acts through a genotoxic carcinogenic mechanism, whereby a dose-response relationship of a linear type has been proposed, with no threshold. Although this hypothesis has recently been cast in doubt, the level of exposure that can be considered "safe" has not been identified, but only a LOAEL of  $32000\text{-}80000 \mu\text{g}/\text{m}^3$  (Bolt et al., 2008; Duarte-Davidson et al., 2001). As to the carcinogenic risk posed by environmental exposure, the concentration of this toxicant judged to pose an acceptable risk in the general population, exposed over a lifetime, has been estimated on the basis of epidemiological studies of workers occupationally exposed to medium-high concentrations of benzene. The WHO estimate, in particular, predicts an excess risk of  $6 \text{ cases} \times 10^{-6}$  for chronic exposure to  $1 \mu\text{g}/\text{m}^3$  throughout life (WHO, 2000).

Absorption of benzene in the general population occurs virtually exclusively by inhalation, that accounts for more than 99% of the daily intake of this toxicant, whereas oral intake plays only a minor role, since the quantities of benzene present in water, foods and consumer products are minimal and skin intake is negligible (Hattemer-Frey et al., 1990). After being absorbed, benzene is partly eliminated as is in exhaled air, accounting for 16.8 % of the quota absorbed, and in urine, albeit only for 0.1% of the absorbed quota (Ghittori et al., 1993; Nomiya & Nomiya, 1974). The remainder is rapidly distributed throughout the organism and undergoes a biotransformation process, prevalently in the liver. This occurs very similar in man and experimental animals, and involves the initial formation of benzene oxide catalyzed by microsomal cytochrome P450 and especially isoenzyme 2E1

(Snyder & Heidli, 1996). After the formation of this reactive intermediate, many different metabolites are then formed in succession through different metabolic pathways, namely phenol, catechol and hydroquinone by oxidation, again catalyzed by cytochrome P450, S-phenylmercapturic acid (SPMA) through the action of the glutathione transferase system, 1,2-benzene dihydrodiol through the action of epoxide hydrolase and the subsequent oxidation to catechol and *t,t*-muconic acid (*t,t*-MA) through the opening of the benzene ring (Snyder et al., 1993). All these metabolites are then excreted in the urine in the form of sulfate or glucuronate derivatives.

For the progressive reduction of the benzene workplace concentrations to which workers are exposed, ever more sensitive and specific biomarkers have needed to be identified, that maintain their validity even in conditions of low or extremely low benzene concentrations. This is why urinary phenol is no longer used in routine practice, because it has poor validity for exposure to less than 16250  $\mu\text{g}/\text{m}^3$ , a value that is still one order of magnitude greater than the TLV-TWA of the ACGIH (Boogard & van Sitter, 1995; Ong et al., 1995). The biological markers currently recommended by the ACGIH are *t,t*-MA and SPMA, that reveal an increased urinary excretion already at levels of exposure to airborne benzene concentrations of 65  $\mu\text{g}/\text{m}^3$ , while the relation becomes unambiguous at concentrations of more than 650  $\mu\text{g}/\text{m}^3$  (Kim et al., 2006). A new marker of internal dose is now under study, namely the determination of benzene as is in the urine (urinary benzene) that may prove particularly useful to monitor exposure to very low concentrations of airborne benzene, since a correlation has been shown with benzene concentrations ranging from 6 to 478  $\mu\text{g}/\text{m}^3$  (Fustinoni et al., 2005).

Despite the ample volume of studies in the occupational field, the behavior and significance of the above-described biomarkers of internal dose are still little known when used to monitor environmental exposure to airborne benzene concentrations near the upper limit of 5  $\mu\text{g}/\text{m}^3$  for air quality. In particular, the validity of these markers needs to be confirmed as a means of excluding environmental exposure beyond this limit, and hence the presence of a carcinogenic risk above what is considered acceptable. The behavior of these biomarkers of internal dose also needs to be assessed in relation to the relative contributions made to the overall intake of benzene by cigarette smoking and urban pollution, the main sources of environmental exposure to the toxicant.

Aim of the present study was to assess the significance and limits of *t,t*-MA, SPMA and urinary benzene for biological monitoring of subjects with non occupational exposure to very low concentrations of benzene as those found in the general environment, as well as to study the influence of the different sources of environmental exposure on these biomarkers.

## 2. Materials and methods

### 2.1 Subjects

The study sample included 123 adult males resident in the cities and hinterland of Bari and Foggia (Apulia - Italy), all in good health and with no occupational exposure to benzene. In the geographic area where this study has been conducted there are no major industries or waste incinerators that could produce benzene emissions. All participants filled out a questionnaire posing questions about personal data, job at the time of the study, smoking habit with particular reference to the number of cigarettes smoked during the environmental sampling, alcohol consumption, personal medical history, time spent in

urban traffic during the environmental sampling, and hobbies. All subjects gave prior written consent to take part in the study.

## 2.2 Air sampling

Exposure to airborne environmental benzene was monitored in all subjects by passive personal sampling, using radial diffusion samplers (Radiello®) containing an active carbon cartridge, worn in the respiratory zone for 8 hours, typically from 8:30 a.m. to 4:30 p.m. After the sampling, the Radiello® vials were preserved at +4°C until the time of analysis.

Analysis of the vials was then performed by gas chromatography – flame ionization detector after desorption of the benzene from the active carbon with a carbon disulfide low benzene content according to the method reported by the manufacturer (Supelco, 2010). The detection limit of the procedure for benzene was 2 µg/m<sup>3</sup>. All analyses were performed blinded.

## 2.3 Urine biomonitoring

Immediately after the environmental sampling, a urine sample was collected from all participants for assays of *t,t*-MA, SPMA and urinary benzene. Each urine sample was subdivided into two aliquots: 30 ml of urine were set aside to determine *t,t*-MA, SPMA and urinary creatinine, preserved in sterile containers, without the addition of preservatives or stabilizers, at -20°C until the time of analysis. The second aliquot, 10 ml of urine used to determine urinary benzene, was immediately transferred to a presealed 20 ml vial containing 4 g of NaCl and preserved at +4°C until the time of analysis. All analyses were conducted blinded.

Urinary *t,t*-MA analysis was carried out with the HPLC-UV method at 264 nm, after solid phase extraction (SPE) (SAX column-Varian) by an analytical method described elsewhere (Aprea et al., 2008). The limit of detection (LOD) of the procedure was 20 µg/L.

The analytical determination of urinary SPMA was performed following the application described in Sabatini et al. (2008). Briefly, after SPE and LC separation, samples were analyzed by a liquid chromatography/electrospray tandem mass spectrometry method (HPLC-ESI-MS/MS), operated in negative ion mode, using isotope-labeled analogs as internal standards. The LOD of the method was 0.03 µg/L.

The analytical determination of benzene in urine was performed by headspace analysis with automated solid phase micro-extraction (SPME), following a previously reported method (Barbieri et al., 2008). A gas chromatograph equipped with a split-splitless injector and coupled to a mass selective detector (GC/MS) was used for analysis. An autosampler (CTC Combi PAL system) was interfaced to GC/MS system for the SPME process. The LOD of the method was 0.02 µg/L.

Analyses of urinary creatinine were performed using the DCA 2000®+ analyzer. The creatinine assay is based on the Benedict/Behre test, and was performed on the same urine samples used for all the other analyses, collected at the end of the environmental sampling (Benedict & Behere, 1936).

All the laboratories conducting the analytical measurements conform to quality assurance procedures and participate in quality control programs.

## 2.4 Statistical analysis

Statistical analyses were done with the SPSS program (14.0 version, Chicago, IL, USA). A value corresponding to one-half of the LOD was assigned to measurements below the

analytical detection limit. A normal distribution of all the variables was checked with the Kolmogorov-Smirnov test. Non normally distributed variables were analyzed by parametric tests after logarithmic transformation or by non parametric tests. Correlation analyses were done with Spearman's test. Multiple linear regression models were applied to assess a dependency relation of the different biomarkers on the independent variables. The level of significance was set at  $p < 0.05$ .

### 3. Results

Of the 123 subjects, 55 were smokers and 68 non smokers, and the population sample featured a wide age (21-62 years) and BMI (19.4-44.4 Kg/m<sup>2</sup>) range (Table 1).

For all the biomarkers determined, the percentage of cases in the entire sample exceeding the LOD was significantly higher in smokers than non smokers. When the whole sample was subdivided according to exposure to urban traffic or not during the environmental sampling, a higher percentage of results above the LOD was found only for airborne benzene and urinary benzene (Table 2).

	N.	Mean±SD	Median	Range
Age (years)	123	41.5±11.8	44.0	21-62
Body mass index (Kg/m <sup>2</sup> )	123	26.8±4.8	25.8	19.4-44.4
Residence				
- urban	79	64.2%		
- rural	44	35.8%		
Alcohol consumption				
- Teetotal	20	16.2%		
- <10 g/day	51	41.5%		
- >10 g/day	52	42.3%		
Smoking habit				
- Smokers	55	44.7%		
- Non smokers	68	55.3%		
N. cigarettes/day*	55	15.8±8.6	15.0	5-40
N. cigarettes smoked during environmental sampling*	55	6.6±4.1	6.0	0-20
Time between last sigarete and urine collection (minutes)*	53	54.3±69.0	30.0	2-360
Exposure to urban traffic during environmental sampling				
- Yes	27	21.9%		
- No	96	78.1%		

\*In smokers

Table 1. Personal data of the 123 subjects examined.

	Smoking habit				Exposure to urban traffic			
	Yes		No		Yes		No	
	N <sub>&gt;LOD</sub> /N	N <sub>&gt;LOD</sub> /N%	N <sub>&gt;LOD</sub> /N	N <sub>&gt;LOD</sub> /N%	N <sub>&gt;LOD</sub> /N	N <sub>&gt;LOD</sub> /N%	N <sub>&gt;LOD</sub> /N	N <sub>&gt;LOD</sub> /N%
Airborne benzene	14/55	25.5%	20/68	29.4%	18/27	66.7% <sup>c</sup>	16/96	16.7%
<i>t,t</i> -MA	55/55	100.0% <sup>a</sup>	62/68	91.2%	27/27	100.0%	90/96	93.7%
SPMA	44/53	83.0% <sup>b</sup>	2/66	3.0%	12/27	44.4%	34/92	37.0%
Urinary benzene	53/54	98.1% <sup>b</sup>	22/65	33.8%	25/27	92.6% <sup>c</sup>	50/92	54.3%

Smokers vs non smokers: <sup>a</sup>p<0.05; <sup>b</sup>p≤0.001; Exposed vs not exposed to urban traffic: <sup>c</sup>p<0.001

Table 2. Percentage of determinations of airborne benzene, *t,t*-MA, SPMA and urinary benzene above the LOD in the sample, subdivided by smoking habit and exposure to urban traffic during sampling.

To assess a simultaneous influence of smoking habit and exposure to urban traffic on the levels of airborne benzene and the urinary biomarkers, two-way analysis of variance was done, including in the model only the environmental and urinary values exceeding the LOD (Table 3). The results demonstrated an evident association between airborne benzene or *t,t*-MA or urinary benzene and cigarette smoking, whereas it was not possible to perform the analysis for SPMA because only 2 determinations were above the LOD in non smokers. In any case, there was no significant difference in the urinary concentrations of SPMA in smokers between subjects with or without exposure to urban traffic.

	Exposure to urban traffic	Smokers				Non smokers				ANOVA
		N <sub>&gt;LOD</sub>	Mean±SD	Median	Range	N <sub>&gt;LOD</sub>	Mean±SD	Median	Range	
Airborne benzene (µg/m <sup>3</sup> )	Yes	10	7.4±3.7	7.3	3.9-16.3	8	4.7±1.0	4.5	3.3-6.2	Model: F=8.3 <sup>a</sup> Smoke: F=8.4 <sup>a</sup> Traffic: F=4.0
	No	4	6.2±3.9	5.7	2.0-11.5	12	3.4±1.1	3.0	2.0-5.7	
<i>t,t</i> -MA (µg/g creat)	Yes	14	80.4±52.9	73.5	13.0-196.0	13	51.0±16.5	46.0	30.0-90.0	Model: F=13.0 <sup>b</sup> Smoke: F=14.1 <sup>b</sup> Traffic: F=0.0
	No	41	102.8±58.6	89.7	19.0-307.0	49	55.7±105.0	35.6	9.6-734.0	
SPMA (µg/g creat)	Yes	11	1.47±1.30	0.99	0.38-4.48	1	-	-	-	-
	No	33	1.54±1.19	1.28	0.29-5.22	1	-	-	-	
Urinary benzene (µg/L)	Yes	14	2.24±3.62	0.41	0.04-11.40	11	0.07±0.02	0.06	0.04-0.10	Model: F=21.2 <sup>b</sup> Smoke: F=58.7 <sup>b</sup> Traffic: F=0.0
	No	39	0.97±1.00	0.60	0.06-4.27	11	0.07±0.03	0.06	0.04-0.12	

<sup>a</sup>p<0.01; <sup>b</sup>p<0.001

Table 3. Concentrations of airborne benzene, *t,t*-MA, SPMA and urinary benzene in the determinations above the LOD in the sample, subdivided by smoking habit and exposure to urban traffic.

In non smokers, a possible association between exposure to passive smoke during the environmental sampling and higher airborne benzene levels or urinary excretion of its metabolites was studied. No association was found in non smokers exposed or not exposed to passive smoke and airborne benzene and its urinary metabolites (data not shown).

Analysis of correlations between the general lifestyle of the participants, variables linked to smoking habit and exposure to urban traffic, the airborne benzene concentrations and the biomarkers studied was done on the entire sample and then subdivided into smokers and non smokers (Table 4). Airborne benzene was found to be correlated to the time spent in urban traffic during the sampling, both in the entire sample and when analyzing smokers and non smokers separately, whereas there was no correlation between airborne benzene and the number of cigarettes smoked per day or during the sampling time (Table 4). Among the biomarkers studied, airborne benzene was correlated only with urinary benzene, and only in the non smoker group. When analyzing the entire sample together, all the biological markers studied were found to be significantly correlated to the number of cigarettes smoked per day or during the sampling time. In smokers this correlation was significant only for SPMA. Moreover, *t,t*-MA and urinary benzene were correlated in non smokers with the time spent in urban traffic; when taking the group as a whole, this correlation was confirmed for urinary benzene only, albeit with a low rho value. Finally, the three biological markers were all mutually correlated in the entire sample taken as a whole (Table 4).

		Age (years)	BMI	N. cig./day	N. cig./sampling	Time since last cig.	Alcohol	Urban traffic (minutes)	Airborne benzene	<i>t,t</i> -MA	SPMA
BMI (Kg/m <sup>2</sup> )	Total	0.35 <sup>c</sup>	-	-	-	-	-	-	-	-	-
	Smokers	0.36 <sup>b</sup>	-	-	-	-	-	-	-	-	-
	Non smokers	0.32 <sup>b</sup>	-	-	-	-	-	-	-	-	-
N. cig./day	Total	-0.22 <sup>a</sup>	-0.05	-	-	-	-	-	-	-	-
	Smokers	-0.08	0.11	-	-	-	-	-	-	-	-
	Non smokers	-	-	-	-	-	-	-	-	-	-
N. cig./sampling	Total	-0.21 <sup>a</sup>	-0.04	0.94 <sup>c</sup>	-	-	-	-	-	-	-
	Smokers	-0.02	0.08	0.55 <sup>c</sup>	-	-	-	-	-	-	-
	Non smokers	-	-	-	-	-	-	-	-	-	-
Time since last cigarette	Total	-	-	-	-	-	-	-	-	-	-
	Smokers	0.23	0.08	-0.48 <sup>c</sup>	-0.22	-	-	-	-	-	-
	Non smokers	-	-	-	-	-	-	-	-	-	-
Alcohol	Total	0.37 <sup>c</sup>	0.25 <sup>b</sup>	0.005	0.01	-	-	-	-	-	-
	Smokers	0.35 <sup>b</sup>	0.14	0.02	0.11	0.12	-	-	-	-	-
	Non smokers	0.38 <sup>c</sup>	0.37 <sup>b</sup>	-	-	-	-	-	-	-	-
Urban traffic	Total	0.14	0.07	0.04	0.04	-	0.11	-	-	-	-
	Smokers	0.46 <sup>c</sup>	0.31 <sup>a</sup>	-0.13	-0.04	0.21	0.25	-	-	-	-
	Non smokers	-0.17	-0.14	-	-	-	-0.03	-	-	-	-
Airborne benzene	Total	0.17	0.09	0.001	0.03	-	0.11	0.51 <sup>c</sup>	-	-	-
	Smokers	0.33 <sup>a</sup>	0.26	-0.03	0.17	-0.01	0.13	0.61 <sup>c</sup>	-	-	-
	Non smokers	-0.01	-0.06	-	-	-	0.10	0.40 <sup>c</sup>	-	-	-
<i>t,t</i> -MA	Total	-0.15	-0.03	0.57 <sup>c</sup>	0.55 <sup>c</sup>	-	-0.02	0.04	0.04	-	-
	Smokers	0.03	0.10	0.23	0.07	-0.11	0.01	-0.22	-0.13	-	-
	Non smokers	-0.12	-0.12	-	-	-	-0.08	0.29 <sup>a</sup>	0.22	-	-
SPMA	Total	0.05	-0.01	0.76 <sup>c</sup>	0.76 <sup>c</sup>	-	0.11	0.003	0.02	0.56 <sup>c</sup>	-
	Smokers	0.32 <sup>a</sup>	0.08	0.42 <sup>c</sup>	0.43 <sup>c</sup>	-0.17	0.26	-0.13	0.04	0.28 <sup>a</sup>	-
	Non smokers	0.33 <sup>b</sup>	0.04	-	-	-	0.09	0.02	-0.05	0.19	-
Urinary benzene	Total	-0.30 <sup>c</sup>	-0.06	0.77 <sup>c</sup>	0.74 <sup>c</sup>	-	-0.13	0.19 <sup>a</sup>	0.16	0.44 <sup>c</sup>	0.56 <sup>c</sup>
	Smokers	-0.12	0.10	0.23	0.17	-0.18	-0.38 <sup>b</sup>	0.01	0.20	0.14	0.23
	Non smokers	-0.19	-0.08	-	-	-	0.07	0.50 <sup>c</sup>	0.31 <sup>a</sup>	0.06	-0.19

<sup>a</sup>p<0.05; <sup>b</sup>p<0.01; <sup>c</sup>p≤0.001

Table 4. Spearman correlations among personal data, lifestyle, airborne benzene and biomarkers in the whole sample and subdivided by smoking habit.

The dependency of *t,t*-MA, SPMA and urinary benzene on the variables age, BMI, number of cigarettes/day, urban traffic and airborne benzene was studied in the sample as a whole,

applying different multiple linear regression models. The results demonstrated a dependency of the urinary concentrations of *t,t*-MA and SPMA on the number of cigarettes smoked per day, and of urinary benzene both on the number of cigarettes/day and on the time spent in urban traffic (Table 5).

To verify the influence of cigarette smoking on the biotransformation of benzene, the urinary benzene/*t,t*-MA ratio was studied in smokers and non smokers, and showed significantly higher levels in smokers (median 0.0071 vs 0.0008;  $p \leq 0.001$ ).

	<i>t,t</i> -MA		SPMA		Urinary benzene		
	t	p	t	p	t	p	
Age (years)	NS		NS		NS		
BMI (Kg/m <sup>2</sup> )	NS		NS		NS		
N. cigarettes/day	5.8	<0.001	11.9	<0.001	12.5	<0.001	
Urban traffic (minutes)	NS		NS		2.1	0.034	
Airborne benzene (µg/m <sup>3</sup> )	NS		NS		NS		
Model	F	p	R <sup>2</sup>	F	p	R <sup>2</sup>	
	34.2	<0.001	0.22	141.0	<0.001	0.55	
					F	p	R <sup>2</sup>
					80.6	<0.001	0.59

NS= non significant

Table 5. Multiple linear regression analysis of the whole sample, taken as a single group, for the dependent variables *t,t*-MA, SPMA and urinary benzene.

#### 4. Discussion

This research analyzed the contribution of both traditional and new biological markers of internal dose, namely *t,t*-MA and urinary SPMA as compared to urinary benzene, to the monitoring of environmental exposure to very low concentrations of benzene. A particular attention was paid to how much a smoking habit and exposure to urban traffic, the main non occupational sources of this toxicant, could condition the urinary excretion of the various biomarkers.

Environmental monitoring demonstrated that overall, the general population studied was exposed to only very low concentrations of benzene, with 72.4% of the air samples showing values of less than 2 µg/m<sup>3</sup>, a much higher percentage than in a recent study of a population with no occupational exposure resident in a large Northern Italian city with very dense road traffic, Milan. In that study, the 5<sup>th</sup> percentile of environmental benzene distribution was equal to 1.5 µg/m<sup>3</sup> and the 25<sup>th</sup> percentile to 3.0 µg/m<sup>3</sup> (Fustinoni et al., 2010). Thus, our results suggest that our study sample, resident in small-medium cities with a population of less than 500 thousand inhabitants, is exposed to generally similar or lower levels of environmental pollution by benzene than inhabitants of other industrial, urban and suburban/residential/rural areas in Italy and worldwide (Table 6). Apart from the lesser traffic volume, this difference is also due to climatic conditions, in particular the strong winds that commonly blow in this area and tend to prevent the stagnation of pollutants. Nevertheless, exposure to road traffic was in any case the main factor determining environmental benzene concentrations exceeding the LOD, even if no significantly higher benzene concentrations could be determined in the traffic-exposed group as compared to the non-exposed group when analyzing only the values obtained in excess of the LOD, possibly due to small sample sizes.

Although the airborne benzene levels measured were generally low, concentrations above the threshold limit for air quality of  $5 \mu\text{g}/\text{m}^3$  were observed in 13 cases, highlighting the fact that exposure to excess benzene levels in the daily environment is still possible in western nations even nowadays. In fact, the threshold of  $5 \mu\text{g}/\text{m}^3$  represents the mean annual exposure posing a carcinogenic risk considered to lie within acceptable limits for the general population with exposure over a lifetime. No definitive level of exposure below which the genotoxic carcinogenic effect of benzene is completely revoked has yet been identified, while adopting a threshold of 0, the only value that could guarantee the absence of risk, seems practically impossible bearing in mind the widespread ubiquity of benzene sources. It is now common practice to use the threshold value for air quality when interpreting the results of personal sampling for carcinogenic environmental agents in both occupational and non-occupational settings. All the same, it must be taken into account that the threshold value represents a mean value referred to annual measurements, and is thus less affected in the long term by the inevitably wide variability of benzene concentrations occurring over such a long period, whereas environmental samplings conducted as in our study refer to a short period, in our case 8 hours, and can therefore be affected by occasional peaks of exposure. Therefore, the finding of values in excess of the  $5 \mu\text{g}/\text{m}^3$  threshold during sampling lasting 8 hours does not necessarily indicate a raised health risk.

As regards the influence of a smoking habit, previous studies have demonstrated that the benzene concentrations measured with personal samplers do not reflect the true level of exposure to benzene induced by cigarette smoke (Fustinoni et al., 2005; Lovreglio et al., 2010). The results of the present study partly confirm reports in literature, since a smoking habit did not affect the percentage of benzene determinations exceeding the LOD, and no correlation was observed between airborne benzene and the number of cigarettes smoked during the sampling period. However, higher concentrations of airborne benzene were observed in smokers when only the values exceeding the LOD were analyzed, even after stratification for exposure to urban traffic.

OCCUPATIONAL EXPOSURE						
Occupational settings	Sampling	Year	Exposure levels ( $\mu\text{g}/\text{m}^3$ )			References
			Mean $\pm$ SD	Median	Range	
Chemicals manufacture Heavy truck drivers Crude petroleum extraction	Personal (8 h) <sup>NA</sup>	1996-2007	357.5 $\pm$ 1300.0 357.5 $\pm$ 195.0 97.5 $\pm$ 260.0	- - -	6.5-1300.0 81.2-487.5 3.25-975.0	Scarselli et al., (2011)
Fuel tanker drivers Filling station attendants	Personal (8 h) <sup>*</sup>	2006	306.7 $\pm$ 266.7 23.5 $\pm$ 17.4	246.6 20.9	7.4-1017.1 4.5-66.3	Lovreglio et al., (2010)
Petrochemical factories Gasoline service stations	Ambient (8 h) <sup>**</sup>	-	214.6 $\pm$ 78.8 209.9 $\pm$ 57.0	34.6 114.3	5.3-1766.2 5.6-773.6	Navasumrit et al., (2005)
Refinery blue collars	Personal (8h) <sup>***</sup>	-		190	60-2310	Fustinoni et al., (2011)
Sewage workplace	Personal (4 d) <sup>*</sup>	2008-2009	19.1 $\pm$ 2.9			Al Zabadi et al.(2011)
Urban policemen	Personal (6 h) <sup>***</sup>	2004		9.6	5.4-22.5	Campo et al., (2011)

ENVIRONMENTAL EXPOSURE						
City	Sampling	Year	Exposure levels ( $\mu\text{g}/\text{m}^3$ )			References
			Mean $\pm$ SD	Median	Range	
INDUSTRIAL AREA						
Lin-Yuan TWN Ping-Tung TWN	Ambient Outdoor (1-2 h)****	2003-2004	25.8 $\pm$ 34.7 5.9 $\pm$ 3.4	8.4 5.0	3.7-120.6 ND-13.7	Hsieh et al., (2006)
La Plata ARG	Ambient Indoor (4 w)** Ambient Outdoor (4 w)**	2000-2002	19.1 16.1	18.0 13.4	max 59.5 max 37.2	Massolo et al., (2009)
Yokohama JPN	Ambient Outdoor ****	2007-2008	6.8 $\pm$ 8.7			Tiwari et al., (2010)
Dunkerque FRA	Ambient Outdoor *	2007	1.9 $\pm$ 2.1		0.8-6.7	Roukos et al., (2009)
URBAN AREA						
Bangkok THA	Ambient Outdoor Roadside** Ambient Outdoor School**	-	109.2 $\pm$ 22.4 26.7 $\pm$ 2.5	91.8 27.9	50.2-212.9 22.5-28.7	Navasumrit et al., (2005)
Naples ITA	Ambient Outdoor (24h)*	2006	9.8 $\pm$ 4.4		4.4-17.2	Iovino et al., (2009)
Milano ITA	Personal (72 h)* Ambient Indoor work (1 w)* Ambient Outdoor work (1 w)*	2003	8.5 $\pm$ 3.0 3.0 $\pm$ 1.5 1.9 $\pm$ 1.4			Bruinen de Bruin et al., (2008)
Catania ITA	Personal (72 h)* Ambient Indoor work (1 w)* Ambient Outdoor work (1 w)*		5.2 $\pm$ 1.6 5.0 $\pm$ 3.4 4.2 $\pm$ 1.8			
Brussels BEL	Personal (72 h)* Ambient Indoor home (1 w)* Ambient Indoor work (1 w)* Ambient Outdoor work (1 w)*		3.2 $\pm$ 1.4 2.7 $\pm$ 1.2 2.1 $\pm$ 0.2 1.9 $\pm$ 0.6			
Helsinki FIN	Personal (72 h)* Ambient Indoor home (1 w)* Ambient Outdoor work (1 w)* Ambient Indoor work (1 w)*		2.0 $\pm$ 0.9 1.7 $\pm$ 1.0 1.0 $\pm$ 0.3 1.0 $\pm$ 0.2			
Bucharest ROM Madrid ESP Lisbon PRT Brussels BEL Ljubljana SVN Dublin IRL	Ambient Outdoor (1 d)*		2003 2003 2002 2002 2003 2004		7.1 4.5 3.8 2.5 3.1 1.1	
Bari ITA	Personal (8 h)*	2006	4.6 $\pm$ 2.6	4.3	<3.0-11.5	Lovreglio et al., (2010)
Milan ITA	Personal (5h)*	-		4.0	1.5-16.1	Fustinoni et al., (2011)
Beijing CHN	Ambient Outdoor (13 h)****	2008	3.7 $\pm$ 3.0			Liu et al., (2009)
LaPlata ARG	Indoor (4 w)** Outdoor (4 w)**	2000-2002	3.6 3.1	3.2 3.1	max 12.7 max 5.6	Massolo et al., (2009)
Bari ITA	Ambient Indoor (24h)*	2005	(weekly average) 1.3-14.8			Bruno et al., (2008)
Bari ITA	Ambient Outdoor (1 w)*	2008 Autumn 2008 Spring			1.3-9.0 0.8-2.7	Caselli et al., (2010)
Antwerp BEL	Ambient Outdoor (5 d)* Ambient Indoor (5 d)*	2002-2003	1.9 1.5		0.7-4.4 0.3-3.1	Stranger et al., (2008)

ENVIRONMENTAL EXPOSURE							
City	Sampling	Year	Exposure levels ( $\mu\text{g}/\text{m}^3$ )			References	
			Mean $\pm$ SD	Median	Range		
URBAN AREA							
Spain	Ambient Outdoor home (1 w)*	2004-2008	1.6 $\pm$ 0.9			Estarlich et al.,(2011)	
Toulouse FRA	Ambient Outdoor*	2001	1.1 $\pm$ 0.3		2.0-0.7	Simon et al., (2004)	
SUBURBAN/ RESIDENTIAL/ RURAL AREA							
Naples ITA	Ambient Outdoor (24h)*	2006	5.7 $\pm$ 3.2		2.3-12.8	Iovino et al., (2009)	
LaPlata ARG	Indoor (4w)**	2000-2002	4.7	3.1	max 13.2	Massolo et al., (2009)	
	Outdoor (4w)**		1.6	1.7	max 1.8		
Spain	Ambient Outdoor home (1 w)*	2004-2008	1.5 $\pm$ 0.7			Estarlich et al., (2011)	
Yokohama JPN	Ambient Outdoor****	2007-2008	1.3 $\pm$ 1.0			Tiwari et al., (2010)	
Antwerp BEL	Ambient Indoor (5 d)*	2002-2003	0.4			0.1-0.7	Stranger et al., (2008)
	Ambient Outdoor (5 d)*						
Auvergne FRA	Ambient Indoor (1w)*	-		0.8	0.3-9.8	Hulin et al., (2010)	

\* passive sampling by Radiello; \*\* passive sampling by 3M OVM 3500; \*\*\* passive sampling by Carbopack B; \*\*\*\* active sampling; <sup>NA</sup> not available; (sampling period): h= hour; d= day; w= week.

Table 6. Levels of occupational and environmental exposure to benzene observed in the last 10 years.

Thanks to progressive improvements in the analytical techniques used to assay biomarkers since the 1970s, it has become possible to measure ever lower concentrations of chemical substances or their metabolites in human biological fluids after exposure to toxicants in the daily environment. For this reason, from being a tool used exclusively to assess occupational risks, biological monitoring has now become a valuable means of evaluating the risk of exposure to toxicants in the daily environment that integrates, but does not replace, environmental monitoring results. It thus could play a central role in public health and environmental medicine policies, helping to identify population groups at higher risk of exposure and/or effects, contributing to measure the internal dose and thus providing helpful information that can direct subsequent corrective measures (Angerer et al., 2007).

The results of biological monitoring in the general public are generally interpreted by comparison with reference values. These are obtained by statistical processing of the results of assays of the concentrations of a toxicant or its metabolites in biological fluids collected from a population or reference group with no occupational exposure to the substance (Apostoli & Minoia, 1999). It must be remembered, however, when assessing the risk of exposure to benzene, that unlike the threshold for air quality that derives from an estimate, albeit indirect, of the carcinogenic risk of exposure to benzene judged acceptable for the general population, the reference values do not bear any relation to the carcinogenic effect of benzene. Although the use of reference values does not therefore allow us to exclude a public health risk, it is important as a means of individuating specific population groups with higher levels of exposure to a toxicant than those of the reference population, due to residence in a highly polluted area, for instance, thus addressing one of the main objectives of biological monitoring in the non occupational field.

Many biological monitoring studies have been conducted to assess occupational exposure to benzene, but only few studies of the living environment. Moreover, these experiences have been gained largely in workers such as traffic wardens or public transport drivers who have occupational exposure to road traffic and so to generally higher concentrations of benzene than those affecting the general population (Barbieri et al., 2008; Campo et al., 2011; Fustinoni et al., 2005; Manini et al., 2008).

In this work the three main biological markers of internal dose currently in use, or under study for validation in biological monitoring of exposure to benzene, namely *t,t*-MA, SPMA and urinary benzene, were studied. Urinary *t,t*-MA is the only one of the three that is not entirely specific for benzene, since it can also derive from the metabolism of sorbic acid, an anti-mycotic preservative present in many foods, and so commonly absorbed in the diet, and in cosmetics. Although only a small percentage of the sorbic acid ingested is biotransformed to *t,t*-MA, the contribution of the diet can induce comparable concentrations of this metabolite to those observed in workers with occupational exposure to benzene levels equal to the TLV-TWA of ACGIH. Indeed, this phenomenon is growing proportionally as environmental exposure to benzene declines (Pezzagno et al., 1999). In 8 urine samples observed in the present population the concentrations of *t,t*-MA were above the upper limit of the reference range for the Italian population, equal to 15.2-163.1 µg/g creatinine (5<sup>th</sup>-95<sup>th</sup> percentile), recently defined in a multicentric national study (Aprea et al., 2010). However, the concentrations observed seem to be in line with the reference values for male subjects subdivided by smoking habit, since only 1 smoker and 3 non smokers had values exceeding the 95<sup>th</sup> percentile established for the relative group (274.7 µg/g creatinine and 117.8 µg/g creatinine, respectively).

The concentrations of *t,t*-MA, unlike those of the other markers of internal dose studied, were nearly always above the LOD, even in the non smokers. Moreover, the highest concentration, 734.0 µg/g creatinine, that is, in fact, higher than the BEI of 500 µg/g creatinine recommended by the ACGIH to control occupational exposure, was observed in a non smoker without exposure to urban traffic. This result could be attributed to the intake of sorbic acid in the diet. However, assessment of the contribution of sorbic acid to the *t,t*-MA levels excreted in the population studied was not possible because no data were available on the population's diet in the days before the study. Despite the reduced specificity of *t,t*-MA, cigarette smoke can condition urinary excretion of this marker, that shows a dependency relation on the number of cigarettes/day characterized by a R<sup>2</sup> of 0.22. This finding is in agreement with those by Pezzagno et al. (1999), who estimated that only 25% of urinary *t,t*-MA can be attributed to the biotransformation of benzene.

This dependency of urinary *t,t*-MA on the dietary intake of sorbic acid and the quantity of benzene absorbed in cigarette smoke explains the absence of a relation between *t,t*-MA and the concentrations of airborne benzene, and hence the poor validity of this marker as a means of monitoring exposure to very low concentrations of benzene like those observed in this study. Nevertheless, a significant correlation was found, when considering only the non smokers, between urinary *t,t*-MA and the time spent in urban traffic, in agreement with the report by Bergamaschi et al. (1999).

Unlike *t,t*-MA, SPMA is a highly specific metabolite of benzene, and in the occupational field it has been shown to be a valid biomarker of exposure to even low concentrations of this toxicant (Hoet et al., 2009). All the SPMA assays made in our sample, both adjusted and non adjusted for creatinine, were below the top limit of the reference values for the Italian population of 0.1-10 µg/L (5<sup>th</sup>-95<sup>th</sup> percentile) (Biolind, 2011), even if in 15 smokers and 2

non smokers the concentrations were higher than the 95<sup>th</sup> percentile of the values recently observed in a population with no occupational exposure resident in a northern Italian city (Milan) (Fustinoni et al., 2011). Cigarette smoke seems to be the only environmental factor causing such a high absorption of benzene as to condition the urinary excretion of SPMA, that was higher than the LOD practically only in smokers, yielding comparable results to those previously observed in a study of a population with no occupational exposure to benzene (Lovreglio et al., 2010). Unlike what was observed in subjects with occupational and non occupational exposure to higher concentrations of benzene, ranging from <3.0 to 66.5  $\mu\text{g}/\text{m}^3$ , in which SPMA was found to be dependent not only on the number of cigarettes smoked per day but also on the levels of airborne benzene, in subjects exposed to concentrations of less than 16.3  $\mu\text{g}/\text{m}^3$ , as in this study, SPMA was found to be conditioned only by a smoking habit (Lovreglio et al., 2011). Thus, urinary SPMA can be considered a very valid biomarker of internal dose when assessing exposure to low concentrations of benzene or excluding occupational exposure to benzene, but less useful for monitoring exposure to very low concentrations of benzene like those present in the living environment. Instead, thanks to its specificity and high sensitivity, urinary benzene has been proposed as a valid biomarker of even very low concentrations of benzene (Fustinoni et al. 2011). There are still some limits to the use of reference values to interpret the results of urinary benzene because the method for the pre-analytical phase of sample collection has not yet been standardized. This could explain why the top limit of the reference values for the Italian population of 0.05-1.45  $\mu\text{g}/\text{L}$  (5<sup>th</sup>-95<sup>th</sup> percentile) was exceeded in 14 determinations (Biolind, 2011), all in smoker subjects, whereas comparison with other recent data on a non occupationally exposed population showed values exceeding the 95<sup>th</sup> percentile only in 7 determinations, again all smokers (Fustinoni et al., 2011). In agreement with the data in literature, cigarette smoking is confirmed to be the main factor conditioning urinary benzene concentrations. In addition, unlike SPMA, it was dosable in 1/3 of the non smokers and showed a higher percentage of determinations above the LOD in subjects exposed to urban traffic than in those with no such exposure. Therefore, urinary benzene seems to be able to reflect even exposure to benzene due to pollution in urban areas, in agreement with what was previously observed in a population with exposure to higher concentrations of benzene attributable to the pollution of urban areas (Fustinoni et al., 2010).

Moreover, urinary benzene was found to be the only biomarker correlated to the concentrations of airborne benzene after excluding the contribution of cigarette smoking, in agreement with the reports by Fustinoni et al. (2010) and Campo et al. (2011). The concentrations of airborne benzene in the latter two studies were higher than those in our study (median values 4.0 and 9.6  $\mu\text{g}/\text{m}^3$  vs. <2.0  $\mu\text{g}/\text{m}^3$ ). The different trend of airborne benzene-urinary benzene correlations in smokers versus non smokers seems to confirm a confounding effect of the relation between exposure to airborne benzene and the biomarkers of internal dose due to a compounding effect of the benzene absorbed with cigarette smoke. This aspect was already emphasized in our previous experience as an important point to be taken into account when interpreting the results of biological monitoring to assess non occupational exposure to benzene.

## 5. Conclusion

Overall, our results seem to highlight that even when assessing the risk of non occupational exposure to benzene to exclude the possibility that a given population is exposed to higher

concentrations of benzene than the reference population, biological monitoring with different biomarkers of internal dose can usefully integrate environmental monitoring results. In this sense, the use of reference values can have a great importance, especially if smokers and non smokers are considered separately. In fact, in agreement with the findings in previous studies, cigarette smoking was shown to be the main non occupational source of benzene in smokers, that conditions the urinary excretion of all the biomarkers studied. Moreover, our results seem to confirm that even in conditions of exposure to very low benzene concentrations there is a competitive inhibitory effect of cigarette smoking on the metabolism of this toxicant, that induces proportionally higher concentrations of urinary benzene in smokers than of the metabolites such as *t,t*-MA. This can be explained by the presence of high quantities of chemical substances in cigarette smoke, that are partly metabolized through the same biotransformation pathways as benzene, especially via the oxidation of CYP2E1 (van Vleet et al., 2001).

Among the biomarkers analyzed in this study, urinary benzene seems to be the most sensitive and so the only one with a dependency relation not only on cigarette smoking but also on the time spent in urban traffic, as well as being the only one found to be correlated with the concentrations of airborne benzene, after excluding the effect of cigarette smoking. Thus, urinary benzene can be considered the biomarker of choice for assessing exposure to benzene in the daily environment, especially in terms of the contribution of urban traffic, but it is essential to take a smoking habit into account when interpreting the results of such studies.

## 6. References

- Agency for Toxic Substances and Disease Registry. (2007). *Toxicological profile for benzene*. ATSDR, US Department of Health and Human Services, Public Health Service, Atlanta, GA, US
- Al Zabadi, H., Ferrari, L., Sari-Minodier, I., Kerautret, M.A., Tiberguent, A., Paris, C., Zmirou-Navier, D. (2011). Integrated exposure assessment of sewage workers to genotoxicants: an urinary biomarker approach and oxidative stress evaluation. *Environmental Health*, 10, pp.23
- American Conference of Governmental Industrial Hygienists. (2010). *Threshold limit values and biological exposure indices*. ACGIH, ISBN 978-1-607260-19-6, Cincinnati, OH, US
- Angerer, J., Ewers, U. & Wilhelm, M. (2007). Human biomonitoring: state of the art. *International Journal of Hygiene and Environmental Health*, 210 (3-4), pp. 201-228
- Apostoli, P. & Minoia, C. (1999). I valori di riferimento in medicina occupazionale ed ambientale. *Giornale Italiano di Medicina del Lavoro ed Ergonomia*, 21(1), pp.25-39
- Aprèa, C., Sciarra, G., Bozzi, N., Pagliantini, M., Perico, A., Bavazzano, P., Leandri, A., Carrieri, M., Scapellato, M.L., Bettinelli, M. & Bartolucci, G.B. (2008). Reference Values of Urinary *Trans,trans*-muconic Acid: Italian Multicentric Study. *Archives of Environmental Contamination and Toxicology*, 55(2), pp.329-340
- Baan, R., Grosse, Y., Straif, K., Secretan, B., El Ghissassi, F., Bouvard, V., Benbrahim-Tallaa, L., Guha, N., Freeman, C., Galichet & L., Coglianò, V. on behalf of WHO International Agency for Research on Cancer Monograph Working Group. (2009). A review of human carcinogens-part F: Chemical agents and related occupations. *Lancet Oncology*, 10, pp.1143-1144

- Barbieri, A., Violante, F.S., Graziosi, F., Sabatini, L. & Mattioli, S. (2008). Urinary biomarkers and low-level environmental benzene concentration: assessing occupational and general exposure. *Chemosphere*, 74(1), pp.64-69
- Barrefors, G. & Petersson, G. (1995). Assessment by gas chromatography and gas chromatography-mass spectrometry of volatile hydrocarbons from biomass burning. *Journal of Chromatography A*, 710, pp.71-77
- Benedict, S.R. & Behre, J.A. (1936). Some application of a new color reaction for creatinine. *The Journal of Biology Chemistry*, 114, pp.515-532
- Bergamschi, E., Brustolin, A., De Palma, G., Manini, P., Mozzoni, P., Andreoli, R., Cavazzini, S. & Mutti, A. (1999). Biomarkers of dose and susceptibility in cyclists exposed to monoaromatic hydrocarbons. *Toxicology Letters*, 108(2-3), pp.241-247
- BIOLIND. (2011). In: *BIOLIND.NET*, Lista Valori di Riferimento SIVR 2005, last accessed 22.03.11, Available from <http://www.biolind.net/default.asp?nc=3346&id=161>
- Bolt, H.M. & Huici-Montagud, A. (2008). Strategy of the scientific committee on occupational exposure limits (SCOEL) in the derivation of occupational exposure limits for carcinogens and mutagens. *Archives of Toxicology*, 82(1), pp.61-64
- Boogaard, P.J. & van Sittert, N.J. (1995). Biological monitoring of exposure to benzene: a comparison between S-phenylmercapturic acid, trans, trans-muconic acid, and phenol. *Occupational and Environmental Medicine*, 52(9), pp.611-620
- Brief, R.S., Lynch, J., Bernath, T. & Scala, R.A. (1980). Benzene in the workplace. *American Industrial Hygiene Association Journal*, 41(9), pp.616-623
- Bruinen de Bruin, Y., Koistinen, K., Kephelopoulos, S., Geiss, O., Tirendi, S., Kotzias, D. (2008). *Environmental Science and Pollution Research International*, 15(5), pp.417-430
- Bruno, P., Caselli, M., De Gennaro, G., Iacobellis, S., Tutino, M. (2008). Monitoring of volatile organic compounds in non-residential indoor environments. *Indoor Air*, 18(3), pp.250-256
- Campo, L., Cattaneo, A., Consonni, D., Scibetta, L., Costamagna, P., Cavallo, D.M., Bertazzi, P.A. & Fustinoni, S. (2011). Urinary methyl tert-butyl ether and benzene as biomarkers of exposure to urban traffic. *Environment International*, 37(2), pp.404-411
- Caselli, M., De Gennaro, G., Marzocca, A., Trizio, L., Tutino, M. (2010). Assessment of the impact of the vehicular traffic on BTEX concentration in ring roads in urban areas of Bari (Italy). *Chemosphere*, 81(3), pp.306-311
- Collins, J.J., Ireland, B.K., Easterday, P.A., Nair, R.S., Braun, J. (1997). Evaluation of lymphopenia among workers with low-level benzene exposure and the utility of routine data collection. *Journal of Occupational and Environmental Medicine*, 39(3), pp.232-237
- Duarte-Davidson, R., Courage, C., Rushton, L. & Levy, L. (2001). Benzene in the environment: an assessment of the potential risks to the health of the population. *Occupational and Environmental Medicine*, 58(1), pp.2-13
- Edgerton, S.A. & Shah, J.J. (1992). Assessing total exposures to gasoline vapor using the source exposure model. *Journal of Exposure Analysis and Environmental Epidemiology*, 2(1), pp.109-115
- Estarlich, M., Ballester, F., Aguilera, I., Fernandez-Somoano, A., Lertxundi, A., Llop, S., Freire, C., Tardon, A., Basterrechea, M., Sunyer, J., Inguez, C. (2011). Residential Exposure to Outdoor Air Pollution during Pregnancy and Anthropometric


- Measures at Birth in a Multicenter Cohort in Spain. *Environmental Health Perspectives*, doi:10.1289/ehp.1002918
- European Commission. (2000). Directive 2000/69/EC of the European Parliament and of the Council of 16 November relating to limit values for benzene and carbon monoxide in ambient air. *Official Journal of the European Communities*, L313/12 of 13/12/2000
- European Parliament. (1999). Council Directive 1999/38/EC of 29 April 1999 amending for the second time Directive 90/394/EEC on the protection of workers from the risks related to exposure to carcinogens at work and extending it to mutagens. *Official Journal of the European Communities*, L138/66 of 01/06/1999
- Fruin, S.A., St. Denis, M., Winer, A.M., Colome, S.D. & Lurmann, F.W. (2001). Reductions in human benzene exposure in the California South Coast Air Basin. *Atmospheric Environment*, 35(6), pp.1069-1077
- Fustinoni, S., Consonni, D., Campo, L., Buratti, M., Colombi, A., Pesatori, A.C., Bonzini, M., Bertazzi, P.A., Foà, V., Garte, S., Farmer, P.B., Levy, L.S., Pala, M., Valerio, F., Fontana, V., Desideri, A. & Merlo, D.F. (2005). Monitoring low benzene exposure: comparative evaluation of urinary biomarkers, influence of cigarette smoking, and genetic polymorphisms. *Cancer Epidemiology, Biomarkers & Prevention*, 14(9), pp.2237-2244
- Fustinoni, S., Rossella, F., Campo, L., Mercadante, R. & Bertazzi, P.A. (2010). Urinary BTEX, MTBE and naphthalene as biomarkers to gain environmental exposure profiles of the general population. *Science of the Total Environment*, 408(14), pp.2840-2849
- Fustinoni, S., Campo, L., Mercadante, R., Consonni, D., Miekzynska, D. & Bertazzi, P.A. (2011). A quantitative approach to evaluate urinary benzene and S-phenylmercapturic acid as biomarkers of low benzene exposure. *Biomarkers*, DOI: 10.3109/1354750X.2011.561499
- Geiss, O., Tirendi, S., Barrero-Moreno, J. & Kotzias, D. (2009). Investigation of volatile organic compounds and phthalates present in the cabin air of used private cars. *Environment International*, 35(8), pp.1188-1195
- Ghittori, S., Fiorentino, M.L., Maestri, L., Cordioli, G. & Imbriani, M. (1993). Urinary excretion of unmetabolized benzene as an indicator of benzene exposure. *Journal of Toxicology and Environmental Health*, 38(3), pp.233-243
- Hattemer-Frey, H.A., Travis, C.C. & Land, M.L. (1990). Benzene: Environmental partitioning and human exposure. *Environmental Research*, 53(2), pp.221-232
- Hoet, P., De Smedt, E., Ferrari, M., Imbriani, M., Maestri, L., Negri, S., De Wilde, P., Lison, D. & Haufroid, V. (2009). Evaluation of urinary biomarkers of exposure to benzene: correlation with blood benzene and influence of confounding factors. *International Archives of Occupational and Environmental Health*, 82(8), pp.985-995.
- Holmberg, B. & Lundberg, P. (1985). Benzene: Standards, occurrence, and exposure. *American Journal of Industrial Medicine*, 7(5-6), pp.375-383
- Hsieh, L.T., Yang, H.H., Chen, H.W. (2006). Ambient BTEX and MTBE in the neighborhoods of different industrial parks in Southern Taiwan. *Journal of Hazardous Materials*, 128(2-3), pp. 106-115
- Hulin, M., Caillaud, D., Annesi-Maesano, I. (2010). Indoor air pollution and childhood asthma: variations between urban and rural areas. *Indoor Air*, 20(6), pp.502-514

- International Agency for Research on Cancer. (1982). Benzene, In: *Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. Vol. 29. Some industrial chemicals and dyestuffs.* pp. 93, IARC, Lyon (France)
- International Agency for Research on Cancer. (2004). *Monographs on the Evaluation of Carcinogenic Risks to Humans. Vol. 83. Tobacco Smoke and Involuntary Smoking*, IARC, ISBN 92 832 1283 5, Lyon (France)
- Iovino, P., Polverino, R., Salvestrini, S., Capasso, S. (2009). Temporal and spatial distribution of BTEX pollutants in the atmosphere of metropolitan areas and neighbouring towns. *Environmental Monitoring and Assessment*, 150(1-4), pp.437-444
- Italian Parliament. Law n. 245 of 5.03.1963. Limitazione dell'impiego del benzolo nelle attività lavorative. *Official Journal of the Italian Republic*, n. 77 of 21.03.1963, Ordinary Supplement n. 108. Istituto Poligrafico Zecca dello Stato, Rome
- Italian Parliament. Law n. 413 of 4.11.1997. Misure urgenti per la prevenzione dell'inquinamento atmosferico da benzene. *Official Journal of the Italian Republic*, n. 282 of 3.12.1997. Istituto Poligrafico Zecca dello Stato, Rome
- Kim, S., Vermeulen, R., Waidyanatha, S., Johnson, B.A., Lan, Q., Smith, M.T., Zhang, L., Li, G., Shen, M., Yin, S., Rothman, N. & Rappaport, S.M. (2006). Modeling human metabolism of benzene following occupational and environmental exposures. *Cancer epidemiology, Biomarkers & Prevention*, 15(11), pp.2246-2252
- Li, S., Chen, S., Zhu, L., Chen, X., Yao, C. & Shen, X. (2009). Concentrations and risk assessment of selected monoaromatic hydrocarbons in buses and bus stations of Hangzhou, China. *Science of the Total Environment*, 407(6), pp.2004-2011
- Liu, J., Mu, Y., Zhang, Y., Zhang, Z., Wang, X., Liu, Y., Sun, Z. (2009). Atmospheric levels of BTEX compounds during the 2008 Olympic Games in the urban area of Beijing. *Science of the Total Environment*, 408(1), pp.109-116
- Lovreglio, P., Barbieri, A., Carrieri, M., Sabatini, L., Fracasso, M.E., Doria, D., Drago, I., Basso, A., D'Errico, M.N., Bartolucci, G.B., Violante, F.S. & Soleo L. (2010). Validity of new biomarkers of internal dose for use in the biological monitoring of occupational and environmental exposure to low concentrations of benzene and toluene. *International Archives of Occupational and Environmental Health*, 83(3), pp.341-356
- Lovreglio, P., Cancanelli, G., Barbieri, A., Sabatini, L., D'Errico, M.N., Scicolone, L., Ghitti, R., Violante, F.S., Apostoli, P. & Soleo, L. (2010). Ruolo dei fattori non occupazionali nel condizionare i livelli di indicatori di dose interna utilizzati per monitorare l'esposizione occupazionale a concentrazioni molto molto basse di benzene. *Giornale Italiano di Medicina del Lavoro ed Ergonomia*, 32(1), pp.49-58
- Lovreglio, P., Carrieri, M., Barbieri, A., Sabatini, L., Fracasso, M.E., Doria, D., Iavicoli, S., Drago, I., D'Errico, M.N., Imbriani, M., Violante, F.S., Bartolucci, G.B. & Soleo, L. (2011). Applicability of urinary benzene to biological monitoring of occupational and environmental exposure to very low benzene concentrations. *Giornale Italiano di Medicina del Lavoro ed Ergonomia*, 33(1), pp.41-46
- Manini, P., De Palma, G., Andreoli, R., Poli, D., Petyx, M., Corradi, M., Mutti, A. & Apostoli, P. (2008). Biological monitoring of low benzene exposure in Italian traffic policeman. *Toxicology Letters*, 181(1), pp.25-30
- Massolo, L., Rehwagen, M., Porta, A., Ronco, A., Herbarth, O., Mueller, A. (2010). Indoor-outdoor distribution and risk assessment of volatile organic compounds in the

- atmosphere of industrial and urban areas. *Environmental Toxicology*, 25(4), pp.339-349
- Navasmrit, P., Chanvaivit, S., Intarasunanont, P., Arayasiri, M., Lauhareungpanya, N., Parnlob, V., Settachan, D., Ruchirawat, M. (2005). Environmental and occupational exposure to benzene in Thailand. *Chemico-biological Interactions*, 153-154, pp. 75-83
- Nomiyama, K. & Nomiyama, H. (1974). Respiratory elimination of organic solvents in man. Benzene, toluene, n-hexane, trichloroethylene, acetone, ethyl acetate and ethyl alcohol. *Internationales Archiv für Arbeitsmedizin*, 32(1), pp.85-91
- Ong, C.N., Kok, P.W., Lee, B.L., Shi, C.Y., Ong, H.Y., Chia, K.S., Lee, C.S. & Luo, X.W. (1995). Evaluation of biomarkers for occupational exposure to benzene. *Occupational and Environmental Medicine*, 52(8), pp.528-533
- Perez Ballesta, P., Field, R.A., Connolly, R., Cao, N., Baeza Caracena, A., De Saeger E. (2006). Population exposure to benzene: One day cross-sections in six European cities. *Atmospheric Environment*, 40, pp.3355-3366
- Pezzagno, G., Maestri, L. & Fiorentino, M.L. (1999). Trans,trans-muconic acid, a biological indicator to low levels of environmental benzene: some aspects of its specificity. *American Journal of Industrial Medicine*, 35(5), pp.511-518
- Roukos, J., Riffault, V., Locoge, N., Plaisance, H. (2009). VOC in an urban and industrial harbor on the French North Sea coast during two contrasted meteorological situations. *Environmental Pollution*, 157(11), pp.3001-3009
- Sabatini, L., Barbieri, A., Indiveri, P., Mattioli, S. & Violante, F.S. (2008). Validation of an HPLC-MS/MS method for the simultaneous determination of phenylmercapturic acid, benzylmercapturic acid and o-methylbenzylmercapturic acid in urine as biomarkers of exposure to benzene, toluene and xylenes. *Journal of Chromatography B*, 863(1), pp.115-122
- Scarselli, A., Binazzi, A., Di Marzio, D. (2011). Occupational exposure levels to benzene in Italy: findings from a national database. *International Archives of Occupational and Environmental Health*, DOI 10.1007/s00420-011-0616-9
- Simon, V., Baer, M., Torres, L., Olivier, S., Meybeck, M., Della Massa, J.P. (2004). The impact of reduction in the benzene limit value in gasoline on airborne benzene, toluene and xylenes levels. *Science of the Total Environment*, 334-335, pp.177-183
- Snyder, R., Witz, G. & Gildstein, B.D. (1993). The toxicology of benzene. *Environmental Health Perspectives*, 100, pp.293-306
- Snyder, R. & Hedli, C.C. (1996). An overview of benzene metabolism. *Environmental Health Perspectives*, 10 (supp 6), pp.1165-1171
- Stranger, M., Potgieter-Vermaak, S.S., Van Grieken, R. (2008). Characterization of indoor air quality in primary schools in Antwerp, Belgium. *Indoor Air*, 18(6), pp.454-463
- Supelco. (2011). Volatile organic compounds (VOCs) chemically desorbed with CS<sub>2</sub>. In: *Application Note Radiello*, 22.03.2011, Available from [http://www.sigmaaldrich.com/etc/medialib/docs/Supelco/Application\\_Notes/radiello\\_d1\\_d6.Par.0001.File.tmp/radiello\\_d1\\_d6.pdf](http://www.sigmaaldrich.com/etc/medialib/docs/Supelco/Application_Notes/radiello_d1_d6.Par.0001.File.tmp/radiello_d1_d6.pdf).
- Tiwari, V., Hanai, Y., Masunaga, S. (2010). Ambient levels of volatile organic compounds in the vicinity of petrochemical industrial area of Yokohama, Japan. *Air Quality, Atmosphere, & Health*, 3(2), pp.65-75

- Van Vleet, T.R., Bombick, D.W. & Coulombe, R.A. Jr. (2001). Inhibition of human cytochrome P450 2E1 by nicotine, cotinine, and aqueous cigarette tar extract in vitro. *Toxicological Sciences*, 64(2), pp.185-191
- Wallace, L.A. (1989). Major sources of benzene exposure. *Environmental Health Perspectives*, 82, pp.165-169
- Wallace, L.A. (1995). Human exposure to environmental pollutants: A decade of experience. *Clinical and Experimental Allergy*, 25(1), pp.4-9
- World Health Organization. (1993). *Environmental Health Criteria 155. Benzene*. WHO, Geneva (Switzerland).
- World Health Organization. (2000). Benzene, In: *Air quality guidelines for Europe. Second Edition*. WHO, pp. 1-18, Copenhagen (Denmark), Available from [http://www.euro.who.int/\\_\\_data/assets/pdf\\_file/0017/123056/AQG2ndEd\\_5\\_2\\_benzene.pdf](http://www.euro.who.int/__data/assets/pdf_file/0017/123056/AQG2ndEd_5_2_benzene.pdf)

IntechOpen



## **Air Quality-Models and Applications**

Edited by Prof. Dragana Popovic

ISBN 978-953-307-307-1

Hard cover, 364 pages

**Publisher** InTech

**Published online** 09, June, 2011

**Published in print edition** June, 2011

Air pollution has been a major transboundary problem and a matter of global concern for decades. High concentrations of different air pollutants are particularly harmful to large cities residents, where numerous anthropogenic activities strongly influence the quality of air. Although there are many books on the subject, the one in front of you will hopefully fulfill some of the gaps in the area of air quality monitoring and modeling, and be of help to graduate students, professionals and researchers. The book is divided in five sections, dealing with mathematical models and computing techniques used in air pollution monitoring and forecasting; air pollution models and application; measuring methodologies in air pollution monitoring and control; experimental data on urban air pollution in China, Egypt, Northeastern U.S, Brazil and Romania; and finally, the health effects due to exposure to benzene, and on the influence of air pollutants on the acute respiratory diseases in children in Mexico.

### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Piero Lovreglio, Maria Nicolà D'Errico, Silvia Fustinoni, Ignazio Drago, Anna Barbieri, Laura Sabatini, Mariella Carrieri, Pietro Apostoli, Leonardo Soleo (2011). Assessment of Environmental Exposure to Benzene: Traditional and New Biomarkers of Internal Dose, *Air Quality-Models and Applications*, Prof. Dragana Popovic (Ed.), ISBN: 978-953-307-307-1, InTech, Available from: <http://www.intechopen.com/books/air-quality-models-and-applications/assessment-of-environmental-exposure-to-benzene-traditional-and-new-biomarkers-of-internal-dose>

**INTECH**  
open science | open minds

### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166  
[www.intechopen.com](http://www.intechopen.com)

### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](#), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen