

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,300

Open access books available

130,000

International authors and editors

155M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



The Role of Aquaporines in Brain Tumors

Tatsuo Sawada

*Department of pathology Tokyo Women's Medical University
Japan*

1. Introduction

As about 80% of the brain is water, cerebral edema, associated with various cerebral injuries is critical and crucial events as the accumulation of excessive fluids will cause fatal cerebral herniation.

Water molecular channels called aquaporines (AQP) which are small membrane-spanning proteins which are expressed at plasma membranes^{1,2}. There are thirteen AQPs which are divided by three groups. Recently AQPs plays roles of cell migration³ and proliferation⁴. In central nervous system, eight AQPs (1,3,4,5,6,8,9, and 11) are expressed. AQP4 is expressed in astrocytes and AQP1 in cholloid plexus, predominantly.

Recent researches elucidated the physiological roles of AQPs expands from water channels to other function associated with tumor biology. Firstly, AQPs were found in high grade malignant tumors^{5,6}. And the expression of AQPs was correlated with metastatic potentials of malignant tumors and made easier to tumor cell migrations³.

To elucidate the characterization of the expression of AQPs especially of AQP 1 and 4, and the relation of the expression of VEGF family and their receptors, we had investigated the expression of these proteins in various tumors of the central nervous system, immunohistochemically

In this review, we survey the physiological roles of AQP 1 and 4 in central nervous system tumor, using our data and recent reports.

2. Materials and methods

Thirty-one tumors and three cerebral tissue for control were used at our study until now^{7,8}. Astrocytomas were reclassified according to WHO classification⁹. Although we could not examine the histology of primary tumors, the histological type of metastatic carcinoma was confirmed by the charts. Non-neoplastic cerebral tissue was obtained from three patients who underwent cerebral resection for cerebral hemorrhage. No information about corticosteroid treatment including dexamemethasone, was obtained in this study.

3. Immunostainings

Tissue specimen were immunostained by ABC methods using antihum an AQP 1, antihum an AQP4 polyclonal antibody (H80, diluted 1:50; Santa Cruz Biotechnology, Santa Cruz, CA, USA), antihuman VEGF polyclonal antibody (C-1, diluted 1:1000; Santa Cruz Biotechnology) and antihuman MIB1-LI monoclonal antibody (diluted 1:100; DAKO, Kyoto Japan)

4. Results

In glioma, immunoreaction for AQP1 and 4 were recognized in tumor cells. Negative reactions for AQP 1 and 4 in tumor cells of grade 1 meningioma, medulloblastoma, neurocytoma, and lymphoma. Some of metastatic carcinoma cells – some of metastatic carcinoma cells were negative for AQP4 but a few tumor cells were positive for AQP1. Positive reactions for AQP 1 and 4 were recognized in reactive astroglia in the surrounded cerebral area of tumor. Immunoreactions for AQP 1 were recognized only endothelial cells in neovasculature of gliomas and grade 2 meningiomas but normal endothelial cells negative for AQP 1 or 4. In grade 2 meningiomas, immunoreactions for AQP1 were detected in tumor cells and endothelial cells in neovasculature. A positive correlation of distinctiveness for immunoreactions between VEGF-A and C were obtained of expression of AQPs of endothelial cells in neovasculature.

5. Discussion

Recent researches elucidated the physiological roles of AQPs expands from water channels to other functions associated with tumor biology. Firstly, AQPs were found in high grade malignant tumors. And the expression of AQPs was correlated with metastatic potentials of malignant tumors and made easier to tumor cell migrations. AQP 4 was first cloned from rat lung¹⁰, played major roles of pathogenesis of neuromyelitis optica¹¹.

In cerebral tissue, metastatic carcinoma show significant cerebral edema in the surrounded area. Reactive astrocytes show positive reactions for AQP1 and 4. The mechanism of AQPs were still unclear., but As AQP4 did not alter blood-brain barrier integrity or cerebral morphology¹² but deletion increased cerebral edema. Tail et al suggested the AQP4 played the elimination of cerebral edema using rat experimental SAH models¹³.

AQP4 was massively up regulated in high grade astrocytoma and the expression of VEGF families tended to show a positive relation to AQP1 and 4 in high grade astrocytoma and grade 2 meningioma. Neovasculature in high grade astrocytoma and the adjacent areas of metastatic carcinoma showed glomeruoid proliferation of neovasculature in which the physiological blood-brain barrier impaired, showing high VEGFs expression¹⁴. In meningiomas, significant peritumoral edema are associated frequently(50-92%)¹⁵. But a few studies are available for the expression of AQPs about meningioma, Ng et al. found increased expression of AQP4 with peritumoral edema in meningioma⁶. But all cases of meningiomas Ng et al studied were grade 1. No grade 2 or 3 meningioma was not included. Our report showed no expression of AQP4 in tumor cells of grade 1 meningiomas⁵. Cerebral edema in high grade glioma, metastatic carcinoma and meningioma was likely the result of the relationship between cerebral blood vessels and AQP4-mediated water transport.

AQP is expressed in the apical membrane of the choroid plexus and plays a role in forming CSF¹⁶ But Marton et al. reported over expression of AQP 1 in atypical meningioma of infancy¹⁷. Nagashima et al suggested the dural invasion of meningiomas facilitated

Expressions of AQP1 played some physiological roles of cystic formation of cholloid plexus tumors¹⁸. As we describes, Nielsen et al reported that normal cerebral endothelial cells were negative for AQP1 but highly expressed in the endothelial cells of peripheral area of injuries, and cultured endothelial cells without astrocytes are also positive for AQP1. These data suggested that astrocytes endofeet may regulates adjacent endothelial cells in the endothelial cells in normal cerebral capillaries o switch off endothelial AQP 1 expression.

Our study could obtain the positive reactions of AQP1 in high grade glioma and grade 2 meningiomas but not in benign tumors.

In tumor biology, AQP1 deletion in mice reduced tumor growth and migration of endothelial and astroglia¹⁹. The migration of cells are closely related tumor malignancy, The biological roles of AQP 1 for arachnoid cell differentiation were uncertain, but our results suggested AQPs played some roles of take a malignant potentials of meningiomas..

In this study we could not take no informations about cerebral edema. Further studies about the relations between cerebral edema and the expressions of AQPs of meningiomas are needed.




Fig. 1. Histological and immunohistochemical findings of glioblastoma.

Histological findings (upper left) and immunohistochemical staining for AQP4 (upper left and lower left and right). Positive findings for AQP4 are recognized in endothelial cells and tumor cells.




Fig. 2. Histological and immunohistochemical findings of metastatic adenocarcinoma. Histological findings (upper left) and immunohistochemical staining for AQP4 (upper left and lower left and right). Positive findings for AQP4 are recognized in endothelial cells and reactive astroglia.





Fig. 3. Immunohistochemical findings of atypical meningioma. Immunohistochemical staining for VEGF (upper left), for AQP1 (upper right and lower left) and for AQP4 (lower right). Positive findings for AQP4 in tumor cells and for AQP1 in tumor and endothelial cells are recognized.

6. References

- [1] Verkman AS (1996) Water transport across mammalian membranes *Am J Physiol* 48:C12-C30
- [2] Ishibashi K, Kuwahara M, Sasaki S: (2000) Molecular biology of aquaporins *Rev Physiol Biochem Pharmacol* 141: 1-32
- [3] Papadopoulos MC, Saadoun and Verkman AS: aquaporins and cell migration *Pflugers Arch-Eur J Physiol* (2008)456: 693-700
- [4] Hoque MO, Soria JC, Woo J et al: Aquaporin 1 is overexpressed in lung cancer and stimulates NIH-3T3 Cell proliferation and anchorage-independent growth *Am J Pathology* (2006) 168:1345-1353
- [5] Saadoun S, Papadopoulos MC, Davis DC et al: Aquaporin-4 expression is increased in oedematous human brain tumors *J Neurol Neurosurg Psychiatry* (2002) 72: 262-265
- [6] Warth A, Kroger S, Wolburg H (2004) redistribution of aquaporin-4 in human glioblastoma correlated with loss of agrin immunoreactivity from brain capillary basal laminae. *Acta Neuropathol(Berl)* 107:311-318
- [7] Sawada T, Kato Y, Kobayashi M: Expression of aquaporin-4 in central nervous system tumors. *Brain Tumor Pathol* (2007) 24:81-84
- [8] Sawada T, Kato Y, Kobayashi M: Expression of aquaporin 1 and 4 in meningiomas. An immunohistochemical study. *Jr. Tokyo Women's Medical University*(2011)81: E9-13
- [9] Astrocytic tumors. WHO Classification of Tumours of the central nervous system.(2007) 13-52. WHO
- [10] Hasegawa H, Sawaya R, Mohanam S et al : Molecular cloning of a mercurial-insensitive water channel expressed in selected water transporting tissue. *J Biol Chem* (1994) 269: 5497-5500
- [11] Graber DJ, Levy M, Kerr et al : Neuromyelitis optica pathogenesis and aquaporin 4 *J of Neuroinflammation* (2008) 5: 22
- [12] Saadoun S, Tait MJ, Reza et al: AQP4 gene deletion in mice does not alter blood-brain barrier integrity or brain morphology *Neuroscience* (2009) 161: 764-772
- [13] Tait MJ, Saadoun S, Bell BA et al: Increased brain edema in AQP4-null mice in an experimental model of subarachnoid hemorrhage. *Neuroscience* (2010) 167: 60-67
- [14] Papadopoulos MC, Saadoun, Davis et al: (2001) Emerging molecular mechanism of brain tumor edema *Br.J neurosurg* 15:101-108
- [15] Kalkanis SN, Carroll RS, Zhang J et al: Correlation of vascular growth factor messenger RNA expression with peritumoral vasogenic cerebral edema in meningiomas. *J neurosurg.*(1996) 85: 1095-1101
- [16] Oshio K et al: Reduced cerebrospinal fluid production and intracranial pressure in mice lacking choroid plexus water channel aquaporin-1 *FASEB J* (2006) 19:76-78
- [17] Marton E, Feletti, Basaldella L et al (2008) Atypical cystic meningioma overexpressing AQP1 in early infancy: case report with literature review *Acta Paediatr* 97: 1145-1149.
- [18] Longatti P, Basaldella L, Orvieto E et al: Aquaporins expression in choroid plexus tumours *Pediatr Neurosurg* (2006) 42: 228-33
- [19] Saadoun SM, Papadopoulos MC, Watanabe H, et al: (2005) Involvement of aquaporin-4 in astroglia cell migration and glial scar formation *J Cell Sci* 118: 5691-5698



Molecular Targets of CNS Tumors

Edited by Dr. Miklos Garami

ISBN 978-953-307-736-9

Hard cover, 674 pages

Publisher InTech

Published online 22, September, 2011

Published in print edition September, 2011

Molecular Targets of CNS Tumors is a selected review of Central Nervous System (CNS) tumors with particular emphasis on signaling pathway of the most common CNS tumor types. To develop drugs which specifically attack the cancer cells requires an understanding of the distinct characteristics of those cells. Additional detailed information is provided on selected signal pathways in CNS tumors.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Tatsuo Sawada (2011). The Role of Aquaporines in Brain Tumors, Molecular Targets of CNS Tumors, Dr. Miklos Garami (Ed.), ISBN: 978-953-307-736-9, InTech, Available from:
<http://www.intechopen.com/books/molecular-targets-of-cns-tumors/the-role-of-aquaporines-in-brain-tumors>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike-3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/), which permits use, distribution and reproduction for non-commercial purposes, provided the original is properly cited and derivative works building on this content are distributed under the same license.

IntechOpen

IntechOpen