

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,300

Open access books available

131,000

International authors and editors

160M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



DNA Repair in Pathogenic Eukaryotic Cells: Insights from Comparative Genomics of Parasitic Protozoan

Laurence A. Marchat^{1,2}, Mavil López-Casamichana³,
Esther Orozco³ and César López-Camarillo³

*¹Escuela Nacional de Medicina y Homeopatía-IPN,
Programa Institucional de Biomedicina Molecular*

²Programa de Doctorado en Biotecnología en Red-IPN

*³Universidad Autónoma de la Ciudad de México,
Posgrado en Ciencias Genómicas
México*

1. Introduction

Numerous external and internal DNA damaging agents can affect genetic material to produce single-strand breaks (SSB), double strand breaks (DSB), inter- and intra-strand cross-links in the form of cyclobutane pyrimidine dimers and (6-4)-photoproducts, oxidation and alkylation of bases, or formation of bulky chemical adducts. Cells possess several biological processes that act in a coordinated way to supervise DNA molecules and properly repair DNA lesions to minimize genetic information loss. This DNA repair system, which has been conserved throughout eukaryotes and prokaryotes evolution, includes various pathways that can be classified according to the type of DNA lesion they can restore: i) DSB, the most detrimental lesions of DNA, can be repaired by homologous recombination (HRR) and non-homologous end joining (NHEJ) pathways [Fleck & Nielsen, 2004]; ii) aberrant bases or nucleotides from a single strand DNA can be repaired by base excision repair (BER), nucleotide excision repair (NER) and mismatch repair (MMR) pathways using the complementary strand as template for DNA synthesis. BER mainly restores non-bulky lesions that result from bases alkylation, oxidation or deamination [Krokan et al., 1997]. The main task of NER pathway, which consists in two subpathways: global genome repair (GGR) to remove damage in the overall genome and transcription-coupled repair (TCR) to specifically repair the transcribed strand of active genes, is to eliminate photoproducts produced by ultraviolet (UV) light and other bulky lesions, such as inter- and intra-strand crosslinks [Prakash & Prakash, 2000]. MMR allows the removal of base mismatches and small insertion/deletion loops (IDL) that are formed during the replication process [Marti et al., 2002].

The genome of protozoan parasites is continuously subjected to the effects of antiparasitic drugs and host immune system attacks, which can affect its stability and therefore parasite survival. Thus, efficient DNA maintenance mechanisms are necessary to detect and accurately repair damaged nucleotides. The fully sequenced genome of the four major human pathogens described here provides new insights into parasite biology, including molecular features of

DNA repair mechanisms, and genome evolution. *Entamoeba histolytica* and *Giardia lamblia* (syn. *G. intestinalis*, *G. duodenalis*) are intestinal parasites that cause diarrheal diseases. *E. histolytica* is responsible for fulminating dysentery, bloody diarrhea, weight loss, fatigue, abdominal pain, which affect 50 million people and provoke 100,000 deaths in developing countries each year. In some cases, *E. histolytica* trophozoites can cross the intestinal wall and use the blood stream to reach different vital organs of the human body, usually the liver (but also the lungs, brain or spleen) to provoke liver abscesses, which can be fatal if untreated [Guo et al., 2007]. *G. lamblia* is another contributor to the enormous burden of diarrheal diseases with over 250 million symptomatic human infections per year worldwide. This anaerobic flagellated protozoa colonises and reproduces in the small intestine of several vertebrates, including human, causing giardiasis, commonly known as Beaver fever, which is characterized by diarrhea, excess gas, stomach or abdominal cramps, upset stomach, and nausea. Additionally, *Giardia* infection has an adverse impact on child linear growth and psychomotor development since the parasite causes iron-deficiency anemia, micronutrient deficiencies and growth retardation associated with diarrhea and malabsorption syndrome [Ankarklev et al., 2010]. Individuals become infected by *E. histolytica* and *G. lamblia* through ingesting or coming into contact with food, soil, or water that have been contaminated by the feces of an infected human or animal. *Plasmodium falciparum* is the protozoan parasite responsible for human malaria, which is one of the most severe infectious diseases with 240 million cases in 2009 and more than 1 million deaths in children each year in Africa alone. The presence of the parasite in red blood cells lead them to stick to blood vessels through a process called cytoadherence, which produce the obstruction of the microcirculation and dysfunction of multiple organs, typically the brain in cerebral malaria. Symptoms usually include fever and headache, in severe cases progressing to coma, and death (Kokwaro, 2009). Trichomoniasis caused by *Trichomonas vaginalis* is the most common nonviral sexually transmitted disease (STD) in the world [WHO, 1995]. It has long been recognized as a frequent cause of vaginitis in women and urethritis in men, but data now link it to cervical cancer and bad pregnancy outcomes [Cotch et al., 1997], as well as to an enhanced risk for human immunodeficiency virus transmission [Sorvillo & Kerndt, 1998]. Here, we combined the use of genomic approaches based on bioinformatic analysis of parasite genome sequence with the review of published reports to perform a comparative description of DNA repair machineries from *E. histolytica*, *G. lamblia*, *P. falciparum* and *T. vaginalis*, which cause high morbidity and mortality in many developed and developing countries.

2. DNA repair machineries in pathogenic eukaryotic cells

2.1 Identification of DNA repair machineries

In order to identify amino acids sequences of *E. histolytica*, *G. lamblia*, *P. falciparum* and *T. vaginalis* proteins related to DNA repair factors, we performed similarity searches in the Eupath database (<http://eupathdb.org/eupathdb/>) using the *Saccharomyces cerevisiae* DNA repair proteins from HRR, NHEJ, BER, NER and MMR machineries as probes [reviewed in Lopez-Camarillo et al., 2009]. Putative gene products were selected from BLAST analysis against each parasite database using the Blosum 62 scoring matrix and the following criteria: (i) at least 20% identity and 35% homology to the query sequence and (ii) e-value lower than 0.002, unless a portion of the protein showed a very strong similarity. All sequences, as well as the *E. histolytica* sequences obtained from previous work [López-Camarillo et al., 2009], were then verified by BLAST against *S. cerevisiae* and *Homo sapiens* databases to confirm their identity. Additionally, we also retrieved data from published reports about *G. lamblia*,

P. falciparum and *T. vaginalis* (Table 1). The absence of a given sequence in the table indicates that the corresponding gene was not identified in the parasite genome or that the sequence was too divergent to be detected by our *in silico* strategy.

None of the protozoan parasites studied here has the complete DNA repair pathways reported in yeast. HRR is the most conserved pathway suggesting that it is the mayor DSB repair pathway in these protozoan parasites. *E. histolytica*, *G. lamblia*, *P. falciparum* and *T. vaginalis* genomes contain most of the RAD52 epistasis group genes, although their functional relevance remains to be determined. Homologs for RAD50, RAD51, MRE11, RAD54 and RPA (lacking the RAD52 interacting domain) have been previously reported in *P. falciparum* [Voss et al., 2002; Malik et al., 2008]. In agreement with its participation in DNA repair, the *PfRad51* gene is overexpressed in the mitotically active schizont stage and in response to methyl methane sulfonate [Bhattacharyya & Kumar, 2003]. In *T. vaginalis*, RAD50 y MRE11 were previously published as components of the meiotic recombination machinery, although meiosis has not been observed in this organism [Malik et al., 2008]. Ramesh et al. [2005] and Malik et al. [2008] identified the *Rad50/Mre11*, *Rad52* and *Dmc1* genes involved in meiotic recombination machinery by HRR in *Giardia*. Intriguingly, *G. lamblia* and *P. falciparum* lack the *nsb1* homologue (*xrs2* in Yeast) that is a component of the MRN complex involved in DSB detection and 3' ssDNA tails conversion. Recently, we published the *E. histolytica* RAD52 epistasis group involved in HRR [Lopez-Casamichana et al., 2007, 2008]. Interestingly, RT-PCR assays evidenced that some genes were down-regulated, whereas others were up-regulated when DSB were induced by UV-C irradiation, which revealed an intricate transcriptional modulation of *E. histolytica* RAD52 epistasis group related genes in response to DNA damage. Particularly, *EhRad51* mRNA expression was 16-, 11- and 4-fold increased at 30 min, 3 h and 12 h, respectively. DNA microarrays assays confirmed the activation of *EhMre11*, *EhRad50*, and *EhRad54* genes at 5 min after DSB induction, suggesting that they represent early sensors of damage in HRR pathway [Weber et al., 2009]. Additionally, the molecular characterization of EhRAD51 showed that the presence of all the functional domains reported in yeast and human homologues. EhRAD51 was upregulated and redistributed from cytoplasm to the nucleus of trophozoites at 3 h after DNA damage and it was able to catalyze specific single-strand DNA (ssDNA) transfer to homologous double strand DNA (dsDNA) forming the three-stranded pairing molecule called D-loop structure, confirming that it is a *bonafide* recombinase in *E. histolytica* [Lopez-Casamichana et al., 2008].

G. lamblia and *P. falciparum* only have three of the eight factors of the NHEJ pathway (including the MNR complex also involved in HRR), which strongly suggest that they preferably use HRR to repair DSB. In contrast, almost all NEHJ pathway factors have been identified in *E. histolytica* and *T. vaginalis*, including the LIF1 ligase, RAD27 nuclease and MRE11/RAD50/NSB1 proteins. However, *E. histolytica* genome does not contain a homologous gene for KU80 subunit [López-Camarillo et al., 2009] and *T. vaginalis* lacks both *ku70* and *ku80* genes [Carlton et al., 2007]. As these proteins form a single KU complex that recognizes DSB sites and recruits other DNA repair factors, our findings could appear contradictory. The absence of conserved KU proteins has also been reported in *Encephalitozoon cunili* [Gill & Fast, 2007] and yeast [Hefferin & Tomkinson, 2005], thus it is possible that these organisms use highly divergent KU proteins to perform the NHEJ pathway.

The other key DNA repair mechanisms represented by BER, NER and MMR pathways operate to repair aberrant bases or nucleotides from a ssDNA using the complementary strand as template for DNA synthesis. As in *E. histolytica* [Lopez-Camarillo et al., 2009], the *G. lamblia* BER pathway appears to be largely incomplete, lacking *apn1*, *mag1*, *ogg1*, *rad10*, *mus81* and *mms4* genes. Both parasites live under oxygen-limiting conditions and have a

highly reduced form of mitochondria called mitosomes [Tovar et al., 1999, 2003]. Then the absence of OGG1 could indicate that they do not suffer oxidative damage to mitochondrial DNA. In contrast, *Plasmodium* Flap endonuclease-1 (PfFEN-1) and Pf DNA Ligase I (PfLigI) have enzymatic activities similar to other species [Gardner et al., 2002; Casta et al., 2008], indicating that BER pathway should be functional in this parasite although several components are lacking.

Most genes involved in NER pathway are represented in *E. histolytica* [Lopez-Camarillo et al., 2009], *G. lamblia*, *P. falciparum* and *T. vaginalis* genomes suggesting that this mechanism could be potentially active in these eukaryotic parasites. PfXPB/RAD25, PfXPG/RAD2 and PfXPD/RAD3 have been previously reported in *P. falciparum* [Gardner et al., 2002; Bethke et al., 2007; Casta et al., 2008]. Additionally, the overexpression of *EhDdb1*, *EhRad23* and *EhRad54* genes after UV-induced DNA damage in *E. histolytica* [Weber et al., 2009] suggested that these genes could be involved in chromatin remodeling complexes as their homologues in human and yeast. *E. histolytica*, *G. lamblia* and *T. vaginalis* have various *rad3* genes to form the NEF3 complex (RAD2, RAD3, RAD25) of the BER pathway. Particularly, we identified six *rad3* genes and an additional truncated gene in *T. vaginalis*. On the other hand, all the parasites studied here lack almost one of the components of the TFIIH complex subunits (TFB1, TFB2 or TFB3).

As in bacteria, *Drosophila melanogaster*, *H. sapiens* and many other organisms [Lisby & Rothstein, 2005], *E. histolytica*, *G. lamblia* [Ramesh et al., 2005], *P. falciparum* [Bethke et al., 2007] and *T. vaginalis* [Malik et al., 2008] have almost all *S. cerevisiae* MMR genes, including the components of the MUTS α (MSH2/MSH6) heterodimer, which strongly suggest that MMR could be an active DNA repair pathway in these parasites. Notably, *E. histolytica* and *P. falciparum* have two *msh2* genes. However, neither *E. histolytica* nor *P. falciparum* present the *msh3* gene that is required for the formation of the MUTS β (MSH2/MSH3) heterodimer. PfMSH2-1, PfMSH2-2, PfMSH6, PfMLH1 and PfPMS1 proteins potentially participating in MMR have been previously reported in *P. falciparum*. Inhibition of *PfMSH2-2* gene increased mutation rate and microsatellite polymorphism, indirectly demonstrating its relevance in MMR and microsatellite slippage prevention. Moreover, antimalarial drug resistance has been recently related to a defective DNA mismatch repair, mainly in PfMutL α content [Castellini et al., 2011], which demonstrated the relevance of this mechanism for the parasite biology.

Gene name	<i>E. histolytica</i>	<i>G. lamblia</i>	<i>P. falciparum</i>	<i>T. vaginalis</i>	<i>S. cerevisiae</i>
Homologous recombination repair (HRR) pathway					
<i>rad50</i>	C4M2L7	Q6WD96	C6KSQ6	A2FAD3	P12753
<i>mre11</i>	Q86C23 C4LVX7 C4M8N7*	Q86C19 A8BR27*	PFA0390w**	A2ECB0	P32829
<i>nbs1</i>	C4M874	-	-	A2DHF7	P33301
<i>rad51</i>	C4M4K4	Q86C21	Q8IIS8 Pf11_0087**	A2FXT7	P25454
<i>rad52</i>	C4M197	Q6WD95	-	-	P06778
<i>rad54</i> <i>rad54b</i>	C4LVM6 C4M7S7	-	Q8IAN4	A2FNE0	P32863
<i>rad51c</i> <i>rad57</i>	C4M5L7	-	-	A2GIB8	P38953 P25301

Gene name	<i>E. histolytica</i>	<i>G. lamblia</i>	<i>P. falciparum</i>	<i>T. vaginalis</i>	<i>S. cerevisiae</i>
<i>rad59</i>	-	-	-	-	Q12223
<i>exo1</i>	C4MBM5	A8BQ11	Q8IBK1	A2E2N7	P39975
<i>rpa1</i>	C4M8G6	-	Q9U0J0 Q8I3A1	A2G5D0	P22336
<i>rpa2</i>	C4LT79	-	-	-	P26754
<i>sgs1</i>	C4M4V5	A8BAJ1 A8B9Y0	Q8I2W7 Q8ILG5	A2DYY2	P35187
<i>rad24</i>	C4M5T7	-	-	A2D9F4	P32641
<i>hpr5</i>	-	-	Q8I3W6	A2F783	P12954
<i>rad17</i>	-	-	-	-	P48581
<i>ddc1</i>	-	-	-	A2F0Q2	Q08949
<i>mec3</i>	-	-	-	-	Q02574
Non homologous end joining (NHEJ) pathway					
<i>ku80</i>	C4MBG9	-	-	-	P32807
<i>ku70</i>	-	-	-	-	Q04437
<i>lif1</i>	-	-	-	-	P53150
<i>dnl4</i>	C4M5H3	-	-	A2DFX6	Q08387
<i>rad27</i>	C4M6G8	A8B672 D3KG58	Q7K734 Q8IJW1	A2GNP0	P26793
Base excision repair (BER) pathway					
<i>apn1</i>	-	-	Q9BMG7	-	P22936
<i>apn21</i>	-	A8BGE2	O97240	-	P38207
<i>mag1</i>	-	-	-	-	P22134
<i>ogg1</i>	-	-	Q8I2Y2	-	P53397
<i>ntg1</i>	C4M764 C4LYM7	-	Q8II68	A2DS55	P31378
<i>ung1</i>	C4LUV5	A8B632	Q8ILU6	A2GFQ7	P12887
<i>pcna</i>	C4M9R9	A8BIU1	P61074 Q7KQJ9	A2DQV2	P15873
<i>rad1</i>	C4LT01	D3KH96	Q8ID22	A2DS24	P06777
<i>rad10</i>	C4LW01	-	O96136	A2DBF5	P06838
<i>cdc9</i>	C4M5H3	A8BWV4	Q8IES4	A2DFX6	P04819
<i>mus81</i>	-	-	-	A2FKU9	Q04149
<i>mms4</i>	-	-	-	A2DHF7	P38257
Nucleotide excision repair (NER) pathway					
<i>rad2</i>	C4M0V9	-	O96154	A2GNP0	P07276
<i>rad3</i>	C4M8K7 C4M8Q4 C4M6T8	A8BYS3 A8B495	Q8I2H7	A2G2G8 A2E4I6 A2F1W2 A2DDD4 A2E1B9 A2ELX1 A2G2G9*	P06839

Gene name	<i>E. histolytica</i>	<i>G. lamblia</i>	<i>P. falciparum</i>	<i>T. vaginalis</i>	<i>S. cerevisiae</i>
<i>rad4</i>	-	-	-	-	P14736
<i>rad7</i>	-	-	-	-	P06779
<i>rad14</i>	-	-	-	-	P28519
<i>rad16</i>	-	A8BL62	Q8I4S6	A2D9P9	P31244
<i>rad23</i>	C4MAR5	D3KF29*	Q8IJS8	A2FM19	P32628
<i>rad25</i>	C4MA19	A8BMI7	Q8IJ31	A2DEA8	Q00578
<i>rad26</i>	C4MAR8	A8BK31	-	A2EXQ4	P40352
<i>rad28</i>	-	-	Q8IJ73	A2DZ24	Q12021
<i>ssl1</i>	C4LV67	A8BA50	Q8IEG6	A2ENQ3	Q04673
<i>tfb1</i>	C4LWV8	-	-	-	P32776
<i>tfb2</i>	C4MIG0	-	Q8I4Y8	A2E2N2	Q02939
<i>tfb3</i>	-	D3KH94	Q8I3Y3	-	Q03290
<i>tfb4</i>	C4M9E2	A8B6C2	Q8IDG5	A2EYI3	Q12004
Mismatch repair (MMR) pathway					
<i>mlh1</i>	C4M5R1	-	Q8IIJ0	A2EGR5	P38920
<i>msh2</i>	C4M9J9 B1N4L6	Q6WD97	Q8ILI9 COH4L8	A2EP54	P25847
<i>msh3</i>	-	-	-	-	P25336
<i>msh6</i>	C4M4T8	A8BC61	Q8I447	A2EA54	Q03834
<i>pms1</i>	C4LW71	A8B4I6	Q8IBJ3	A2G2B4	P14242

Table 1. Comparison of DNA repair machineries from *E. histolytica*, *G. lamblia*, *P. falciparum*, *T. vaginalis* and *S. cerevisiae*. * fragment, ** PlasmoDB database.

2.2 Conservation of DNA repair pathways

To investigate the degree of conservation of DNA repair pathways in protozoan parasites, we next determined the values of Smith-Waterman identity scores between *E. histolytica* proteins and their corresponding orthologues in *G. lamblia*, *P. falciparum* and *T. vaginalis* by BLAST analysis based in pairwise sequence alignments and calculated the mean value for each DNA repair machinery (Fig. 1). Data of the MNR complex which participate in HRR and NHEJ pathways were included in both mechanisms. DSB repair pathways were generally more conserved than Excision Repair mechanisms. Considering amino acids identity, mean values for HRR and NHEJ pathways were higher in *E. histolytica*/*P. falciparum* comparison, suggesting that *E. histolytica* machinery was closer to *P. falciparum* than to *G. lamblia* and *T. vaginalis* machineries. The comparison *E. histolytica*/*G. lamblia* evidenced that HRR is highly conserved between both parasites, whereas components of the other pathways were more divergent. In the case of *E. histolytica*/*P. falciparum* comparison, NHEJ appeared to be more conserved than HRR, while the identity of HRR and NHEJ factors was very similar in *E. histolytica*/*T. vaginalis*. In all the parasites, the RAD51 recombinase is the most conserved protein (51%, 58% and 64% when *E. histolytica* protein sequence was compared with *G. lamblia*, *P. falciparum* and *T. vaginalis* orthologues, respectively), which is consistent with its relevant role in HRR mechanism.




Fig. 1. Conservation of DNA repair pathways between *E. histolytica* and *G. lamblia* (A), *P. falciparum* (B) and *T. vaginalis* (C). Amino acids sequences from orthologous proteins were compared by Blast and the percentage of identity was determined through pair wise alignment of the most conserved region. Average identity of all pathways is indicated above each graph.

2.3 DNA repair activity in cell free lysates evidences the functionality of DNA repair proteins

Although insights about the activity of DNA repair proteins in protozoa have been mainly obtained from experimental evidence based in heterologous expression and characterization of recombinant proteins, some reports showed that DNA repair activity could be detected in whole cell extracts, supporting the notion that DNA repair pathways already operates *in vivo*. For instance, Haltiwanger et al., (2000) reported the characterization of an AP

endonuclease activity in a *P. falciparum* cell free lysate. Authors provide evidence for the presence of class II, Mg^{2+} -dependent and independent AP endonucleases in the extracts. Moreover, they detected that *Plasmodium* AP endonuclease(s) possessed a 3'-phosphodiesterase activity similar to those described in other class II AP endonucleases [Demple et al., 1986]. In a related study, it was reported that a *P. falciparum* lysate contained uracil DNA glycosylase, AP endonuclease, DNA polymerase, flap endonuclease, and DNA ligase activities [Haltiwanger et al., 2000]. In contrast, DNA repair activities in cell lysates have not been detected in *Entamoeba*, *Giardia* and *Trichomonas* parasites. These data remark the utility of cell free lysates to understand DNA repair pathways, and pointed out to the urgency to investigate endogenous DNA repair activities using whole cell extracts in parasites where no data is available.

3. Functional categorization of *Entamoeba histolytica* DNA repair genes

To define the putative functions of *E. histolytica* DNA repair genes in unrelated DNA repair processes, we investigated the functional diversity of genomic maintenance pathways using Gene Ontology (GO) annotations. Functional related gene groups were predicted by the David bioinformatic resources (<http://david.abcc.ncifcrf.gov/gene2gene.jsp>), using a functional classification tool which generates a gene-to-gene similarity matrix based in shared functional annotation using over 75,000 terms from 14 functional annotation sources, allowing the classification of highly related genes in functionally related groups. Results from this analysis revealed that a large number of DNA repair genes were miss-annotated in parasites genome databases (43%). However, our analysis clearly showed that the majority of these genes seems to participate in DNA repair related processes. Besides, 57% of genes were predicted to function in DNA repair related process. 11% of genes participates in DNA damage repair, and 18% and 8% have helicase and endonuclease functions, respectively (Fig. 2).




Fig. 2. **Functional categorizations of *E. histolytica* DNA repair genes.** Biological processes and molecular functions were determined using David software (<http://david.abcc.ncifcrf.gov/gene2gene.jsp>). Percentage of genes included in individual categories is given.

4. Duplicated genes: The case of *rad3*

Gene duplicates represent for 8-20% of the genes in eukaryotic cells, and the rates of gene duplication are estimated at between 0.2% and 2% per gene per million years. Gene duplications are one of the major motors in the evolution of genetic systems and may occur in homologous recombination, retrotransposition event, or duplication of an entire chromosome [Zhang, 2003]. Duplicated genes are believed to be a main system for the establishment of new gene functions generating evolutionary novelty [Long & Langley, 1993; Gilbert et al., 1997].

A detailed examination of **Table 1** revealed that several DNA repair genes are duplicated in protozoan parasites, while there is only one gene in yeast. For example, the HRR machinery includes two *rad51* genes in *P. falciparum*, two *rad54* and *mre11* genes in *E. histolytica* [Lopez-Casamichana et al., 2008], two *rpa1* genes in *T. vaginalis*, and two *sgs1* genes in *G. lamblia* and *P. falciparum*. We also identified two *rad27* genes in *P. falciparum* and *G. lamblia* NHEJ pathway, two *E. histolytica* *ntg1* and *P. falciparum* *pcna* genes in the BER pathway, as well as two *msh2* genes for the MMR pathway in *E. histolytica* and *P. falciparum*. But the most duplicated gene was the *rad3* gene from the NER mechanism, since there are three genes in *E. histolytica*, two in *G. lamblia* and six in *T. vaginalis*, whereas *P. falciparum* has only one *rad3* gene, alike yeast. Remarkably, gene duplication is evident for many other genes in *T. vaginalis* and reflexes the massive gene expansion inside the large genome of this pathogen [Hartl & Wirth, 2006]. In yeast, the RAD3 protein is involved in mitotic recombination and spontaneous mutagenesis, becoming essential for cell viability in the absence of DNA injury. Furthermore, this protein participates in the repair of UV-irradiated DNA via NER, and constitutes a subunit of RNA polII initiation factor TFIIH [Moriel-Carretero & Aguilera, 2010]. *S. cerevisiae* RAD3 is related to the *H. sapiens* XPD, also known as ERCC2. Defects in human XPD result in a wide range of diseases, including Xeroderma pigmentosum (XP), Cockayne's syndrome, and Trichothiodystrophy characterized by a wide spectrum of symptoms ranging from cancer susceptibility to neurological and developmental defects [Liu et al., 2008].

In order to describe the inferred evolutionary relationships among the most abundant duplicated gene found through the analysis of DNA repair machineries from the human pathogens studied here, we have undertaken a phylogenetic analysis of RAD3 helicase orthologues in *S. cerevisiae*, *E. histolytica*, *T. vaginalis*, *G. lamblia* and *P. falciparum*. We evaluated the minimum evolution of RAD3 proteins through the construction of Neighbor-Joining phylogenetic tree using the MEGA version 5.05 [Tamura et al., 2011]. The robustness was established by bootstrapping test, involving 500 replications of the data based on the criteria of 50% majority-rule consensus (**Fig. 3**). Two main branches that came from a common ancestor can be observed. On one branch, *T. vaginalis* RAD3 paralogues are clustered into two sister proteins pairs (A2E1B9 and A2ELX1, A2E4I6 and A2DDD4), that have each evolved from the same ancestor. Besides, *E. histolytica* C4M6T8 is closer to *T. vaginalis* A2E4I6 and A2DDD4, than to its own paralogues. The other branch supports *T. vaginalis* A2G2G8 that is closely related to yeast and *P. falciparum* RAD3 proteins that came off the same node. Interestingly, these two organisms only have one *rad3* gene. This branch also includes *E. histolytica* C4M8K7 and C4M8Q4 sister proteins pair. Intriguingly, the two *Giardia* RAD3 proteins have emerged from different nodes and appeared to be more related to orthologues from other species than to each other; particularly, the branch supporting *Giardia* A8B495 also includes *Trichomonas* A2E1B9 and A2ELX1, while *Giardia* A8BYS3 is on the other branch, isolated from the other proteins, such as *Trichomonas* A2F1W2, which suggested that these proteins have evolved early.




Fig. 3. **Phylogenetic relationships between RAD3 from *S. cerevisiae*, *E. histolytica*, *T. vaginalis*, *G. lamblia* and *P. falciparum*.** The unrooted tree was created with the MEGA 5.05 program using the Neighbor Joining algorithm based on ClustalW. Numbers above the tree nodes indicate the percentage of times that the branch was recovered in 500 replications.

5. Molecular organization of the MNR complex

The MRE11-RAD50-NBS1 (MRN) complex is considered to have an imperative function in DSB repair. This protein complex operates as DSB sensor, co-activator of DSB-induced cell cycle checkpoint signaling, and as a DSB repairs effector in both the HRR and NHEJ pathways [Taylor et al., 2010; Rass et al., 2009]. Additionally, it has also been found to associate with telomeres maintenance at the ends of linear chromosomes. MRE11 and RAD50 orthologues have been reported in all taxonomic Kingdoms. MRE11, RAD50, and XRS2 (the *S. cerevisiae* homologue of vertebrate-specific NBS1) were initially recognized through yeast resistance to DNA damage induced by UV light and X-rays and meiotic recombination studies [Ogawa et al., 1995]. To efficiently perform these functions, this complex has shown particular enzymatic roles. Biochemical experiments have revealed that the phosphoesterase domain of MRE11 works as both a single- and double-stranded DNA endonuclease, besides as 3'-5' dsDNA exonuclease [D'Amours & Jackson, 2002]. Furthermore, RAD50 and NBS1/Xrs2 are able to promote the activity of MRE11, in an ATP dependent manner [Paul & Gellert, 1998]. ATP binding by RAD50 stimulates the binding of the MR complex to 3' overhangs and, also, ATP hydrolysis is required to arouse the cleavage of DNA hairpins, inducing modification of endonuclease specificity via DNA relaxing [Paull & Gellert, 1998; de Jager et al., 2002].

In this chapter, we have identified the presence of *Mre11* and *Rad50* genes in the genome of *E. histolytica*, *T. vaginalis*, *G. lamblia* and *P. falciparum*. However, all analyzed pathogenic eukaryotic cells, with the exception of *E. histolytica*, lack the *Xrs2* homologue. The absence of a NBS1/*Xrs2* homologous sequence in the other parasites might seem antagonistic to the idea of the existence of an active MRN complex. However we cannot discard the possibility that these microorganisms use a very divergent NBS1 protein, or even that this third component could be unessential. In order to initiate the characterization of components of MRN complex in these parasites, we studied the structural and evolutionary relationships between MRE11, RAD50 and NBS1 through PSI-BLAST analysis in comparison to human and yeast orthologues. This program generates a weighted profile from the sequences detected in the first pass of a gapped-BLAST search and iteratively searches the database using this profile as the query, allowing the inclusion of sequences with e-value cut off higher than 0.01 [Alschult et al., 1997]. Using the e-value threshold as a similarity measure, we evidenced a close relation between putative EhMRE11, HsMRE11, ScMRE11, TvMRE11 and PfMRE11. Conversely, GIMRE11 turned out to be less similar to the others, being closer to *E. histolytica* and *T. vaginalis* proteins (Fig. 4). On the other hand, analysis of RAD50 orthologues exposed a great conservation of these proteins, since all e-value threshold were <0.0001. As we have previously reported, EhNBS1 is closer to its human homologue than yeast [Lopez-Casamichana et al., 2007].




Fig. 4. Individual protein relationships of MRN complex in pathogenic eukaryotic cells. Similarity was evaluated through PSI-BLAST analysis. The width of connecting lines indicates similarity level.

To better understand the functionality of MRN complex in these parasites, predicted amino acid sequences of RAD50 and MRE11 were compared through multiple alignment using ClustalW software (<http://www.ebi.ac.uk/clustalw/>). Reported functional and structural domains were surveyed using Prosite (<http://www.expasy.org/tools/scanprosite/>), Pfam (<http://www.sanger.ac.uk/Software/Pfam/>), SMART (<http://smart.emblheidelberg.de/>) and Motif Scan (<http://myhits.isb-sib.ch/cgi-bin/motifscan>) programs. For all studied parasites, our search revealed that the MRE11 orthologues contain the N-terminal Mn²⁺/Mg²⁺-dependent nuclease domain including the five conserved phosphoesterase motifs described in yeast protein [Hopkins & Paull, 2008]. Moreover, C-terminal DNA binding domains were also identified [Williams et al., 2007; D'Amours & Jackson, 2002] (Fig. 5A).

RAD50 proteins displayed sequence and organizational homology to structural maintenance of chromosome (SMC) family members that control the higher-order structure and dynamics of chromatin. The N-terminal Walker A and C-terminal Walker B nucleotide binding motifs, which associate one with another to form a bipartite ATP-binding cassette (ABC)-type ATPase domain, were predicted [Hopfner et al., 2000; Hopfner et al., 2001]. Furthermore, amino acids flanking Walker motifs form coiled-coil configurations that converge with the cysteine zinc hook (CysXXCys) motif [Hopfner et al., 2002] (Fig. 5B). In the interphase of Walker domains, there are two MRE11 binding sites. Formation of the stable MRE11-RAD50 complex is reached by each unit of the MRE11 dimer binding a RAD50 molecule at the intersection of its globular and coiled-coil domains [de Jager et al., 2001a]. Scanning force microscopy experiments have demonstrated that whereas the globular head of the Mre112Rad502 complex links with the ends of linear dsDNA, the two coiled-coil regions of RAD50 are stretchy "arms", and project outward away from the DNA [Hopfner et al., 2002].

The third member of the MRN complex is NBS1 protein that was only detected in *E. histolytica*, but not in *G. lamblia*, *P. falciparum* neither *T. vaginalis*. We have previously reported that EhNBS1 consists of an FHA domain and adjacent BRCT domains at its N-terminus [Lopez-Casamichana et al., 2007]. In *Homo sapiens*, the FHA domain binds phosphorylated threonine residues in Ser-X-Thr motifs present in DNA damage proteins, including CTP1 and MDC1. The BRCT domains in human NBS1 fix Ser-X-Thr motifs when the serine residue is phosphorylated. These phospho-dependent interactions are significant for recruiting repair machineries and checkpoint proteins to DNA DSBs [Lloyd et al., 2009; Williams et al., 2009]. In reconstitution studies, the affinity of MRE11-RAD50 for DNA and its nuclease activity is further enhanced by the addition of NBS1 [Paull & Gellert, 1999].

6. Molecular organization of the RAD51 recombinase

RAD51 recombinase is an essential protein in HRR pathway that catalyzes strand transfer between a broken DNA and its undamaged homologous strand, allowing damaged region to be repaired [Thacker, 2005]. Strand exchange reaction is initiated by RAD51-coating of ssDNA released from DSBs, to generate a nucleoprotein filament. This active thread binds the intact dsDNA substrate, searching and locating homologous sequences, and promoting DNA strand exchange in an ATP-dependent manner, forming a heteroduplex structure [Paques & Haber, 1999]. After DNA damage, RAD51 protein has been observed in nuclear complexes forming discrete foci, which are considered as the recombinational DNA repair




Fig. 5. Comparison of the amino acids sequence of MRE11, RAD50 and RAD51 proteins of *S. cerevisiae*, *E. histolytica*, *T. vaginalis*, *P. falciparum* and *G. lamblia*. (A). Functional and structural domains of MRE11 proteins. MRE11 phosphoesterase motifs I-V (black

rectangles) and DNA-binding domains (gray rectangles). **(B)**. Functional and structural domains of RAD50 proteins. Walker A (N terminus) and Walker B (C terminus) are marked in the ABC-ATPase domains (gray rectangles), while MRE11-binding sites flank Walker motifs (dark gray rectangles) and the CysXXCys hook domain (white rectangles) are labeled in the coiled-coil central regions. **(C)**. Functional and structural domains of RAD51 proteins. Polymerization motif (PM), as well as Walker A and B motifs, L1 and L2 regions and ATP cap appear as colored rectangles. Boxshade panels: black boxes, identical aa; grey boxes, conserved substitutions.

sites [Tashiro et al., 2000]. RAD51 proteins have been identified in *Trypanosoma brucei* and *Plasmodium falciparum* parasites, which perform HRR to switch the expression of genes encoding surface membrane glycoproteins and generate antigenic variation [Conway et al., 2002; Freitas-Junior et al., 2000]. Furthermore, recombinational rearrangements are responsible for amplification of the multidrug resistance *pfmdr1* gene in *P. falciparum* [Triglia et al., 1991] demonstrating the relevance of HRR to generate genomic versatility and plasticity in protozoan parasites. Molecular analysis and functional assays confirmed that recombinant EhRAD51 is a *bonafide* recombinase that is able to catalyze specific ssDNA transfer to homologous dsDNA forming the three-stranded pairing molecule called D-loop structure. In addition, *E. histolytica* RAD51 sequence conserves the typical architecture of RECA/RAD51 family members [Lopez-Casamichana et al., 2008]. Amino acid sequences multiple alignment of RAD51 orthologues from *E. histolytica*, *S. cerevisiae*, *T. vaginalis*, *G. lamblia* and *P. falciparum* revealed that all these proteins share functional and structural conserved motifs (**Fig. 5C**). Each of them contains the putative polymerization motif (PM), which tethers individual subunits to form quaternary assemblies in human RAD51 protein [Bell, 2005]. We also identified the ATPase Walker A or phosphate binding loop (P-loop) and Walker B motifs residues, the ssDNA binding loops L1 and L2, as well as the ATP stacking motif or ATP cap at the C terminus, which are essential for nucleofilament assembling and ATP hydrolysis in RAD51/RECA-like recombinases [Shin et al., 2003; Conway et al., 2004].

7. Conclusions

Protozoan parasites are continuously subjected to the effects of antiparasitic drugs and host immune system attacks, which can affect their genome stability and therefore, their survival. In order to maintain the integrity of their DNA molecules, parasites have developed several mechanisms that are efficient to detect and accurately repair damaged nucleotides. Bioinformatic analyses of fully sequenced genomes are useful to identify molecular machineries for DNA repair in protozoan parasites of clinical relevance such as *Entamoeba histolytica*, *Giardia lamblia*, *Plasmodium falciparum* and *Trichomonas vaginalis*, which have a world-wide distribution with a high prevalence in developing countries. The computational data presented here provide new information on the evolution of DNA repair proteins and their potential relevance for DNA damage response in these major human pathogens. Future directions would include functional assays, as well as protein expression and protein-protein interactions analysis for the most relevant proteins, in order to contribute to the further elucidation of mechanisms regulating genome integrity in these organisms.

8. Acknowledgments

Genes and predicted protein sequences were obtained from the Eupath database (<http://eupathdb.org/eupathdb/>) and Uniprot Knowledgebase (<http://www.uniprot.org/>). This work was supported by UACM (México), CONACyT (México), COFAA-IPN (Mexico), SIP-IPN (Mexico) and European Community grants.

9. References

- Alschult, S.F., Madden, T.L., Schaffer, A.A., Zang, J., Zang, Z., Miller, W., Lipman, D.J. (1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acid Research*, (September 1997), Vol.25, pp. 3389-3402, ISSN 0305-1048
- Ankarklev, J., Jerlstrom-Hultqvist, J., Ringqvist, E., Troell, K., Svard, S.G. (2010). Behind the smile: cell biology and disease mechanisms of Giardia species. *Nature Reviews Microbiology*, Vol.8, (June 2010), pp. 413-422. ISSN: 1740-1526
- Bethke, L., Thomas, S., Walker, K., Lakhia, R., Rangarajan, R., Wirth, D. (2007). The role of DNA mismatch repair in generating genetic diversity and drug resistance in malaria parasites. *Molecular and Biochemical Parasitology*, Vol. 155, No. 1, (September 2007), pp. 18-25. ISSN 0166-6851
- Bell, C.E. (2005). Structure and mechanism of Escherichia coli RecA ATPase. *Molecular Microbiology*, Vol.58, No.2, (October 2005, pp.358-366, ISSN 0950-382X
- Bhattacharyya, M.K., Kumar, N. (2003). Identification and molecular characterisation of DNA damaging agent induced expression of Plasmodium falciparum recombination protein PfRad51. *International Journal for Parasitology*, Vol.33, (October 2002), pp. 1385-1392. ISSN 0020-7519.
- Carlton, J.M., Hirt, R.P., Silva, J.C., Delcher, A.L., Schatz, M., Zhao, Q., Wortman, J.R., Bidwell, S.L., Alsmark, U.C., Besteiro, S., et al. (2007). Draft genome sequence of the sexually transmitted pathogen Trichomonas vaginalis. *Science*, Vol. 315, No. 5809, (January 2007), pp. 207-212. ISSN 0036-8075
- Casta, L.J., Buguliskis, J.S., Matsumoto, Y., Taraschi, T.F. (2008). Expression and biochemical characterization of the Plasmodium falciparum DNA repair enzyme, flap endonuclease-1 (PffEN-1). *Molecular and Biochemical Parasitology*, Vol. 157, No. 1, (January 2008), pp. 1-12. ISSN 0166-6851
- Castellini, M.A., Buguliskis, J.S., Casta, L.J., Butz, C.E., Clark, A.B., Kunkel, T.A., Taraschi, T.F. (2011). Malaria drug resistance is associated with defective DNA mismatch repair *Molecular and Biochemical Parasitology*, Vol. 177, No. 2 (June 2010), pp. 143-147. ISSN 0166-6851
- Conway, C., Proudfoot, C., Burton, P., Barry, J.D., McCulloch, R. (2002). Two pathways of homologous recombination in Trypanosoma brucei. *Molecular Microbiology*, Vol.45, No.6, (September 2002), pp. 1687-1700, ISSN 0950-382X
- Conway, A.B., Lynch, T.W., Zhang, Y., Fortin, G.S, Fung, C.W., Symington, L.S., Rice, P.A. (2004). Crystal structure of a Rad51 filament. *Nature structural & molecular biology*, Vol.11, No.8, (August 2004), pp. 791-796, ISSN 1545-9993

- Cotch, M.F., Pastorek, J.G., Nugent, R.P., Hillier, S.L., Gibbs, R.S., Martin, D.H. et al. (1997). *Trichomonas vaginalis* associated with low birth weight and preterm delivery. The Vaginal Infections and Prematurity Study Group. *Sexually transmitted diseases*, Vol. 4, No.6, (July 1997), pp. 353-360. ISSN0148-5717
- D'Amours, D., Jackson, S.P. (2002). The Mre11 complex: at the crossroads of DNA repair and checkpoint signalling. *Nature Reviews Molecular Cell Biology*, Vol. 3, (May 2002), pp. 317-327, ISSN 1471-0072
- de Jager, M., Dronkert, M.L., Modesti, M., Beerens, C.E., Kanaar, R., van Gent, D.C. (2001). DNA-binding and strand-annealing activities of human Mre11: implications for its roles in DNA double-strand break repair pathways. *Nucleic Acids Research*, Vol.29, (March 2001), pp. 1317-1325, ISSN 0305-1048
- de Jager, M., van Noort, J., van Gent, D.C., Dekker, C., Kanaar, R., Wyman, C. (2001a). Human Rad50/Mre11 is a flexible complex that can tether DNA ends. *Molecular Cell*, Vol.8, (November, 2001 pp. 1129-113, ISSN 1097-2765
- Demple B, Johnson A, Fung D. Exonuclease III and endonuclease IV remove 3' blocks from DNA synthesis primers in H₂O₂-damaged *Escherichia coli*. 1986 *Proceedings of the National Academy of Sciences of the United States of America*. Voln 83, No.20, (October 1986), pp. 7731-7735, ISSN 0027-8424
- Fleck, O., Nielsen, O. (2004) DNA repair. *Journal of Cell Science*, Vol. 117, Pt. 4, (February), pp. 515-517, ISSN: 0021-9533
- Freitas-Junior, L.H., Bottius, E., Pirrit, L.A., Deitsch, K.W., Scheidig, C., Guinet, F., Nehrbass, U., Wellems, T.E., Scherf, A. (2000). Frequent ectopic recombination of virulence factor genes in telomeric chromosome clusters of *P. falciparum*. *Nature*, Vol.407, No.6807, (October 2000), pp. 1018-1022, ISSN 0028-0836
- Gardner, M.J., Hall, N., Fung, E., White, O., Berriman, M., Hyman, R.W. et al. (2002). Genome sequence of the human parasite *Plasmodium falciparum*. *Nature*, Vol. 419, No. 6906, (October 2002), pp. 498-511. ISSN 0028-0836
- Gilbert, W., de Souza, S. J., Long, M. Y. (1997). Origin of genes. *Proceedings of the National Academy of Sciences of the United States of America*, Vol.94, No.15, (July 1997), pp. 7698-7703, ISSN 0027-8424
- Gill, E.E., Fast, N.M. (2007). Stripped-down DNA repair in a highly reduced parasite. *BMC Molecular Biology*, Vol. 8, (March), pp. 24. ISSN 1471-2199
- Guo, X., Houpt, E., Petri, W.A.Jr. (2007). Crosstalk at the initial encounter: interplay between host defense and ameba survival strategies. *Current Opinion in Immunology*, Vol.19, (August 2007), pp. 376-384, ISSN 0952-7915
- Haltiwanger, B.M., Karpinich, N.O, and Taraschi, T.F. (2000) Characterization of class II apurinic/apyrimidinic endonuclease activities in the human malaria parasite, *Plasmodium falciparum*. *Biochemical Journal*, Vol. 345, (January 2000), pp. 85-89, ISSN 0264-6021.
- Haltiwanger, B.M, Matsumoto Y., Nicolas E., Dianov G.L., Bohr V.A., Taraschi T.F. (2000) DNA base excision repair in human malaria parasites is predominantly by a long-patch pathway. *Biochemistry*. Voln. 39, No. 4, (February 2000), pp. 763-772, ISSN 0006-2960.

- Hartl, D.L., Wirth, D.F. (2006), Duplication, gene conversion, and genetic diversity in the species-specific acyl-CoA synthetase gene family of *Plasmodium falciparum*. *Molecular and Biochemical Parasitology*, Vol.150, No.1, (November, 2006), pp.10-24, ISSN 0166-6851
- Hefferin, M.L., Tomkinson, A.E. (2005). Mechanism of DNA double-strand break repair by non-homologous end joining. *DNA Repair (Amsterdam)*, Vol. 4, No. 6, (June 2005), pp. 639-648. ISSN 1568-7864
- Hopfner, K.P., Karcher, A., Shin, D.S., Craig, L., Arthur, L.M., Carney, J.P., Tainer, J.A. (2000) Structural biology of Rad50 ATPase: ATP-driven conformational control in DNA double-strand break repair and the ABCATPase superfamily. *Cell*, Vol.101, No.7, (June 2000), pp. 789-800, ISSN 0092-8674
- Hopfner, K.P., Karcher, A., Craig, L., Woo, T.T., Carney, J.P., Tainer, J.A. (2001) Structural biochemistry and interaction architecture of the DNA double-strand break repair Mre11 nuclease and Rad50-ATPase. *Cell*, Vol.105, No.4, (May 2001), pp.473-485, ISSN 0092-8674
- Hopfner, K.P., Craig, L., Moncalian, G., Zinkel, R.A., Usui, T., Owen, B.A. et al. (2002). The Rad50 zinc-hook is a structure joining Mre11 complexes in DNA recombination and repair. *Nature*, Vol.418, No. 6897, (August 2002), pp. 562-566, ISSN 0028-0836
- Hopkins, B.B., Paull, T.T. (2008). The *P. falciparum* mre11/rad50 complex promotes 5' strand resection at a DNA double-strand break. *Cell*, Vol.135, (October 2008), pp. 250-260, ISSN 0092-8674
- Kokwaro, G. (2009). Ongoing challenges in the management of malaria. *Malaria Journal*, Vol.8, Suppl 1, (October 2009), pp. S2. ISSN 1475-2875
- Krokan, H.E., Standal, R., Slupphaug, G. (1997). DNA glycosylases in the base excision repair of DNA. *The Biochemistry Journal*, Vol. 325, No.1, (July 1997), pp. 1-16. ISSN 0264-6021
- Lisby, M., Rothstein, R. (2005). Localization of checkpoint and DNA repair proteins in eukaryotes. *Biochimie*, Vol.87, No.7, (July), pp. 579-589, ISSN 0300-9084
- Liu H, Rudolf J, Johnson KA, McMahon SA, Oke M, Carter L, McRobbie AM, Brown SE, Naismith JH, White MF. (2008), Structure of the DNA repair helicase XPD. *Cell*, Vol.133, No.5, (May 2008), pp.801-12, ISSN 0092-8674
- Lloyd, J. et al. (2009) A supramodular FHA/BRCT-repeat architecture mediates Nbs1 adaptor function in response to DNA damage. *Cell*, Vol.139, No. , (October 2009), pp. 100-111, ISSN 0092-8674
- Long, M. Y., Langley, C. H. (1993). Natural selection and the origin of jingwei, a chimeric processed functional gene in *Drosophila*. *Science*, Vol.260, No.91, (April 1993), pp.91-5, ISSN 0036-8075
- López-Camarillo, C., López-Casamichana, M., Weber, C., Guillen, N., Orozco, E., Marchat, L.A. (2009). DNA repair mechanisms in eukaryotes: Special focus in *Entamoeba histolytica* and related protozoan parasites. *Infection, Genetics and Evolution*, Vol.9, No.6, (December 2009), pp. 1051-1056. ISSN 1567-1348.
- López-Casamichana, M., Orozco, E., Marchat, L.A., López-Camarillo, C., (2007). Insights in DNA repair and homologous recombination in *Entamoeba histolytica*: in silico

- characterization of EhMRN complex. *Proceedings of the 5th European Congress on Tropical Medicine and International Health*, Monduzzi Editore s.r.l, pp. 7-11 , Medimond, ISBN: 13602276, Italy
- López-Casamichana, M., Orozco, E., Marchat, L.A., López-Camarillo, C. (2008). Transcriptional profile of the homologous recombination machinery and characterization of the EhRAD51 recombinase in response to DNA damage in *Entamoeba histolytica*. *BMC Molecular Biology*, Vol.9, (April 2008), pp.35. ISSN 1471-2199
- Malik, S.B., Pightling, A.W., Stefaniak, L.M., Schurko, A.M., Logsdon, J.M. Jr. (2007). An expanded inventory of conserved meiotic genes provides evidence for sex in *Trichomonas vaginalis*. *PLoS One*, Vol.3, No.8, (August 2007), e2879. ISSN 1932-6203
- Marti, T.M., Kunz, C., Fleck, O. (2002). DNA mismatch repair and mutation avoidance pathways. *Journal of Cellular Physiology*, Vol.191, (April 2002), pp. 28-41. ISSN 0021-9541
- Moriel-Carretero M, Aguilera A. (2010). Replication fork breakage and re-start: New insights into Rad3/XPD-associated deficiencies. *Cell Cycle*, Vol.9, No.15, (August 2010), pp.2958-62, ISSN 1538-4101
- Ogawa, H., Johzuka, K., Nakagawa, T., Leem, S.H., Hagihara, A.H. (1995). Functions of the yeast meiotic recombination genes, MRE11 and MRE2. *Advances in Biophysics*, Vol.31, pp. 67-76. ISSN 0065-227X
- Paques, F., Haber, J.E. (1999). Multiple pathways of recombination induced by double-strand breaks in *Saccharomyces cerevisiae*. *Microbiology and Molecular Biology Reviews*, Vol.63, No.2, (June 1999), pp. 349-404, ISSN 1092-2172
- Paull, T.T., Gellert, M. (1999). Nbs1 potentiates ATP-driven DNA unwinding and endonuclease cleavage by the Mre11/Rad50 complex. *Genes & Development*, Vol.13, No.10, (May 1999). pp. 1276-1288. ISSN 0890-9369
- Paull, T.T., Gellert, M. (1998). The 30 to 50 exonuclease activity of Mre11 facilitates repair of DNA double-strand breaks. *Molecular Cell*, Vol. 1, (June 1998), pp. 969-979, ISSN 1097-2765
- Prakash, S., Prakash, L. (2000). Nucleotide excision repair in yeast. *Mutation Research*, Vol.451, No.1-2, (June 2000), pp. 13-24. ISSN 1383-5742
- Ramesh, M.A., Malik, S.B., Logsdon, J.M.Jr. (2005). A phylogenomic inventory of meiotic genes: evidence for sex in *Giardia* and an early eukaryotic origin of meiosis. *Current Biology*, Vol.15, (January 2005), pp. 185-191. ISSN 0960-9822
- Rass, E., Grabarz, A., Plo, I., Gautier, J., Bertrand, P. , Lopez, B.S. (2009). Role of Mre11 in chromosomal nonhomologous end joining in mammalian cells. *Nature Structural & Molecular Biology*, Vol.16, (August 2009), pp. 819-824, ISSN 1545-9985
- Shin, D.S., Pellegrini, L., Daniels, D.S., Yelent, B., Craig, L., Bates, D. et al. (2003). Full-length archaeal Rad51 structure and mutants: mechanisms for RAD51 assembly and control by BRCA2. *EMBO Journal*, Vol.22, (September 2003), pp. 4566-4576, ISSN 0261-4189


- Sorvillo, F., Kerndt, P. (1998). *Trichomonas vaginalis* and amplification of HIV-1 transmission. *The Lancet*, Vol.351, (January 1998), pp. 213-214. ISSN 0140-6736
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., Kumar, S. (2011). MEGA5: Molecular Evolutionary Genetics Analysis using Maximum Likelihood, Evolutionary Distance, and Maximum Parsimony Methods. *Molecular Biology and Evolution* (submitted), ISSN 0737-4038
- Tashiro, S., Walter, J., Shinohara, A., Kamada, N., Cremer, T. (2000). Rad51 accumulation at sites of DNA damage and in postreplicative chromatin. *Journal of Cell Biology*, Vol.150, No.2, (July 2000), pp. 283-291, INSS 1540-8140
- Taylor, E.M., Cecillon, S.M., Bonis, A., Chapman, J.R., Povirk, L.F., Lindsay, H.D. (2010). The Mre11/Rad50/Nbs1 complex functions in resection-based DNA end joining in *Xenopus laevis*. *Nucleic Acids Research*, Vol.38, (November 2009), pp. 441-454, ISSN 0305-1048
- Thacker, J. (2005). The RAD51 gene family, genetic instability and cancer. *Cancer Letters*, 2005 Mar 10; Vol. 219, No. 2, (March 2005), pp. 125-135, ISSN 0304-3835
- Tovar, J., Leon-Avila, G., Sanchez, L.B., Sutak, R., Tachezy, J., et al. (2003). Mitochondrial remnant organelles of *Giardia* function in iron-sulphur protein maturation. *Nature*, Vol. 426, No. 6963, (November 2003), pp. 172-176; ISSN 0028-0836
- Tovar J, Fischer A, Clark CG. (1999). The mitosome, a novel organelle related to mitochondria in the amitochondrial parasite *Entamoeba histolytica*. *Molecular Microbiology*, Vol. 32, No. 5, (June 1999), pp. 1013-1021. ISSN: 0950-382X
- Triglia, T., Foote, S.J., Kemp, D.J., Cowman, A.F. (1991). Amplification of the multidrug resistance gene *pfmdr1* in *Plasmodium falciparum* has arisen as multiple independent events. *Molecular and Cellular Biology*, Vol.11, No.10, (October 2010), pp. 5244-5250, ISSN 0270-7306
- Voss, T.S., Mini, T., Jenoe, P., Beck, H.P. (2002). *Plasmodium falciparum* possesses a cell cycle-regulated short type replication protein A large subunit encoded by an unusual transcript. *The Journal of Biological Chemistry*, Vol.277, No.20, (May 2002), pp. 17493-17501. ; ISSN 0021-9258
- Weber, C., Marchat, L.A., Guillen, N., López-Camarillo, C. (2009). Effects of DNA damage induced by UV irradiation on gene expression in the protozoan parasite *Entamoeba histolytica*. *Molecular & Biochemical Parasitology*, Vol. 164, No. 2, (April 2009), pp. 165-169. ISSN 0166-6851
- World Health Organization (WHO). (1995). Global Programme on AIDS Report 1995. Geneva: WHO; 1995. An overview of selected curable sexually transmitted diseases
- Williams, R.S., Dodson, G.E., Limbo, O., Yamada, Y., Williams, J.S., Guenther, G. et al. (2009). Nbs1 flexibly tethers Ctp1 and Mre11-Rad50 to coordinate DNA double-strand break processing and repair. *Cell*, Vol.139, No.1, (October 2009), pp. 87-99, ISSN 0092-8674
- Williams, R.S., Williams, J.S., Tainer, J.A. (2007). Mre11-Rad50-Nbs1 is a keystone complex connecting DNA repair machinery, double-strand break signaling, and the

chromatin template. *Biochemistry and Cell Biology*, Vol.85, (August 2007), pp. 509-520, ISSN 0829-8211

Zhang, J. (2003). Evolution by gene duplication: an update. *Trends in Ecology & Evolution*, (June 2003) Vol. 18, No. 6, pp.292-298, ISSN 0169-5347

IntechOpen

IntechOpen



Selected Topics in DNA Repair

Edited by Prof. Clark Chen

ISBN 978-953-307-606-5

Hard cover, 572 pages

Publisher InTech

Published online 26, October, 2011

Published in print edition October, 2011

This book is intended for students and scientists working in the field of DNA repair, focusing on a number of topics ranging from DNA damaging agents and mechanistic insights to methods in DNA repair and insights into therapeutic strategies. These topics demonstrate how scientific ideas are developed, tested, dialogued, and matured as it is meant to discuss key concepts in DNA repair. The book should serve as a supplementary text in courses and seminars as well as a general reference for biologists with an interest in DNA repair.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Laurence A. Marchat, Mavil López-Casamichana, Esther Orozco and César López-Camarillo (2011). DNA Repair in Pathogenic Eukaryotic Cells: Insights from Comparative Genomics of Parasitic Protozoan, Selected Topics in DNA Repair, Prof. Clark Chen (Ed.), ISBN: 978-953-307-606-5, InTech, Available from: <http://www.intechopen.com/books/selected-topics-in-dna-repair/dna-repair-in-pathogenic-eukaryotic-cells-insights-from-comparative-genomics-of-parasitic-protozoan>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen