

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,300

Open access books available

130,000

International authors and editors

155M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Retinoids and Bone

H. Herschel Conaway¹ and Ulf H. Lerner^{2,3}

¹*Department of Physiology and Biophysics, University of Arkansas for Medical Sciences,*

²*Department of Molecular Periodontology, Umeå University*

³*Center for Bone and Arthritis Research at Institute of Medicine, Sahlgrenska Academy at*

University of Gothenburg,

¹*USA*

^{2,3}*Sweden*

1. Introduction

Vitamin A (retinol) can be produced in the body by hydrolysis of retinyl esters or reduction of retinal. Liver and eggs, which are good animal sources of vitamin A, contain retinyl esters. Plant sources such as carrots and spinach contain pro-vitamin A carotenoids, which can be cleaved to retinal. Retinal, also called retinaldehyde, is interconvertible with retinol (Fig. 1). Retinal also serves as an intermediate in the irreversible production of all-trans-retinoic acid (ATRA), which is considered the major biologically active derivative of vitamin A [Moise *et al.* 2007, Chambon 1996]. Another important derivative of vitamin A is the visual chromophore 11-cis-retinal [Wald 1968]. Binding of 11-cis-retinal to proteins called opsins is the chemical basis of vision. Vitamin A formed from retinyl esters or carotenoids in the normal diet, or ingested in fortified foods or dietary supplements, is stored in the liver and transported to tissues as a complex bound to retinol binding protein [Moise *et al.* 2007].




Fig. 1. Formation of all-trans retinoic acid from vitamin A and structures of 9-cis retinoic acid and 11-cis-retinal.

Effects of retinoids are mediated primarily by two families of nuclear hormone receptors, retinoic acid receptors (RARs) and the retinoid X receptors (RXRs) [Bastein *et al.* 2004]. Each receptor family is made up of three isoforms (α , β and γ), produced by separate genes. RARs can be activated by ATRA and the isomer 9-*cis* retinoic acid (9-*cis* RA), while RXRs are activated by 9-*cis* RA. RARs form heterodimers with RXRs and these heterodimers and RXR homodimers function as transcription factors, activating RAREs in the promoter regions of target genes. Most retinol signaling in cells is thought to be mediated by ATRA binding RAR in RAR/RXR heterodimers [Mic *et al.* 2003]. It is still not clear if 9-*cis* RA is formed physiologically in cells and what role this isomer may play as a specific ligand for RXR [Mic *et al.* 2003]. Activated retinoid receptors function as transcription factors, activating specific RA response elements (RAREs) for transcriptional regulation of target genes [Bastein *et al.* 2004, Mic *et al.* 2003].

Peroxisome proliferator-activated receptors (PPARs), α , β/δ , and γ , represent another group of nuclear hormone receptors that forms heterodimers with RXR. PPAR/RXR heterodimers also function as transcription factors, activating specific response elements of target genes [Mangelsdorf and Evans 1995, Bocher *et al.* 2002, Wilson *et al.* 2000]. Recent studies have indicated that ATRA not only can bind RARs, but can serve as a ligand for PPAR β/δ as well [Berry *et al.* 2007]. Channelling of ATRA to RARs or PPAR β/δ is suggested to depend on the intracellular binding proteins, RA binding protein II or keratinocyte fatty acid binding protein 5, which deliver ATRA to either RAR or PPAR β/δ , respectively [Schug *et al.* 2007].

Vitamin A plays an essential role in numerous biological processes, including vision, cellular proliferation, differentiation, and apoptosis, organ development and function, and immunity [Moise *et al.* 2007, Mark *et al.* 2006, Mark *et al.* 2009]. Numerous malformations and impaired vision, growth, organ function, and reproduction have been noted with vitamin A deficiency. Vitamin A is used in developing countries to help correct vitamin A deficiency [Moise *et al.* 2007]. Supplementation with vitamin A has had an enormous worldwide impact, saving countless lives, at minimal costs per patient. There is also widespread use of vitamin A derivatives (retinoids) for treatment of various skin conditions, such as acne [Peck *et al.* 1979], and for different cancers, including acute promyelocytic leukemia (APL), Kaposi's sarcoma, head and neck squamous cell carcinoma, ovarian carcinoma, and neuroblastoma [Siddikuzzaman *et al.* 2010]. Use of retinoids has proved particularly useful for treatment of APL [Siddikuzzaman *et al.* 2010]. Pharmacological concentrations (10^{-7} - 10^{-6} M) of ATRA block repression by a PML- RAR α fusion protein that interferes with normal RAR α transcriptional regulation in APL.

Besides serving as an intermediate in retinoic acid formation, recent studies have shown that retinal (retinaldehyde) is present at biologically active concentrations in fat tissue, where it antagonizes PPAR- γ activity, inhibits adipogenesis, and improves insulin sensitivity [Ziouzenkova *et al.* 2007]. These observations suggest that retinal may be an additional vitamin A derivative that plays an important role as a mediator of biological processes.

Hypervitaminosis A in experimental animals has been linked to increased bone resorption, decreased bone mass, and increased fractures [Frankel *et al.* 1986, Hough *et al.* 1988, Johansson *et al.* 2002]. There are also case studies in humans showing that significantly increased intake of vitamin A increases bone resorption, causes hypercalcemia, and induces skeletal pain [Frame *et al.* 1974]. Supplementation of the diet with vitamins is a common occurrence in developed countries and there is presently debate over whether more modest

increases in vitamin A might promote skeletal abnormalities. Some studies have shown that increased vitamin A intake or elevated serum retinol levels [Michaelsson *et al.* 2003] are associated with an increased incidence of hip fracture or decreased bone mass [Melhus *et al.* 1998, Feskanich *et al.* 2002, Promislow *et al.* 2002]; however, other studies have shown no deleterious effect on bone mass or fracture risk, and, in some instances, protection from bone loss because of increased vitamin A has been reported [Ribaya-Mercado and Blumberg 2007, Caire-Juvera *et al.* 2009]. In studies where vitamin A analogues have been evaluated, decreases in bone mass have been reported following isotretinoin and acitretin usage in some instances, but a recent, large scale, case-control study has found no increased risk of fracture with these agents [Vestergaard *et al.* 2010].

2. Bone tissue

Skeletal tissue is comprised primarily of either cortical (compact) or trabecular (cancellous) bone. In the adult, approximately 80% of skeletal mass is cortical bone and 20% trabecular bone. Cortical bone is found in the shafts of long bones and the outer surfaces of all other bones in the body. It is organized around blood vessels into cylinders of consolidated bone called osteons, or Haversian systems. Unlike trabecular bone, cortical bone does not contain bone marrow. Trabecular bone is characterized as a network of thin spicules of bone ("spongy bone"), which is found in the interior of vertebrae, flat bones in the skull, the pelvis and sacrum, and the distal and proximal parts of long bones. Both cortical and trabecular bone are important for bone strength. In the adult skeleton, both types of bone are remodelled and modelled [Martin and Seeman 2008]. Modelling of bone is a process that changes the size and shape of bone either by bone resorption without subsequent bone formation, or bone formation without prior bone resorption. Remodelling of bone is the process by which old bone is replaced by new bone. It is initiated by bone resorption to remove damaged bone, followed by new bone formation in the area resorbed. Remodelling does not change the size or shape of the bones. Remodelling is more frequent in trabecular bone than in cortical bone, one of the reasons why a metabolic bone disease like postmenopausal osteoporosis affects primarily bones with proportionally increased amounts of trabecular bone, e.g. vertebrae, distal radius, and proximal femur. Remodelling is believed to be initiated by microcracks in the mineralized bone extracellular matrix and subsequent osteocytic apoptosis, which triggers osteoclast formation and resorption of the micro-damaged area [Martin and Seeman 2008]. The process causes release of "coupling factors" which attract and activate osteoblasts to form new bone under a canopy of bone lining cells [Martin and Sims 2005, Boyce *et al.* 2009, Martin *et al.* 2009]. When osteoblasts fill the resorption lacunae made by osteoclasts with new bone in the bone multicellular unit (BMU), remodelling is in balance and bone mass remains constant. This equilibrium involves bidirectional interactions between osteoclasts and osteoblasts to fine tune the balance between bone formation and bone resorption.

The skeleton is a support structure that serves as a reservoir for calcium, phosphate, and numerous other minerals, protects vital organs such as brain, heart, lung, and bone marrow, and recently has been implicated in glucose- and energy metabolism [Confavreux *et al.* 2009]. Three systemic hormones that play primary roles in calcium homeostasis and bone formation and resorption are parathyroid hormone (PTH), 1,25(OH)₂-vitamin D₃, and calcitonin. PTH and 1,25(OH)₂-vitamin D₃ stimulate bone resorption when serum calcium decreases, whereas calcitonin inhibits bone resorption when serum calcium is increased. Sex

hormones also play important roles, with increased bone resorption and decreased bone formation occurring when levels of either estrogen or testosterone are decreased. Other agents known to play significant roles in bone turnover are the thyroid hormones, glucocorticoids, follicle stimulating hormone, and the retinoids. Interestingly, there is also a great deal of cross talk between bone cells and the immune system. This occurs during both normal physiological remodelling and in pathological inflammatory conditions involving bone, like rheumatoid arthritis and periodontitis [Takayanagi 2009].

3. Osteoclast differentiation

Osteoclasts are multinucleated giant cells found on bone surfaces that stain for tartrate resistant acid phosphatase (TRAP). They are formed by fusion of mononucleated progenitor cells derived from hematopoietic myeloid stem cells, which also give rise to macrophages and dendritic cells. For fusion of osteoclast precursors to occur, they must be differentiated specifically along the osteoclastic lineage [Lorenzo and Horowitz 2008, Edwards and Mundy 2011]. Formation of osteoclasts is controlled by osteoblasts at the bone surface. Thus, osteoblasts are responsible for not only bone formation, but regulate bone resorption as well. In addition to hematopoietic cells, bone marrow also contains pluripotent mesenchymal cells which are able to support differentiation of osteoclasts [Askmyr *et al.* 2009]; however, mature osteoclasts do not form within bone marrow, but enter the circulation as mononucleated osteoclast progenitors and home to periosteal and endosteal tissues. The details of this attraction are not known at present, but stromal cell-derived factor-1 (SDF-1 or CXCL12) produced by osteoblasts and CXCR4 expressed by osteoclast progenitors may play important roles [Kollet *et al.* 2006]. To what extent the circulating osteoclast progenitor cells are primed in the bone marrow, or to what extent priming occurs in the periosteum and endosteum is also not known at present. Nor is it known if osteoclast formation in trabecular bone in the close vicinity of bone marrow is different from osteoclast formation in periosteal tissues, which are always some distance from bone marrow. Recently, one group has presented evidence for the existence of a unique periosteal macrophage - osteomac which not only can form osteoclasts, but is also able to control bone formation [Chang *et al.* 2008]. Increasing evidence indicates that osteoclast formation does not follow a common pathway and that osteoclasts are different in different parts of the skeleton [Everts *et al.* 2009, Henriksen *et al.* 2011].

Crude bone marrow cultures, or co-cultures of periosteal osteoblasts and purified osteoclast progenitors from either bone marrow or spleens, have shown that osteoclast formation requires close physical contact of progenitor cells with either stromal cells in the bone marrow or osteoblasts. Molecularly, it has been found that macrophage colony-stimulating factor (M-CSF) and receptor activator of nuclear factor κ B ligand (RANKL) are two products of stromal cells/osteoblasts that play key roles in regulating osteoclast formation. M-CSF supports progenitor cell proliferation and survival, and RANKL is responsible for progenitor differentiation to osteoclasts, rather than to macrophages or dendritic cells [Takayanagi 2009, Lorenzo and Horowitz 2008, Nakashima and Takayanagi 2009]. RANKL, which is a member of the TNF superfamily, functions by binding to RANK on osteoclast progenitor cells. Osteoprotegerin (OPG), a soluble protein released from stromal cells/osteoblasts, is another key factor regulating osteoclastogenesis. OPG functions as a decoy receptor for RANKL, blocking interaction between RANKL and RANK. The expression of RANKL is thought to be restricted to a relatively small number of cells: bone marrow stromal cells, osteoblasts,

periodontal ligament cells, synovial fibroblasts, T-, B- and NK-cells, while OPG appears to be ubiquitously expressed. It is the relative expression of RANKL and OPG which determines whether osteoclast formation will take place. Enhancement of the RANKL/OPG ratio in stromal cells or osteoblasts by hormones and cytokines causes stimulation of bone resorption. Following RANKL binding to trimeric RANK, association of the cytoplasmic tail of RANK with TNF receptor-associated factor 6 (TRAF 6) leads to activation of several kinases, including IKK β , MAPKs such as p38 and JNK, and PI3K (Fig. 2). Subsequently, transcription factors such as NF- κ B, AP-1, MITF, NFATc1 and Akt are activated. Within the NF- κ B family of transcription factors, p50 and p52 seem to be the most important subunits [Boyce *et al.* 2010]. RANK signaling leads to induction and activation of c-fos, which functions as a component of the transcription factor, activator protein-1 (AP-1). Mice with double knockouts of p50/p52 or with deletion of the c-Fos gene lack osteoclasts and exhibit osteopetrosis. The induction of the master transcription factor for osteoclastogenesis, NFATc1, is dependent on NF- κ B and c-fos signaling by RANK [Negishi-Koga and Takayanagi 2009]. In cooperation with RANK, immunoglobulin-like receptors associated with adaptor proteins harboring the immunoreceptor tyrosine-based activation motif (ITAM) activate phospholipase C γ , calcium signaling, and the formation of calcineurin required for activation of NFATc1 [Nemeth *et al.* 2011]. The activation of specific genes necessary for osteoclastogenesis and osteoclast function is regulated by NFATc1 cooperating with other transcription factors, such as AP-1, CREB, PU.1, and MITF. Important osteoclast genes include those encoding calcitonin receptor and TRAP, which serve as osteoclast markers, cathepsin K, which is involved in breakdown of bone matrix collagen, Atp6i and chloride channel-7, involved in acidification and dissolution of bone mineral crystals, and the integrins α_v and β_3 , which are important for attachments of osteoclasts to bone surfaces.




Fig. 2. Osteoclastogenic pathways, transcription factors, and genes stimulated by M-CSF and RANKL.

4. Retinoids and bone resorption in vitro

The effect of vitamin A on bone resorption has been studied the most thoroughly in rodent models. It has been known since the 1920's that excess vitamin A has effects on the skeleton. In classical experiments in 1952 employing cultured fetal mouse limb bones, Dame Honor Fell showed that hypervitaminosis A stimulated osteoclast formation and bone resorption [Fell and Mellanby 1952]. Thirteen years later, Larry Raisz showed that vitamin A could stimulate bone resorption in organ cultures of fetal rat long bones [Raisz 1965]. Since those early experimental reports, there have been additional studies in organ cultures of fetal rat long bones [Scheven and Hamilton 1990], fetal rat calvariae [Delaissé *et al.* 1988] and calvarial bones from both fetal [Delaissé *et al.* 1988] and newborn [Raisz *et al.* 1977, Togari *et al.* 1991, Kindmark *et al.* 95, Conaway *et al.* 1997] mice confirming the osteoclastogenic effects of vitamin A and derivatives. An investigation indicating that ATRA increases mRNA expression of RANKL in primary human osteoblast-like cultures and decreases mRNA expression and protein formation of OPG in human MG-63 osteosarcoma cells [Jacobson *et al.* 2004] led to the suggestion that bone resorption stimulated by the retinoid is due to an increased RANKL/OPG ratio; however, until recently, there had been no data from bone or bone culture systems showing how vitamin A or derivatives stimulated bone resorption. In a recent investigation, we have reported that stimulation of osteoclastogenesis and bone resorption by retinoids in neonatal mouse calvarial bone is due to increased RANKL mRNA and protein expression. Supporting evidence for the role of RANKL was the observation that exogenous OPG administration blocked retinoid induced calvarial bone resorption [Henning *et al.* 2011].

Resorption in the bone organ culture systems depends on increased osteoclast formation and/or enhanced osteoclast activity. Attempts to elucidate effects of vitamin A on mature osteoclasts have generated conflicting results. It has been reported that bone resorbing activity of rabbit osteoclasts incubated on dentine slices is enhanced by ATRA [Saneshige *et al.* 1995]. Additionally, the vitamin A derivative isotretinonin has been observed to increase activity of rat osteoclasts incubated on cortical bone slices [Lakkakorpi and Väänänen 1991]. In contrast, bone resorbing activity of embryonic chicken osteoclasts on either bovine cortical bone or sperm whale dentine has been reported to be decreased by ATRA [O'Neill *et al.* 1992]. Furthermore, degradation of bone particles incubated with chicken osteoclasts is said to be enhanced by retinol and ATRA [Oreffo *et al.* 1988], which differs from experiments using osteoclast-like cell lines from human giant cell bone tumors, where Colucci *et al.* have found that ATRA inhibits degradation of bone matrix in rat bone particles [Colucci *et al.* 1996].

Evaluations of the effects of vitamin A on osteoclast formation have also generated conflicting data. In human bone marrow cultures, ATRA has been reported to stimulate formation of osteoclast-like multinucleated cells that are incapable of resorbing bone [Thavarajah *et al.* 1991]. Formation of multinucleated osteoclasts unable to resorb bone has also been reported following ATRA treatment of bone marrow cultures from egg laying hens [Chiba *et al.* 1996]. Scheven and Hamilton [1990] found no effect of ATRA on osteoclast formation in rat bone marrow cultures; however, when Hata *et al.* [1992] stimulated osteoclast formation in rat bone marrow cultures with 1,25(OH)₂-vitamin D₃, they found inhibition with ATRA. Using an acyclic retinoid, geranylgeranoic acid, Wang *et al.* [2002] also found that osteoclast formation in mouse bone marrow cultures stimulated by 1,25(OH)₂-vitamin D₃ was inhibited. Inhibition by the retinoid was most likely due to

inhibition of osteoclast progenitor cells, since inhibition of osteoclast formation by geranylgeranoic acid was also observed in mouse bone marrow macrophage (BMM) cultures stimulated by RANKL. In agreement with these latter observations, we have recently shown that ATRA, as well as 9-cis retinoic acid, inhibit osteoclast formation in mouse bone marrow cell cultures stimulated with either 1,25(OH)₂-vitamin D₃ or PTH [Conaway *et al.* 2009], an effect associated with decreased expression of the osteoclast genes *Calcr*, *Acp5*, and *Catsk*, which code for the calcitonin receptor, TRAP, and cathepsin K, respectively. ATRA did not affect 1, 25(OH)₂-vitamin D₃ stimulated mRNA expression of *Rankl*, nor did the retinoid affect the decrease of *Opg* expression induced by 1,25(OH)₂-vitamin D₃, suggesting that the inhibitory effect was at the level of osteoclast progenitors rather than stromal cells. Osteoclast formation in both crude bone marrow cultures and spleen cell cultures stimulated by RANKL was also inhibited by ATRA and 9-cis retinoic acid. Moreover, osteoclast formation stimulated by RANKL in highly purified mouse BMM cultures was inhibited by ATRA and 9-cis retinoic acid, providing good evidence that the retinoids inhibited osteoclastogenesis by directly affecting osteoclast progenitor cells.

Inhibition of osteoclast progenitor cell differentiation by ATRA was due to inhibition of AP-1 and Nfatc1 pathways, but did not involve the NF-κB pathway [Conaway *et al.* 2009]. ATRA inhibited mRNA and protein expression of the transcription factors c-Fos and Nfatc1 induced by RANKL. RANKL also enhanced the mRNA expression of *Fra-1*, *Fra-2* and *JunB* in mouse BMM, but ATRA inhibited only *Fra-2*. The decrease of the macrophage transcription factor MafB caused by RANKL during osteoclastogenesis was blunted by ATRA, suggesting that ATRA arrested precursor cells at the macrophage stage. In agreement with these observations, it has been observed that ATRA inhibits Nfatc1 expression, translocation, and DNA-binding in RAW264.7 cells and mouse BMM [Balkan *et al.* 2011].

Mouse BMM express RARα and RARβ, but little RARγ. By use of different agonists and antagonists, it was demonstrated that ATRA induced inhibition of RANKL stimulated osteoclast formation in BMM was mediated by RARα [Conaway *et al.* 2009]. As discussed above, ATRA can also serve as a ligand for PPARβ/δ. However, ligands specific for this receptor did not mimic the inhibitory effect of ATRA.

Effects of retinoids on osteoclast formation have also been studied using osteoclast progenitor cells isolated from peripheral blood. Woods *et al.* [1995] found that monocytes from chicken peripheral blood spontaneously formed giant cells with osteoclast like features and that this process was inhibited by ATRA. Recently, it was shown that ATRA inhibited osteoclast formation in RANKL stimulated cultures of human CD14⁺ monocytes from peripheral blood [Hu *et al.* 2010]. In agreement with observations in RANKL stimulated mouse BMM, inhibition of osteoclast formation in human CD14⁺ cells was associated with decreased mRNA expression of osteoclastic genes such as *ACP5* and *CATSK*, and the transcription factor *NFATc1*, indicating that retinoids inhibit osteoclast formation by inhibiting osteoclast differentiation rather than fusion of differentiated progenitor cells. The authors attributed the inhibition to decreased expression of RANK mRNA and protein; however, it remains to be shown if this is the primary event.

5. Retinoids and bone resorption in vivo

In short term *in vivo* studies where rodents were treated with increased concentrations of retinoids (ATRA or Ro 13-6298), it has been reported that thinning of cortical bone due to significant stimulation of periosteal resorption occurs at the same time that cancellous bone

resorption is inhibited [Kneissel *et al.* 2005]. Furthermore, in a recent one week study where rats were fed increased vitamin A, increased cortical osteoclasts and cortical bone thinning were also observed; however, in these animals, endosteal osteoclasts disappeared because of impaired endosteal/marrow blood flow, which resulted in hypoxia and pathological endosteal mineralization [Lind *et al.* 2011]. This resulted in thinner, more brittle bones with little apparent affect on bone mass. Vitamin A and retinoids stimulate resorption in cultured fetal mouse and rat limb bones [Fell and Mellanby 1952, Raisz 1965, Scheven *et al.* 1990], but these studies are normally based on calcium release to medium and do not distinguish between cortical and cancellous bone breakdown. On the other hand, calvarial bone is considered to be a good model for periosteal resorption of cortical bone and it is established that vitamin A and ATRA are effective stimulators of osteoclastogenesis and bone resorption in calvarial bone [Raisz *et al.* 1977, Togari *et al.* 1991, Kindmark *et al.* 1995, Conaway *et al.* 1997]. As stated previously, retinoids are also good inhibitors of osteoclastogenesis in mouse bone marrow cell and BMM cultures. Thus, when comparing results of *in vitro* and *in vivo* experimental studies attempting to access the effects of increased vitamin A, it appears there may be good agreement regarding cortical osteoclast formation and function stimulated by vitamin A. In contrast, it seems a continuum of effects may occur in the endosteum, with increased vitamin A inhibiting endosteal osteoclast formation and function, and impaired blood flow and hypoxia promoting osteoclast death.

6. Conclusion

In developed countries, the diet is often supplemented with vitamin A and there is presently controversy over whether increased intake of vitamin A might promote skeletal fragility [Michaelsson *et al.* 2003, Melhaus *et al.* 1998, Feskanich *et al.* 2002, Promislow *et al.* 2002, Ribaya-Mercado and Blumberg 2007, Caire-Juvera *et al.* 2009]. If cortical bone thinning and suppression of osteoclastogenesis in cancellous bone play prominent roles in fracture incidence in humans following increased intake of vitamin A, this would be substantially different from conditions such as postmenopausal osteoporosis and glucocorticoid excess, where the increased incidence of fracture is due primarily to cancellous bone loss. These different paradigms for bone fragility may help explain some of the differing outcomes in studies evaluating vitamin A intake and fracture risk. It also seems possible that a duality of retinoid action on osteoclast precursors in cortical and cancellous bone might be manifest as different degrees of cortical resorption and cancellous inhibition, depending on other systemic and environmental factors. This would also affect bone mass and fracture risk, as would the development of hypoxia and death of endosteal osteoclasts. Our improving understanding of vitamin A action in bone cells is not only promising to be extremely valuable for future experimentation, but appears to warrant new evaluations of bone mass and fracture risk in patients with increased intake of vitamin A as well.

7. Acknowledgements

Studies performed in the author's laboratories were supported by grants from the Swedish Research Council, Swedish Dental Society, the Royal 80 Year Fund of King Gustav V, the Swedish Rheumatism Association, Medical Faculty at Umeå University, Combine, ALF/LUA grants from the Sahlgrenska University Hospital and the County Council of Västerbotten.


8. References

- Askmyr M, Sims NA, Martin TJ, Purton LE. (2009) What is the true nature of the osteoblastic hematopoietic stem cell niche? *Trends Endocrinol Metab* 20:303-309
- Balkan W, Rodríguez-Gonzales M, Pang M, Fernandez I, Troen BR. (2011) Retinoic acid inhibits Nfatc1 expression and osteoclast differentiation. *J Bone Miner Metab* Epub ahead of print 2011
- Bastien, J., and Rochette-Egly, C. (2004) Nuclear retinoid receptors and the transcription of retinoid-target genes. *Gene* 328, 1-16
- Berry, D.C., and Noy, N. (2007) Is PPAR β/δ a retinoid receptor? *PPAR. Res.* 73256:1-5
- Bocher, V., Pineda-Torra, I., Fruchart, J.C., and Staels, B. (2002) PPARs: transcription factors controlling lipid and lipoprotein metabolism. *Ann. N. Y. Acad. Sci.* 967:7-18
- Boyce BF, Yao Z, Xing L. (2009) Osteoclasts have multiple roles in bone in addition to bone resorption. *Crit Rev Eukaryot Gene Expr* 19:171-180
- Boyce BF, Yao Z, Xing L. (2010) Functions of nuclear factor κ B in bone. *Ann NY Acad Sci* 1192:367-375
- Caire-Juvera, G., Ritenbaugh, C., Wactawski-Wende, J., Snetselaar, L., G. and Chen, Z. (2009) Vitamin A and retinol intakes and the risks of fractures among participants of the Women's Health Observational Study. *Am. J. Clin. Nutr.* 89: 323-330
- Chambon, P.(1996) A decade of molecular biology of retinoic acid receptors *FASEB J* 10, 940-954
- Chang MK, Raggatt LJ, Alexander KA, Kuliwaba JS, Fazzalari NL, Schroder K, Maylin ER, Ripoll VM, Hume DA, Pettit AR. (2008) Osteal tissue macrophages are intercalated throughout human and mouse bone lining tissues and regulate osteoblast function in vitro and in vivo. *J Immunol* 181:1232-44
- Chiba M, Teitelbaum SL, Cao X, Ross FP. (1996) Retinoic acid stimulates expression of the functional osteoclast integrin $\alpha_v\beta_3$: transcriptional activation of the β_3 but not the α_v gene. *J Cell Biochem* 62:467-475
- Colucci S, Grano M, Mori G, Scotlandi K, Mastrogiacomo M, Mori C, Zambonin Zallone A. (1996) Retinoic acid induces cell proliferation and modulated gelatinases activity in human osteoclast-like cell lines. *Biochem Biophys Res Commun* 227:47-52
- Conaway HH, Grigorie D, Lerner UH. (1997) Differential effects of glucocorticoids on bone resorption in neonatal mouse calvariae stimulated by peptide and steroid-like hormones. *J Endocrinol* 155:513-521
- Conaway HH, Persson E, Halén M, Granholm S, Svensson O, Pettersson U, Lie A, Lerner UH. (2009) Retinoids inhibit differentiation of hematopoietic osteoclast progenitors. *FASEB J* 23:3526-3538
- Confavreux CB, Levine RL, Karsenty G. (2009) A paradigm of integrative physiology, the crosstalk between bone and energy metabolism. *Mol Cell Endocrinol* 310:21-29
- Delaisé JM, Eeckhout Y, Vaes G. (1988) Bone-resorbing agents affect the production and distribution of procollagenase as well as the activity of collagenase in bone tissue. *Endocrinology* 123:264-276
- Edwards JR, Mundy GR. (2011) Advances in osteoclast biology: old findings and new insights from mouse models. *Nat Rev Rheumatol* EPUB ahead of print
- Everts V, de Vries TJ, Helfrich MH. (2009) Osteoclast heterogeneity: lessons from osteopetrosis and inflammatory conditions. *Biochim Biophys Acta* 1792:757-765

- Fell HB, Mellanby E. (1952) The effect of hypervitaminosis A on embryonic limb-bones cultivated in vitro. *J Physiol* 116:320-349
- Feskanich, D., Singh, V., Willett, W.C. and Colditz, G.A. (2002) Vitamin A intake and hip fractures among postmenopausal women. *JAMA* 287:47-54
- Frame, B., Jackson, C.E., Reynolds, W. A. and Umphrey, J. E. (1974) Hypercalcemia and skeletal effects in chronic hypervitaminous A. *Annals of Internal Medicine* 80:44-48
- Frankel, T.L., Seshadri, M.S., McDowall, D.B., and Cornish, C.J. (1986) Hypervitaminosis A and calcium-regulating hormones in the rat. *J. Nutr.* 116:578-587
- Hata K, Kukita T, Akamine A, Kukita A, Kurisu K. (1992) Trypsinized osteoclast-like multinucleated cells formed in rat bone marrow cultures efficiently form resorption lacunae on dentine. *Bone* 13:139-146
- Henning P, Pirhayati A, Pettersson U, Svensson O, Conaway HH, Lerner UH. (2011) Stimulation of mouse calvarial periosteal resorption and inhibition of osteoclast formation in human peripheral CD14⁺ cells by vitamin A. Abstract at 3rd Joint Meeting European Calcified Tissue Society and International Bone and Mineral Society, Athens May 7-11, 2011.
- Henriksen K, Bollerslev J, Everts V, Karsdal MA. (2011) Osteoclast activity and subtypes as a function of physiology and pathology – implications for future treatments of osteoporosis. *Endocr Rev* 32:31-63
- Hough, S., Avioli, L.V., Muir, H., Gelderblom, D., Jenkins, G., Kurasi, H., Slatopolsky, E., Bergfeld, M.A. and Teitelbaum, S. L. (1988) Effects of hypervitaminosis A on the bone and mineral metabolism of the rat. *Endocrinology* 122:2933-2939
- Hu L, Lind T, Sundqvist A, Jacobson A, Melhus H. (2010) Retinoic acid increases proliferation of human osteoclast progenitors and inhibits RANKL-stimulated osteoclast differentiation by suppressing RANK. *PLoS One* 5:e13305
- Jacobson A, Johansson S, Branting M, Melhus H. (2004) Vitamin A differentially regulates RANKL and OPG expression in human osteoblasts. *Biochem Biophys Res Commun* 322:162-167
- Johansson, S., Lind, P.M., Håkansson, H., Oxlund, H., Orberg, J., and Melhus, H. (2002) Subclinical hypervitaminosis A causes fragile bones in rats. *Bone* 31:685-689
- Kindmark A, Melhus H, Ljunghall S, Ljunggren Ö. (1995) Inhibitory effects of 9-cis and all-trans retinoic acid on 1,25(OH)₂ vitamin D₃-induced bone resorption. *Calcif Tissue Int* 57:242-244
- Kollet O, Dar A, Shivtiel S, Kalinkovich A, Lapid K, Sztainberg Y, Tesio M, Samstein RM, Goichberg P, Spiegel A, Elson A, Lapidot T. (2006) Osteoclasts degrade endosteal components and promote mobilization of hematopoietic progenitor cells. *Nat Med* 12:657-664
- Kneissel, M., Studer, A. and Cortesi, M. S. (2005) Retinoid-induced bone thinning by subperiosteal osteoclast activity in adult rodents. *Bone* 36, 202-214
- Lakkakorpi PT, Väänänen HK. (1991) Kinetics of the osteoclast cytoskeleton during the resorption cycle in vitro. *J Bone Miner Res* 6:817-826
- Lind, T., Lind, P.M., Jacobson, A., Hu, L., Sundqvist, A., Risteli, J., Yebra-Rodriguez, A., Rodriguez-Navarro, A., Andersson, G. and Melhaus, H. (2011) High dietary intake of retinol leads to bone marrow hypoxia and diaphyseal endosteal mineralization in rats. *Bone* 48, 496-506

- Lorenzo J, Horowitz M, Choi Y. (2008) Osteoimmunology: interactions of the bone and immune system. *Endocr Rev* 29: 403-440
- Mangelsdorf, D.J., and Evans, R.M. (1995) The RXR heterodimers and orphan receptors. *Cell* 83:841-850
- Mark, M., Ghyselinck, N.B., and Chambon, P. (2006) Function of retinoid nuclear receptors: lessons from genetic and pharmacological dissections of the retinoic acid signaling pathway during mouse embryogenesis. *Annu. Rev. Pharmacol. Toxicol.* 46: 451-480
- Mark, M., Ghyselinck, N.B., and Chambon, P. (2009) Function of retinoic acid receptors during embryonic development. *Nuclear Receptor Signaling* 7, e002
- Martin TJ, Seeman E. (2008) Bone remodeling: its local regulation and the emergence of bone fragility. *Best Pract Res Clin Endocrinol Metab* 22:701-722
- Martin TJ, Sims NA. (2005) Osteoclast-derived activity in the coupling of bone formation to resorption. *Trends Mol Med* 11:76-81
- Martin TJ, Gooi JH, Sims NA. (2009) Molecular mechanisms in coupling of bone formation to resorption. *Crit Rev Eukaryot Gene Expr* 19:73-88
- Melhus, H., Michaelsson, K., Kindmark, A., Bergstrom, R., Holmberg, L., Mallmin, H., Wolk, A. and Ljunghall, S. (1998) Excessive dietary intake of vitamin A is associated with reduced bone mineral density and increased risk of hip fracture. *Ann. Intern. Med.* 129:770-778
- Mic, F. A., Molotkov, A., Benbrook, D. M., and Duester, G. (2003) Retinoid activation of retinoid acid receptor but not retinoid X receptor is sufficient to rescue lethal defect in retinoic acid synthesis. *Proc. Natl. Acad. Sci. U. S. A.* 100, 7125-7140
- Michaelsson, K., Lithell, H., Vessby, B. and Melhus, H. (2003) Serum retinol levels and the risk of fracture. *N. Engl. J. Med.* 348:287-294
- Moise, A.R., Noy, N., Palcqwski, K., and Blaner, W.S. (2007) Delivery of retinoid-based therapies to target tissues. *Biochemistry* 46, 4449- 4458
- Nakashima T, Takayanagi H. (2009) Osteoclasts and the immune system. *J Bone Miner Metab* 27:519-529
- Negishi-Koga T, Takayanagi H. (2009) Ca²⁺-NFATc1 signalling is an essential axis of osteoclast differentiation. *Immunol Rev* 231:241-256
- Nemeth K, Schoppet M, Al-Fakhri N, Helas S, Jessberger R, Hofbauer LC, Goettsch C. (2011) The role of osteoclast-associated receptor in osteoimmunology. *J Immunol* 186:13-18
- O'Neill RPJ, Jones JS, Boyde A, Taylor ML, Arnett TR. (1992) Effect of retinoic acid on the resorptive activity of chick osteoclasts in vitro. *Bone* 13:23-27
- Oreffo RO, Teti A, Triffitt JT, Francis MJ, Carano A, Zallone AZ. (1988) Effect of vitamin A on bone resorption: evidence for direct stimulation of isolated chicken osteoclasts by retinol and retinoic acid. *J Bone Miner Res* 3:203-210
- Peck, G.L., Olsen, T.G., Yoder, F.W., Peck GL, Olsen TG, Yoder FW, Strauss JS, Downing DT, Pandya M, Butkus D, Arnaud-Battandier J.(1979) Prolonged remissions of cystic and conglobate acne with 13-cis-r retinoic acid . *N. Engl. J. Med.* 300: 329-333
- Promislow, J.H.E., Goodman-Gruen, D., Slymen, D.J. and Barrett-Connor, E. (2002) Retinol intake and bone mineral density in the elderly: The Rancho Bernardo study. *J. Bone Miner. Res.* 17:1349-1358
- Raisz LG. (1965) Bone resorption in tissue culture. Factors influencing the response to parathyroid hormone. *J Clin Invest* 44:103-116

- Raisz LG, Simmons HA, Gworek SC, Eilon G. (1977) Studies on congenital osteopetrosis in microphthalmic mice using organ cultures: impairment of bone resorption in response to physiologic stimulators. *J Exp Med* 145:857-865
- Ribaya-Mercado, J. D. and Blumberg, J. B. (2007) Vitamin A: Is it a risk factor for osteoporosis and bone fracture? *Nutr. Rev.* 65: 425-438
- Saneshige S, Mano H, Tezuka K, Kakudo S, Mori Y, Honda Y, Itabashi A, Yamada T, Miyata K, Hakeda Y, Ishi J, Kumegawa M. (1995) Retinoic acid directly stimulates osteoclastic bone resorption and gene expression of cathepsin K/OC-2. *Biochem J* 309:721-724
- Schug, T.T., Berry, D.C., Shaw, N.S., Travis, S.N. and Noy, N. (2007) Opposing effects of retinoic acid on cell growth result from alternate activation of two different nuclear receptors. *Cell* 129:723-733
- Siddikuzzaman, Guruvayoorappan, C. and Berlin Grace, V., M. (2010) All trans retinoic acid and cancer. *Immunopharmacol. and Immunotoxicol.* EPUB ahead of print 1-9
- Scheven BAA, Hamilton NJ. (1990) Retinoic acid and 1,25(OH)₂-vitamin D₃ stimulate osteoclast formation by different mechanisms. *Bone* 11:53-59
- Takayanagi H. (2009) Osteoimmunology and the effects of the immune system on bone. *Nat Rev Rheumatol* 5:667-676
- Thavarajah M, Evans DB, Kanis JA. (1991) 1,25(OH)₂D₃ induces differentiation of osteoclast-like cells from human bone marrow cultures. *Biochem Biophys Res Commun* 176:1189-1195
- Togari A, Kondo M, Arai M, Matsumoto S. (1991) Effects of retinoic acid on bone formation and resorption in cultured mouse calvaria. *Gen Pharmacol* 22:287-292
- Vestergaard, P., Rejnmark, L. and Mosekilde, L. (2010) High-dose treatment with vitamin A analogues and risk of fractures. *Arch. Dermatol.* 146:478-482.
- Wald, G. (1968) Molecular basis of visual excitation. *Science* 162, 230-239
- Wang X, Wu J, Shidoji Y, Muto Y, Ohishi N, Yagi K, Ikegami S, Shinki T, Udagawa N, Suda T, Ishimi Y. (2002) Effects of geranylgeranoic acid in bone: induction of osteoblast differentiation and inhibition of osteoclast formation. *J Bone Miner Res* 17:91-100
- Wilson, T.M., Brown, P.J., Sternbach, D.D., and Henke, B.R. (2000) The PPARs: from orphan receptors to drug discovery. *J. Med. Chem.* 43:527-550
- Woods C, Domenget C, Solari F, Gandrillon O, Lazarides E, Jurdic P. (1995) Antagonistic role of vitaminD₃ and retinoic acid on the differentiation of chicken hematopoietic macrophages into osteoclast precursor cells. *Endocrinology* 136:85-95
- Ziouzenkova, O., Orasanu, G., Sharlach, M., Adiyama T. E., Berger, J.P., Viereck, J., Hamilton, J. A., Tang, G., Dolnikowski, G. G., Vogel, S., Duester, G. and Plutzky, J. (2007) Retinaldehyde represses adipogenesis and diet-induced obesity. *Nature Med.* 13: 695-702



Contemporary Aspects of Endocrinology

Edited by Dr. Evanthia Diamanti-Kandarakis

ISBN 978-953-307-357-6

Hard cover, 454 pages

Publisher InTech

Published online 30, November, 2011

Published in print edition November, 2011

This book aims to provide readers with a general as well as an advanced overview of the key trends in endocrine disorders. While covering a variety of topics ranging from thyroid carcinogenesis and pituitary adenomas to adrenal tumors and metabolic bone disease, this book also focuses on more specific issues not yet fully elucidated (e.g. the molecular pathways involved in thyrotropin beta gene regulation or monogenic phosphate balance disorders). Readers of different fields and background will have the opportunity to update their knowledge and more importantly to clarify areas of uncertainty and controversies in several topics of endocrine disorders.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

H. Herschel Conaway and Ulf H. Lerner (2011). Retinoids and Bone, Contemporary Aspects of Endocrinology, Dr. Evanthia Diamanti-Kandarakis (Ed.), ISBN: 978-953-307-357-6, InTech, Available from: <http://www.intechopen.com/books/contemporary-aspects-of-endocrinology/retinoids-and-bone>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2011 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen