

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,200

Open access books available

129,000

International authors and editors

150M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# Exercise and the Immune System – Focusing on the Effect of Exercise on Neutrophil Functions

Baruch Wolach

*The Sackler School of Medicine, Tel Aviv University,  
Israel*

## 1. Introduction

A relationship between intense exercise, leukocytosis and susceptibility to illness was already reported at the beginning of the past century (1-3). Today there is a consensus among researchers and clinicians that exercise have effects on various aspects of the immune function (4). The complexity of the underlying mechanisms and the clinical implications and directions need continuous evaluation. Investigators face challenges associated with immune measures and the interpretation of their changes. They should bear in mind that there is inter-individual variability of the exercise capacity, recovery, stress tolerance and immunocompetence. Short exposure to exercise could promote beneficial and appropriate physiological response of the immune system, while heavy exertion could be detrimental to health. In recent years, the development of advanced laboratory techniques contributed to enrich our knowledge and deepened the understanding of the mechanisms underlying the immune system in sports medicine. The development of fluorescent antibodies techniques allow identifying cell sub-types and receptors. Molecular technology and new cytokine methods of identification have permitted the detection of humoral factors present in the body at low concentrations, for short periods of time and to study the effect of exercise on gene expression profiles (5,6).

Studies on recreational and elite athletes should be systematic and well controlled in order to formulate evidence-based guidelines to preserve a balanced immune function.

## 2. The immune system

The immune response can be divided into innate, natural-non-adaptive immunity and acquired-adaptive immunity. Innate immunity is the first response to physical or chemical foreign agents and it occurs naturally and immediately, providing the first line of defense in early stages of the infection. The innate immunity is comprised of phagocyte cells, natural killer cells, soluble factors as the complement and acute phase proteins, as well as the mucosal immune responses. The acquired immunity occurs after an adaptive, specific response to a pathogen and involves the antigen-antibody response. It includes B and T lymphocytes and the immunoglobulines (7).

In the innate immunity phagocytes can recognize and act immediately against the foreign agent without prior exposure, while the adaptive immunity is characterized by a specific response to the infectious agent, becoming fully activated after a lag period. The innate mucosal defenses are the first line of defense against pathogens present at the mucosal surfaces. The 'Common Mucosal Immune System' is a network of organized structures that protect the oral cavity, the respiratory, the gastrointestinal and the urogenital systems. The major effector function of this system is the secretory IgA (8,9).

The adaptive immunity involves the action of specialized immune cells, as are the lymphocytes, which generate antibodies against specific microorganisms, killing them directly or activating other cells through the secretion of cytokines. This adaptive response generates memory which is the basis of the preventive immunization. Both systems of immunity, the innate and the acquired, work synergistically and are essential for an optimal function of the immune response. Phagocytes play an important role in the initiation of the adaptive response by presenting antigens and secreting cytokines that stimulate cells of the adaptive system.

Neutrophils (55-65% of blood leukocytes) and monocytes (5-10%) play an important role in innate immunity and provide a major defense system against microorganisms. They act as the first line of defense against infectious agents and are involved in the muscle tissue inflammatory response to exercise-induced injury (10). The multi-step phagocytic process is activated in response to invasion of foreign microorganisms and includes the rolling and adherence of neutrophils to the blood vessel endothelium, the diapedesis and chemotaxis towards the invading organism, the ingestion, degranulation and the oxidative burst, ending with killing of the pathogen (11). Experienced and well equipped laboratories in this specific field, have established the normal range values, based on a large number of subjects examined, in health and disease. Today it is possible to assess the different steps of the phagocytic process and to detect dysfunctions at each level (11-13).

### **3. Exercise and the immune response – Clinical implications**

The potential influence of exercise on the immune system could be beneficial, detrimental or neutral. The immune response depends on the type of the particular exercise, its intensity, volume and duration. While mild or moderate exercise was shown to be beneficial, acute intense or prolonged exercise elicits depression of several aspects of the immune response. The fitness level of the performers could also exert influence in their immunological response. It seems that there is a combination of physiological and psychological factors, known to exert their influences on the immune system. Appropriate interpretation of the immune response is vital for determining the clinical directions and the integral training program for each athlete.

The physical activity could affect one or all three arms of the immune system, the humoral, the phagocytic and the cellular arm. Eventually, dysfunction of one or more arms of the immune system could lead to the outburst of an infection. In general, the etiology of infections is usually of bacterial origin when the humoral or phagocytic arm is affected, while viral-parasitic infections are usually originated when the cellular arm is involved. Excessive, prolonged training and major competitions have been long considered factors affecting the susceptibility to infections in athletes (14-16), however, in shorter and less

competitive events infections are less common (7). Frequent illness has been associated with the overtraining syndrome in athletes (17-19). During heavy exertion could be an immune suppression that creates an 'open window' of decreased host protection. Bacteria or viruses may gain a foothold, increasing the risk of subclinical and clinical infections (17, 20). In team sports or in other sports where participants are in close physical contact before, during or after the sporting event, both the infected individual and the fellow sportsmen may become infected. Some infections may appear in clusters in the sports setting, such as gastroenteritis, herpes simplex, meningitis, viral hepatitis, skin infections, tonsillo-pharyngitis (21,22). A large number of viruses and bacteria can give rise of myocarditis that can be aggravated by physical exertion (15).

There is consistent data suggesting that male endurance athletes may develop after 1 to 2-wk period increased rates of Upper Respiratory Tract Infection (URTI), following marathon or ultramarathon race events (16,23,24). URTI appears to be the most common minor viral infection in athletes. The current consensus is that the cause of URTI in athletes is uncertain (4). There is today disagreement whether 'sore throats', frequently reported by athletes, are caused by infections or are a reflexion of other inflammatory stimuli mimicking URTI (25,26). Cytokines play an important role in modulating the immune function, inducing changes that increase the risk of infection or the appearance of inflammatory symptoms (27). The physician diagnosis of URTI is based on clinical symptoms and signs, rather than by determining the infectious etiology. In few studies the pathogen was identified as the usual respiratory pathogens associated with URTI in the general population (4). The salivary IgA concentrations and secretion rates have been shown to be significantly decreased in athletes with prolong high intensity exercise (28,29). We could hypothesize that their immunity is reduced with an increase tendency to develop URTI. Other markers of infection as antimicrobial proteins in saliva ( $\alpha$ -amylase, lactoferrin, and lysozyme) have been identified (26,30). Further, viral infections as URTI may lead to a debilitating state and an unexplained deterioration in athletic performance. Viral infections could run a protracted course of easy fatigability, myalgia and lethargy for weeks or even months (31). Additionally, it seems that athletes are more susceptible to develop Infectious Mononucleosis (32).

Infections of non-viral origin, as bacterial pneumonia, mycoplasma and Chlamydia myocarditis, sinusitis, etc., although uncommonly reported in athletes, could also develop following intense exercise (2,15,33). Athletes could aggravate the course of the disease during incubation periods of infections (34,35).

Neutrophils comprise the majority of circulating leukocytes and represent the early body's response in the battle against bacterial and fungal infections. Multi-factorial elements could be involved in the neutrophil behavior and in the immune responses to exercise, as neuro-endocrine mediators (36), corticosteroid release, interleukin production (37) and oxy-reduction processes associated with free radical production (38). Most studies show that of all subsets of circulating leukocytes, mainly neutrophils and lymphocytes, increase dramatically during exercise (39,40). The magnitude is related to the exercise intensity and duration, being more persistent with intense, prolonged exercise (40, 41). Neutrophil count may exhibit a biphasic response, characterized by an initial small increase, followed by a decline to resting values 30-60 minutes after the cessation of exercise. A delayed larger increase in neutrophil numbers could be observed

2 to 4 hours post-exercise (42). This leukocyte trafficking reflect recruitment into the circulation of neutrophils and could be related to hemodynamic changes as increased cardiac output, hyperthermia or could reflect changes in circulating stress hormones, particularly epinephrine and cortisol, released during exercise (42). Resting leukocyte number is generally normal in athletes, although long periods of high-volume training may be associated with long lasting suppression of circulating cell numbers, which may persist low over weeks (43). This may be attributed to migration of leukocytes out of the circulation to possible damaged skeletal muscle (44).

Disproportionate changes in lymphocyte subsets occur during exercise. Usually, during prolonged exercise the NK and CD8 T- cells increase far more than B cells and CD4 T- cells counts (40). Significant decline in the CD4:CD8 ratio was reported after 60 min of treadmill running. Although neutrophil counts may remain elevated for several hours after exercise cessation, lymphocyte number may decline below baseline values for up to 6 hours post exercise (7). Following vigorous exercise it was reported a transient fall of circulating natural killer (NK) cell count (45).

In summary, exercise increases neutrophil numbers and may reflect an appropriate response to exercise-induced stress rather than an impaired immunocompromised state. In contrast, the post-exercise decrease in the absolute lymphocyte counts, the NK decrease and the inversion of the CD4 to CD8 ratio, could indicate immunosuppression.

It is remarkable that no significant changes were reported in B-cell circulating lymphocytes and only local salivary IgA reduction was shown following intense, prolonged exercise. The relationship between the leukocyte dynamics and the clinical implications is still unclear.

#### **4. Immunological studies – Our experience**

Scant information exists on exercise-induced changes in the immune system among children. We investigated the effect of aerobic exercise on several aspects of cellular and humoral functions among 10-12 year-old highly trained female gymnasts and untrained girls (46). All girls were pre-pubertal. Venous blood samples were drawn before, immediately after and 24 h following 20 min of treadmill running (heart rate 170-180 beats.min<sup>-1</sup>). White blood cells' number rose significantly following exercise and remained elevated for 24 h. The increase in leukocyte number was due to an increase in granulocytes as well as an increase in lymphocytes and monocytes. While neutrophil count returned to basal values after 24 h, lymphocytes and monocytes number remained elevated 24 h following exercise. Exercise resulted in a significant elevation of T cell lymphocytes, T helpers, T suppressors and natural killer cells. All values returned to normal after 24 h. There were no changes in B cell lymphocytes following exercise (Table 1). Exercise had no effect on serum immunoglobulin's and sub-types of IgG {IgG1, IgG2, IgG3 and IgG4} (Table 2). No differences were observed between gymnasts and untrained girls (46). The changes observed were similar to those found in adults (40).

Our laboratory for leukocyte functions focused within the last 30 years on granulocyte functions and dysfunctions in healthy subjects and in patients suffering from recurrent, severe, opportunistic infections. For the last 15 years, our group focused on neutrophil functions following a single bout of submaximal aerobic exercise (47-50). Neutrophil

functions as chemotaxis, oxidative burst, and bactericidal activity were unaffected immediately post-exercise, however neutrophil chemotactic activity was found significantly decreased 24 h after the cessation of the exercise (Table 3) (47).

		Pre exercise		Immediately after exercise		24 h following exercise	
		X	SD	X	SD	X	SD
WBC	gymnasts	5561	596	*7046	1306	*6505	436
	untrained	6340	1043	*8263	1225	*6738	1150
Neutrophils	gymnasts	2710	349	*3554	692	3126	498
	untrained	3570	1000	*4658	1184	3490	805
Lymphocytes	gymnasts	1930	235	*2690	360	*2365	524
	untrained	2106	544	*2671	834	*2464	401
T cell	gymnasts	1590	493	*1860	682	723	340
	untrained	1413	219	*1953	357	1677	458
Th cells	gymnasts	943	251	*1030	248	1003	155
	untrained	792	197	*1059	303	992	334
Ts cells	gymnasts	663	278	* 851	410	776	205
	untrained	550	98	* 784	193	669	173
B cells	gymnasts	306	101	320	96	344	93
	untrained	295	56	329	92	350	103
Natural killer cells	gymnasts	161	132	* 256	119	231	124
	untrained	124	47	* 267	85	139	43
Monocytes	gymnasts	293	61	* 391	127	# 424	72
	untrained	373	82	* 462	125	375	18

\* significantly different from pre-exercise value in both groups

# significantly different from pre-exercise value in the gymnasts group only

Table 1. Changes in the cellular components of the immune system. WBC and lymphocyte subpopulations (cells/ $\mu$ l) following exercise in gymnasts and untrained girls (46).

	IgM (mg/dl)		IgA (mg/dl)		IgE (unit/ml)		IgG (mg/dl)		IgG-1 (mg/dl)		IgG-2 (mg/dl)		IgG-3 (mg/dl)		IgG-4 (mg/dl)	
	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD
pre exercise	207.0	83.3	120.6	26.1	81.6	53.2	1194.0	138.2	673.7	108.0	235.3	63.7	69.7	35.5	61.2	24.1
post exercise	199.3	67.6	120.4	21.6	81.1	49.8	1180.7	151.2	705.9	132.6	242.1	66.8	59.1	18.5	63.4	25.1
24 h post exercise	179.4	73.2	116.4	36.4	83.1	56.5	1181.4	147.3	669.9	144.3	232.4	69.0	63.6	22.1	61.4	24.2

Table 2. Immunoglobulin levels pre-exercise, immediately post-exercise among gymnasts (46).

		Pre-exercise (Basal)	Post- exercise	24 h post- exercise
Neutrophils (cells/ $\mu$ l)	trained	2710 $\pm$ 349	3554 $\pm$ 692	3126 $\pm$ 498
	untrained	3570 $\pm$ 1000	4658 $\pm$ 1184	3490 $\pm$ 805
Chemotaxis (cells/field)	trained	58 $\pm$ 11	55 $\pm$ 13	36 $\pm$ 11 *
	untrained	47 $\pm$ 7	52 $\pm$ 15	42 $\pm$ 8 *
Killing (log decrease of colonies) with autologous serum	trained	0.8 $\pm$ 0.3 #	0.8 $\pm$ 0.2 #	0.8 $\pm$ 0.1 #
	untrained	1.1 $\pm$ 0.1	1.1 $\pm$ 0.1	1.0 $\pm$ 0.2
Killing (log decrease of colonies) with homologous serum	trained	0.7 $\pm$ 0.2 #	0.8 $\pm$ 0.1 #	0.8 $\pm$ 0.2
	untrained	1.0 $\pm$ 0.1	1.0 $\pm$ 0.02	0.9 $\pm$ 0.2
Superoxide production (nmol O <sub>2</sub> /10 <sup>6</sup> PMNs/min) with PMA stimulation	trained	5.7 $\pm$ 0.4	4.4 $\pm$ 1.0 *	4.8 $\pm$ 1.0 *
	untrained	5.1 $\pm$ 0.7	4.7 $\pm$ 1.3	4.9 $\pm$ 1.2
Superoxide production (nmol O <sub>2</sub> /10 <sup>6</sup> PMNs/min) with fMLP stimulation	trained	3.4 $\pm$ 1.9	3.2 $\pm$ 1.3	2.6 $\pm$ 1.3
	untrained	4.1 $\pm$ 1.3	3.3 $\pm$ 0.5	3.7 $\pm$ 0.6

\*)<0.05 compared with basal values (before vs. following exercise).

#)<0.05 compared with control group (trained vs. untrained).

Table 3. Effect of exercise on neutrophil count and neutrophil functions (mean+SD), Pre-exercise (basal), immediate post-exercise and 24 h post-exercise (47).

A consistent decrease of neutrophil migration was detected 24 h post-exercise in trained and untrained subjects, children and adults, male and female (47-50). The following studies focused on the recovery time of the impaired neutrophil chemotaxis, using various chemoattractants. We also aimed to learn about the possible mechanisms involved in the post exercise-associated chemotactic defect. We found that the transient impairment shown in the chemotactic activity 24 h post-exercise, returned to normal after 48 h (Figure 1) (49).




Fig. 1. Kinetics of the neutrophil chemotactic activity in 16 athletes; Pre-exercise, immediately post-exercise and 24 h post-exercise. The chemotaxis was induced by the chemoattractant: fMLP (1  $\mu$ M), IL-8 (10 nM), or C5a (10 nM). Random migration was conducted in the presence of medium M199. The results were expressed as the number of migrating cells per field (mean $\pm$ SE) (49).

Looking at the response of the neutrophil specific membrane receptors to the different chemoattractants, we repeatedly found reduction of the chemotaxis following intense exercise, regardless of the chemoattractant used, including Formylated peptides (fMLP), the

chemokine IL-8, and the activated complement component(C5a) (49). These chemokines attach to their specific receptors, fMLP-R (N- formyl-Met-Leu-Phe), IL-8-R (CXCR1 and CXCR2), and C5aR, which belong to the seven-transmembrane helix surface receptor family (“serpentine receptors”) that transduces signals downstream the cytoskeleton by coupling to heterotrimeric G-proteins (51). Once the signal has been triggered, rapid cytoskeletal rearrangement and chemotaxis take place. “Target” chemoattractants (fMLP, C5a) function primarily through a common signal-transduction pathway by stimulating p38 MAPK, whereas “host” intermediary chemoattractants (IL-8, LTB4) primarily function via the PI3K/Akt pathway (52). The surface density of the chemotactic receptor (C5aR), which serves as a representative model of receptor availability, was not affected 24 h after exercise. Moreover, the integrin CD11b/CD18, which represent one of the main receptors for neutrophil adhesiveness and crucial for normal chemotaxis, was also unaffected by exercise. Therefore, the chemotactic defect is not dependent on the specific receptor of activation, or on its specific pathway of transduction. We could speculate that the chemotactic impairment was related to a common defect at the membrane level, leading to decreased receptor availability or to other factors yet to be elucidated. For achieving appropriate chemotactic responses, an intact cytoskeleton structures are necessary (53,54). Continuous reorganization of the cytoskeleton is required for efficient F-actin polymerization and polarization. Both are important steps in the skeletal rearrangement during migration (55). Consequently, we studied the neutrophil F-actin neutrophil polarization and polymerization (49). Following fMLP stimulation, the cell undergoes sequential morphological changes from round to elongated geometrical forms (figure 2A). These changes reflect the cell activation and the ability to migrate against the chemotactic gradients toward the target (49,53). Using the green phalloidin test we found no correlation between the chemotactic defect and the ability to polymerize F-actin, indicating that the reduction in chemotaxis following exercise was not a result of the F-actin dysfunction. Despite the fact that positive correlation between chemotaxis and F-actin polymerization usually occurs, a lack of correlation in certain conditions has been reported (55).

To elucidate other cell skeletal responses to aerobic exercise, we studied the neutrophil polarization, known to be in tight correlation with the chemotactic activity.

Indeed, the neutrophil polarization was significantly decreased 24 h following aerobic exercise. This change also correlated with the decrease in chemotactic activity ( $r= 0.945$ ;  $P= 0.001$ ) (figure 2B) (49,50).

Since the neutrophil bactericidal activity and the oxidative burst were found to be normal, it seems that the signal transduction pathways are not affected following 30 min of intense aerobic exercise. Rather, it seems that aerobic exercise causes a skeletal impairment, and this eventually could leads to a reduction of the chemotactic activity. Most probably the impaired chemotaxis event, following a short bout of submaximal exercise, occurred at the effectors' machinery level, rather than at the level of the neutrophil membrane receptors. Others found no change in chemotaxis 24 h after a graded exercise to exhaustion (56). Giraldo et. al. reported increased chemotaxis immediately after moderate (45 min of 55% VO<sub>2</sub>max) and intense (1 hr of 70% VO<sub>2</sub>max) aerobic exercise that returned to basal values after 24 h (57). These discrepancies are probably related to differences in the type, intensity, and duration of exercise, timing of blood sampling, or use of different laboratory assays.




Fig. 2. Neutrophil polarization. A. The cell shape changes that occur following fMLP-stimulation. Three different cell shapes were recorded: non-activated - round cells (R), partially activated - intermediate cells (I), and fully activated - polarized cells (P). B. Analysis of the cells' morphological changes following fMLP-stimulation, in 11 athletes, before and after effort (mean $\pm$ SE) (49).

## 5. Therapeutic approach to exercise-induced immune suppression

Dietary and drug intervention have been reported to boost performance in athletes (58). They could block the transient immune changes, to prevent the oxidative stress and the inflammation induced by prolonged, intense exercise or excessive training. Some supplements as flavonoids were reported to benefit the immune system (59). In endurance events, iron and mineral supplements, together with antioxidant vitamins, help to prevent muscle damage (60). Carbohydrates enhance muscle glycogen stores. Glutamine and aminoacid supplementation did not prove to be beneficial (61).

Vitamin E (VE) and vitamin C, as antioxidants, play an important role in protecting the cells and muscles from damage (62-65). It is well-established that exercise exerts imbalance on the oxidative state by increasing Reactive Oxygen Species (ROS) and decreasing the level of antioxidants (63). As previously shown, intense or prolonged exercise can adversely affect the function of the immune system. It was found that submaximal aerobic activity (1h swim at 75-80% of VO<sub>2</sub>max) could produce oxidative damage within the neutrophils (64), which lose the appropriate antioxidant defense mechanisms, leading to a defective chemotactic ability (65). This impairment could rise from the increased levels of ROS and lipid peroxidation; both potentially could damage neutrophil function. The enhanced production of ROS, mainly by mitochondria, is associated with excessive oxidation of lipids, proteins, and nucleic acids, causing damage to cell membranes and to the physiological function of proteins and DNA (66-68). To defend themselves from ROS induced damage, cells contain

complex antioxidant mechanisms including enzymatic (e.g. superoxide dismutase, glutathione peroxidase, catalase) and non-enzymatic antioxidants (e.g., vitamin E (VE), vitamin C, beta-carotene). VE is the most important lipid-soluble antioxidant due to its abundance in cell and mitochondrial membranes and its ability to act directly on ROS and stop lipid peroxidation. This antioxidant is known to decrease the exercise-induced oxidative stress (69-71) and has been shown to protect against exercise-induced muscle damage (70). Neutrophils play a dual role in exercise-induced oxidative damage. On the one hand, they contribute to ROS formation during intense or prolonged exercise; on the other hand, intense exercise can produce oxidative damage within neutrophils. VE has an important role as anti-oxidant and an important role in maintaining normal neutrophil function. Chemotaxis, adherence, and phagocytic capacities of neutrophils were shown to be reduced in VE deficiency, improving after antioxidant treatment (65,72,73).

Our research focused on the phagocytic immune response to exercise, showing prevention of the impairments by vitamin E supplementation. The results of chemotaxis and polarization are shown in Figure 3, representing the mean  $\pm$  SEM of 7 trained men pre- and post-exercise, before and after 28 days of daily VE supplementation (74). We can see that daily supplementation of 800 IU d-alpha tocopheryl succinate, indeed corrected the defective neutrophil chemotaxis and polarization observed 24 hr post-exercise.

A relatively small number of studies have dealt with the effect of VE on neutrophil functions following exercise. To the best of our knowledge, there are no studies addressing the effect of VE on exercise-induced impaired chemotaxis. However, a beneficial effect of VE on chemotaxis was shown in other populations, such as healthy elderly men and women and elderly women with coronary heart disease or major depressive disorder (72,73). Improvement in chemotactic ability after VE supplementation was also found in rats with VE deficiency and in periparturient dairy cows (75,76).




Fig. 3. A. Correction of the defective neutrophil chemotactic activity (observed 24 h post-exercise) by vitamin E supplementation. B. Correction of the defective neutrophil polarization (observed 24 h post-exercise) by vitamin E supplementation (74).

Of note is that recent reports have emphasized that ROS production, through the reversible oxidation of thiol groups, has also important physiological influences on gene transcription and protein synthesis, as part of the adaptive processes that occurs after exercise. High dose antioxidant supplementation may interfere with these processes. Cooper's Group reported gene reorganization after intense exercise (77, 78).

In recent years, it has been reported that regular physical activity can have beneficial role in cancer's prevention and therapy (79,80). There is evidence of a protective effect of physical activity on colon and postmenopausal breast cancer (81). Further, it is also mounting that physical activity reduces risks of lung tumor metastases (82). It has been reported that exercise prevent the loss of muscle mass and functional capacity in chronic deteriorating conditions, beyond the beneficial psychological effects, which certainly improve the quality of life (83-85). Athletes are not immunocompromised by clinical definition, but could suffer from transitory, persistent immunosuppression, eventually leading to subclinical or clinical diseases. Recovery time is imperative in elite athletes involved in intense training and competitions. The temporary, sometimes multiple, mild impairments of the immune system could change into a chronic more severe immune dysfunction.

The approach should be multidisciplinary, including all care givers as sport medicine physicians, physiologists, immunologists, physiotherapists, nutritionists, psychologists and coaches. To achieve the main goals, an integrated model with programmed activities and clear guidelines for any specific type of sport is imperative. Recommendations should be directed to elite athletes and recreational sports, for sedentary individuals, for moderate and well trained subjects. The target is to maintain the balance of the immune system for the health of the athlete and his optimal performance.

## 6. References


- [1] Larrabee RC, Leukocytosis after violent exercise. *J Med Res*, 7:76-82, 1902
- [2] Cowles WN. Fatigue as a contributory cause of pneumonia. *Boston Med Surg J*, 179:555, 1918
- [3] Baetjer AM. The effect of muscular fatigue upon resistance. *Physiol Rev*, 12:453-468, 1932
- [4] Walsh NP, Gleeson M, Shephard RJ, Gleeson M, Woods JA, Bishop NC, Fleshner M, Green C, Pedersen BK, Hoffman-Goetz L, Rogers CJ, Northoff H, Abbasi A, Simon P. Part one: Immune function and exercise. *Exercise Immunol Rev*, 17:6-63, 2011
- [5] Petersen AM, Pedersen BK. The anti-inflammatory effect of exercise. *J Appl Physiol*, 98:1154-1162, 2005
- [6] Buettner P, Mosig S, Lechtermann A, Funke H, Mooren FC. Exercise affects the gene expression profiles of human white blood cells. *J Appl Physiol* 102: 26-36, 2007
- [7] Nieman DC, Nehlsen-Cannarella SL. The immune response to exercise. *Sem Hematol*, 31:166-179, 1994
- [8] Lamm ME. Current concepts in mucosal immunity. How epithelial transport of IgA antibodies relates to host defense. *Am J Physiol*, 274:G614-G617, 1998
- [9] Gleeson M, Pyne DB. Special feature for the Olympics: effects of exercise in the immune system: exercise effect on mucosal immunity. *Immunol Cell Biol*, 78:536-544, 2000
- [10] Mackinnon LT. Exercise and resistance to infectious diseases. In: Bahrke, Drews, Wentworth (Ed.), *Advances in Exercise Immunology*. Champaign, IL, pp. 1-26, 1999
- [11] Wolach B, Baehner RL, Boxer LA. Review: clinical and laboratory approach to the management of neutrophil dysfunction. *Isr J Med Sci*, 18:897-916, 1982

- [12] Boxer LA, Blackwood RA. Leukocyte disorders: quantitative and qualitative disorders of the neutrophil, Part 2. *Pediatr Rev*, 17:47-50, 1996
- [13] Dinauer MC. Disorders of neutrophil function: an overview. *Methods Mol Biol*, 412:489-504, 2007
- [14] Mackinnon LT. Chronic exercise training effects on immune function. *Med Sci Sports Exerc*, 32(7 Suppl):S369-376, 2000
- [15] Friman G, Wesslén L. Special feature for the Olympics: effects of exercise on the immune system: infections and exercise in high-performance athletes. *Immunol Cell Biol*, 78:510-522, 2000
- [16] Peters EM, Bateman ED. Ultramarathon running and URTI. *South African Med J*, 64:583-584, 1983
- [17] Nieman DC. Immune response to heavy exertion. *J Appl Physiol*, 82:1385-94, 1997
- [18] Lehmann M, Foster C, Keul J. Overtraining in endurance athletes: a brief review. *Med Sci Sport Exerc*, 25:854-862, 1993
- [19] Fitzgerald L. Overtraining increases the susceptibility to infection. *Int J Sport Med*, Suppl 1:S5-8, 1991
- [20] Pedersen BK, Ullum H. NK-cell response to physical activity. Possible mechanisms of action. *Med Sci Sports Exerc*, 26:140-146, 1994
- [21] Baron RD, Hutch MH, Kleeman K, Maccromac JN. Aseptic meningitis among members of high school football team. *JAMA*, 248:1724-1727, 1982
- [22] Morse LJ, Bryan JA, Murle JP. The Holly Cross College Football team Hepatitis outbreak. *JAJA*, 219:706-708, 1972
- [23] Nieman DC, Johanssen LM, Lee JW, Arabatzis K. Infectious episodes in runners before and after the Los Angeles Marathon. *J Sports Med Phys Fitness*, 30:316-328, 1990
- [24] Nieman DC. Prolonged aerobic exercise, immune response and risk of infection. In: *Exercise and Immune Function*, CRC Press, pp.143-161, 1996
- [25] Cox AG, Gleeson M, Pyne DB, Callister R, Hopkings WG, Fricker PA. Clinical and laboratory evaluation of upper respiratory symptoms in elite athletes. *Clin J Sports Med*, 18: 438-445, 2008
- [26] Suzuki K, Nakaji S, Yamada M, Totsuka M, Sato K, Sugawara K. Systemic inflammatory response to exhaustive exercise. *Cytokine kinetics. Exerc Immunol Rev*, 8:6-48, 2002
- [27] Cox AJ, Pyne DB, Saunders PU, Callister R, Gleeson M. Cytokine responses to treadmill running in healthy and illness-prone athletes. *Med Sci Sports Med*, 39:1918-1926, 2007
- [28] Bishop NC, Gleeson M. Acute and chronic effects of exercise on markers of mucosal immunity. *Front Biosci*, 14:4444-4456, 2009
- [29] Blain AK, Robson PJ, Walsh NP, Clark AM, Glennon L, Gleeson M. The effect of exercising to exhaustion at different intensities on saliva immunoglobulin A, protein and electrolyte secretion. *Int J Sports Med*, 19:547-552, 1998
- [30] Allgrove JE, Gomes E, Hough J, Gleeson M. Effects of exercise intensity on saliva antimicrobial proteins and markers of stress in active men. *J Sports Sci*, 26:653-661, 2008
- [31] Sharp JCM. Viruses and the Athlete. *Br J Sport Med*, 23:47-48, 1989
- [32] Eichner ER. Infectious mononucleosis-recognising the condition, 'reactivating' the patient. *Physician Sports Med*, 24:49-54, 1996
- [33] Friman G, Wesslén L, Karjalainen J, Rolf C. Infectious and lymphocytic myocarditis: epidemiology and factors relevant to sports medicine. *Scand J Med Sci Sports*, 5:269-278, 1995

- [34] Russell WR. Poliomyelitis, the paralytic stage and the effect of physical activity on the severity of paralysis. *Br Med J*, 1:465-471, 1949
- [35] Gautmanitan BG, Ghason JL, Lerner AM. Augmentation of the virulence of murine coxsackie virus B3 myocardiopathy by exercise. *J Exp Med*, 131:1121-1125, 1970
- [36] Ortega E. Neuroendocrine mediators in the modulation of phagocytosis by exercise: physiological implications. *Exerc Immunol Rev*, 9:70-93, 2003
- [37] Shephard RJ. Adhesion molecules, catecholamines and leucocyte redistribution during and following exercise. *Sports Med*, 33:261-284, 2003
- [38] Aoi W, Naito Y, Takanami Y, Kawai Y, Sakuma K, Ichikawa H, Yoshida N, Yoshikawa T. Oxidative stress and delayed onset muscle damage after exercise. *Free Radic Biol Med*, 37:480-487, 2004
- [39] Soppi E, Varjo J, Eskola J, Laitinen LA. Effect of strenuous physical stress on circulating lymphocyte number and function before and after training. *J Clin Lab Immunol*, 8:43-46, 1982
- [40] Gabriel H, Schwarz L, Born P, Kindermann W. Differential mobilization of leucocyte and lymphocyte subpopulations into the circulation during endurance exercise. *Eur J Appl Physiol*, 65:529-534, 1992
- [41] Nieman DC, Henson DA, Sampson CS, Herring JL, Stulles J, Conley M, Stone MH, Buttersworth DE, Davis JM. The acute immune response to exhaustive resistance exercise. *Int J Sports Med*, 16:322-328, 1995
- [42] Busse WW, Anderson CL, Hanson PG, Folts JD. The effect of exercise in the granulocyte response to isoproterenol in the trained athlete and unconditioned individual. *J Allergy Clin Immunol*, 65:358-364, 1980
- [43] Hooper S, Mackinnon LT, Howard A, Gordon RD, Bachmann AW. Markers for monitoring overtraining and recovery in elite swimmers. *Med Sci Sports Exerc*, 27:106-112 1995
- [44] Pizza FX, Mitchell JB, Davis BH, Starling RD, Holtz RW, Bigelow N. Exercise-induced muscle damage: Effect on circulating leukocyte and lymphocyte subsets. *Med Sci Sports Exerc*, 27:363-370, 1995
- [45] Shinkai S, Shore S, Shek PN, Shephard RJ. Acute exercise and immune function. Relationship between lymphocyte activity and changes in subset counts. *Int J Sports Med*, 13:452-461, 1992
- [46] Eliakim A, Wolach B, Kodesh E, Gavrieli R, Radnay J, Ben-Tovim T, Yarom Y, Falk B. Cellular and humoral immune response to exercise among gymnasts and untrained girls. *Int J Sports Med*, 18:208-212, 1997
- [47] Wolach B, Eliakim A, Gavrieli R, Kodesh E, Yarom Y, Schlesinger M, Falk B. Aspects of leukocyte function and the complement system following aerobic exercise in young female gymnasts. *Scand J Med Sci Sports*, 8:91-97, 1998
- [48] Wolach B, Falk B, Gavrieli R, Kodesh E, Eliakim A. Neutrophil function response to aerobic and anaerobic exercise in female judoka and untrained subjects. *Br J Sports Med*, 34:23-28, 2000
- [49] Wolach B, Gavrieli R, Ben-Dror SG, Zigel L, Eliakim A, Falk B. Transient decrease of neutrophil chemotaxis following aerobic exercise. *Med Sci Sports Exerc*, 37:949-954, 2005
- [50] Gavrieli R, Ashlagi-Amiri T, Eliakim A, Nemet D, Zigel L, Berger-Achituv S, Falk B, Wolach B. The effect of aerobic exercise on neutrophil functions. *Med Sci Sports Exerc*, 40:1623-1628, 2008

- [51] Prossnitz ER, Gilbert TL, Chiang S, Campbell JJ, Qin S, Newman W, Sklar LA, Ye RD. Multiple activation steps of the N-formyl-peptide receptor. *Biochemistry*, 38:2240–2247, 1999
- [52] Schraufstatter IU, Chung J, Burger M. IL-8 activates endothelial cell CXCR1 and CXCR2 through Rho and Rac signaling pathways. *Am J Physiol Lung Cell Mol Physiol*, 280:L1094–L1103, 2001
- [53] Chodniewicz D, Zhelev DV. Novel pathways of F-actin polymerization in the human neutrophil. *Blood*, 102:2251–2258, 2003
- [54] Weiner OD, Servant G, M. D. Welch MD, Mitchison TJ, Sedat JW, Bourne HR. Spatial control of actin polymerization during neutrophil chemotaxis. *Nature Cell Biol*, 1:75–81, 1999
- [55] Howard TH, Meyer WH. Chemotactic peptide modulation of actin assembly and locomotion in neutrophils. *J Cell Biol*, 98:1265–1271, 1984.
- [56] Hack V, Strobel G, Rau JP, Weicker H. The effect of maximal exercise on the activity of neutrophil granulocytes in highly trained athletes in a moderate training period. *European journal of applied physiology and occupational physiology*, 65:520-524, 1992
- [57] Giraldo E, Garcia JJ, Hinchado MD, Ortega E. Exercise intensity-dependent changes in the inflammatory response in sedentary women: role of neuroendocrine parameters in the neutrophil phagocytic process and the pro-/anti-inflammatory cytokine balance. *Neuroimmunomodulation*, 16:237-244, 2009
- [58] Walsh NP, Gleeson M, Pyne DB, Nieman DC, Dhabhar FS, Shephard RJ, Oliver SJ, Berman S, Kojouharova A. Position statement. Part two: Maintaining immune health. *Exercise Immunol Rev*, 17: 64-103, 2011
- [59] Nieman DC. Quercetin's bioactive effects in human athletes. *Curr Topic Nutraceut Res*, 8:33-44, 2010
- [60] Jackson MJ. Redox regulation of adaptive responses in skeletal muscle to contractile activity. *Free Radical Biology & Medicine*, 47:1267-1275, 2009
- [61] Gleeson M. Dosing and efficacy of glutamine supplementation in human exercise and sport training. *J Nutr*, 138: 2045S-2049S, 2008
- [62] Gomez-Cabrera MC, Domenech E, Romagnoli M, Arduini A, Borrás C, Pallardo FV, Sastre J, Viña J. Oral administration of vitamin C decreases muscle mitochondrial biogenesis and hampers training-induced adaptations in endurance performance. *American Journal of Clinical Nutrition*, 87:142-149, 2008
- [63] Finaud J, Scislawski V, Lac G, Durand D, Vidalin H, Robert A, Filaire E. Antioxidant status and oxidative stress in professional rugby players: evolution throughout a season. *Int J Sports Med*, 27:87-93, 2006
- [64] Ferrer, MD, Tauler, P, Sureda, A, Tur, JA and Pons, A. Antioxidant regulatory mechanisms in neutrophils and lymphocytes after intense exercise. *J Sports Sci* 27: 49-58, 2009
- [65] Baehner RL, Boxer LA. Role of membrane vitamin E and cytoplasmic glutathione in the regulation of phagocytic functions of neutrophils and monocytes. *American Journal of Pediatric Hematology/Oncology*, 1:71-16, 1979
- [66] Powers, SK, Lennon SL. Analysis of cellular responses to free radicals: Focus on exercise and skeletal muscle. *Proc Nutr Soc*, 58:1025-1033, 1999
- [67] Sachdev S, Davies KJ. Production, detection and adaptive responses to free radicals in exercise. *Free Radic Biol Med*. 44:215-223, 2008
- [68] Finaud J, Lac G, Filaire E. Oxidative stress: relationship with exercise and training *Sports Med*, 36:327-358, 2006

- [69] Silva LA, Pinho CA, Silveira PC, Tuon T, De Souza CT, Dal-Pizzol F, Pinho RA. Vitamin E supplementation decreases muscular and oxidative damage but not inflammatory response induced by eccentric contraction. *Journal of Physiological Sciences*, 60:51-57, 2010
- [70] Bloomer RJ, Goldfarb AH, McKenzie MJ. Oxidative stress response to aerobic exercise: comparison of antioxidant supplements. *Medicine & Science in Sports & Exercise*, 38:1098-1105, 2006
- [71] Williams SL, Strobel NA, Lexis LA, Coombes JS. Antioxidant requirements of endurance athletes: implications for health. *Nutrition Reviews*, 64:93-108, 2006
- [72] De la Fuente M, Ferrández MD, Burgos MS, Soler A, Prieto A, Miquel J. Immune function in aged women is improved by ingestion of vitamins C and E. *Canadian Journal of Physiology and Pharmacology*, 76:373-380, 1998
- [73] De la Fuente M, Hernanz A, Guayerbas N, Victor VM, Arnalich F. Vitamin E ingestion improves several immune functions in elderly men and women. *Free Radical Research*, 42:272-280, 2008
- [74] Gavrieli R, Berger-Achituv S, Ashlagi-Amiri T, Zigel L, Nemet D, Eliakim A, Falk B, and Wolach B. Vitamin E Prevents Neutrophil Impairment after a Single Bout of Intense Aerobic Exercise. Submitted for publication.
- [75] Harris RE, Boxer LA, Baehner RL. Consequences of vitamin-E deficiency on the phagocytic and oxidative functions of the rat polymorphonuclear leukocyte. *Blood*, 55:338-343, 1980
- [76] Politis I, Hidioglou N, White JH, Gilmore JA, Williams SN, Scherf H, Frigg M. Effects of vitamin E on mammary and blood leukocyte function, with emphasis on chemotaxis, in periparturient dairy cows. *American Journal of Veterinary Research*, 57:468-471, 1996
- [77] Cooper DM, Radom-Aizik S, Zaldivar F, Leu SY, Galassetti P. Effects of 30 min of aerobic exercise on gene expression and in human neutrophils. *J Appl Physiol*, 104:236-243, 2008
- [78] Connolly PH, Caiozzo VJ, Zaldivar F, Nemet D, Larson J, Hung S, Heck JD, Hatfield GW, Cooper DM. Effects of exercise on gene expression in human peripheral blood mononuclear cells. *J Appl Physiol*, 97:1461-1469, 2004
- [79] Ben-Eliyahu S, Page GG, Schleifer SJ. Stress, NK cells, and cancer: Still a promissory note. *Brain Behav Immun, Review* 21:881-887, 2007
- [80] Mc Tiernan A. Mechanisms linking physical activity with cancer. *Nat Rev Cancer*, 8:205-211, 2008
- [81] Friedenreich CM, Gregory J, Kopciuk KA, Mackey JR, Courneya KS. Prospective cohort study of lifetime physical activity and breast cancer survival. *Int J Cancer*, 124:1954-1962, 2009
- [82] Davis JM, Kohut ML, Jackson DA, Colbert LH, Mayer EP, Ghaffar A. Exercise effects on lung tumor metastases and in vitro alveolar macrophage antitumor cytotoxicity. *Am J Physiol*, 274:R1454-1459, 1998
- [83] Speck RM, Courneya KS, Masse LC, Duval S, Schmitz KH. An update of controlled physical activity trials in cancer survivors: a systematic review and meta-analysis. *J Cancer Surviv*, 4:87-100, 2010
- [84] Hoffman-Goetz L, Husted J. Exercise and Cancer. Do the biology and epidemiology correspond? *Exercise Immunology Rev*, 1: 81-96, 1995
- [85] World Cancer Research Fund. *Food, Nutrition, Physical Activity and the prevention of Cancer: a Global Perspective*. Washington DC, AICR, 2007



## **An International Perspective on Topics in Sports Medicine and Sports Injury**

Edited by Dr. Kenneth R. Zaslav

ISBN 978-953-51-0005-8

Hard cover, 534 pages

**Publisher** InTech

**Published online** 17, February, 2012

**Published in print edition** February, 2012

For the past two decades, Sports Medicine has been a burgeoning science in the USA and Western Europe. Great strides have been made in understanding the basic physiology of exercise, energy consumption and the mechanisms of sports injury. Additionally, through advances in minimally invasive surgical treatment and physical rehabilitation, athletes have been returning to sports quicker and at higher levels after injury. This book contains new information from basic scientists on the physiology of exercise and sports performance, updates on medical diseases treated in athletes and excellent summaries of treatment options for common sports-related injuries to the skeletal system.

### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Baruch Wolach (2012). Exercise and the Immune System – Focusing on the Effect of Exercise on Neutrophil Functions, An International Perspective on Topics in Sports Medicine and Sports Injury, Dr. Kenneth R. Zaslav (Ed.), ISBN: 978-953-51-0005-8, InTech, Available from: <http://www.intechopen.com/books/an-international-perspective-on-topics-in-sports-medicine-and-sports-injury/exercise-and-the-immune-system-focusing-on-the-effect-of-exercise-on-neutrophil-functions>

**INTECH**  
open science | open minds

### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166  
[www.intechopen.com](http://www.intechopen.com)

### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen