

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,000

Open access books available

125,000

International authors and editors

140M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# Hepatic Lipid Accumulation by High Cholesterol Diet is Inhibited by the Low Dose Fish Oil in Male C57BL/6J Mice

Satoshi Hirako, Miki Harada, Hyoun-Ju Kim\*, Hiroshige Chiba and Akiyo Matsumoto  
*Josai University  
Japan*

## 1. Introduction

Atherosclerotic diseases such as ischemic heart disease and cerebral infarction account for the major causes of death in many countries. It is well known that hyperlipidemia is closely associated with these diseases (Kodama, 1990). Fish oil consumption reduces lipogenesis and induces fatty acid oxidation in liver (Rustan, 1990; Halminski, 1991), and ameliorates plasma and hepatic lipid levels (Nestel, 1990). These effects of fish are greatly attributed to the action of n-3 polyunsaturated fatty acids, such as eicosapentaenoic acid (EPA) or docosahexaenoic acid (DHA). Several studies have reported that n-3 PUFA decreases the expression of genes coding for lipogenesis enzymes such as fatty acid synthase (FAS), acetyl-CoA carboxylase (ACC), and stearoyl-CoA desaturase (SCD) through the lowering of sterol regulatory element binding protein-1 (SREBP-1) (Kim, 1999; Shimano, 1999; Nakatani, 2002). In addition, n-3 PUFA activates the peroxisome proliferator-activated receptor  $\alpha$  (PPAR $\alpha$ ) (Krey, 1997). PPAR $\alpha$  is a ligand-activated transcription factor that regulates the expression of genes involved in triglyceride hydrolysis and fatty acid oxidation (Latruffe, 1997; Nakatani, 2002; Schoonjans, 1996), such as acyl-CoA oxidase (AOX), lipoprotein lipase (LPL), medium-chain acyl-CoA dehydrogenase (MCAD), acyl-CoA synthetase (ACS), and uncoupling protein-2 (UCP-2).

Cholesterol is crucial for the components of animal cell membranes, and for the synthesis of bile acid and steroid hormone. Cholesterol in the inner cavity of the small intestine are entered in to epithelial cells by Niemann-Pick C1 like-1 (NPC1L1) protein on the brush border of epithelial cells of the small intestine (Davis, 2004), and cholesterol is released into the lymph duct. It is well known that cholesterols, including 22- and 24-hydroxy cholesterol, act as ligands for liver X receptor (LXR)  $\alpha$  (Schultz, 2000). LXR $\alpha$ , a nuclear receptor, forms a heterodimer together with the retinoid X receptor (RXR) and combines with LXR element (LXRE) at the promoter of target genes to control transcription. LXR target genes include cholesterol 7 $\alpha$ -hydroxylase (CYP7A1), ATP-binding cassette transporter (ABC)A1, ABCG1, and SREBPs, which are highly involved in cholesterol metabolism (Repa, 2000; Peet, 1998).


We previously indicated that 20% fish oil diet dramatically inhibited hepatic lipids accumulation in female C57BL/6J mice fed 2% cholesterol (Hirako, 2010). In this study, we

determined whether 2 or 5en% fish oil, equivalent to 10 or 25% of the total fat energy, improves lipid metabolism in high cholesterol diet fed male C57BL/6J mice. Mice were given SO diet consisted of 20en% safflower oil, 2FO diet consisted of 2en% fish oil plus 18en% safflower oil, 5FO diet consisted of 5en% fish oil plus 15en% safflower oil, and SO/CH, 2FO/CH, 5FO/CH are consisted of SO, 2FO, 5FO with 2%cholesterol. The body fat composition of mice was examined radiographically using an X-ray CT for experimental animals. Blood parameter and hepatic lipids were measured using enzymatic methods. Hepatic mRNA expression levels were measured by real-time RT-PCR.

## 2. Results and discussion

### 2.1 Body weight and tissue weight did not change with the low dose fish oil feeding in male C57BL/6J mice

Final body weights were not significantly different among the groups. No large difference in weights of the white adipose tissue around the uterus and brown adipose tissue from the interscapular region between the groups was observed. In previous study, we reported that body weight gain was significantly decreased in female C57BL/6J mice fed 20 en% fish oil (Hirako, 2010). However, our study observed that 5 en% fish oil feeding in female C57BL/6J mice did not modify body weights (data not shown). These results revealed that 2 and 5 en% fish oil feeding in male C57BL/6J mice did not modify body weight and adipose tissue weight




Representative X-ray CT images of mice fed SO, SO/CH, 2FO, 2FO/CH, 5FO, and 5FO/CH for 8 weeks, at the L3 level (upper). The areas indicated with pink, yellow, and light-blue are visceral fat, subcutaneous fat, and muscle, respectively. CT-estimated amounts of visceral fat and subcutaneous fat in the abdominal area of L2–L4 (bottom). Values represent means  $\pm$  S.D. (n=5). Means with different letters are different at the  $p < 0.05$  level by Fisher's protected least significant difference (PLSD) test.

Fig. 1. CT-based fat tissues composition analysis of 16-week-old male C57BL/6J mice.

as well as female C57BL/6J mice. CT scan analysis showed that visceral and subcutaneous fat mass were significantly increased in the 5FO and 5FO/CH groups, but not changed by cholesterol feeding (Figure 1). Meanwhile, these levels were high in all group compared with the fat mass in the female C57BL/6J mice ( $0.45 \pm 0.28 \text{ cm}^3$  visceral fat in male C57BL/6J mice fed the SO diet and  $0.24 \pm 0.13 \text{ cm}^3$  visceral fat in female C57BL/6J mice fed the SO diet).

## 2.2 Hepatic tissue histology and lipid levels modified with the low dose fish oil feeding in male C57BL/6J mice

Although the liver weight significantly increased in the SO/CH group compared with the SO group, there was no difference in the fish oil groups compared with the SO group (SO:  $0.96 \pm 0.10$ , SO/CH:  $1.31 \pm 1.2$ , 2FO:  $1.00 \pm 0.13$ , 2FO/CH:  $1.11 \pm 0.12$ , 5FO:  $0.92 \pm 0.08$ , 5FO/CH:  $1.12 \pm 0.12$ ,  $p < 0.05$ ). The liver size in the SO/CH group had enlarged, and the entire surface had a pale color, suggestive of increased lipid storage. In contrast, the livers of the 2FO/CH and 5FO/CH groups were less pale and had a normal reddish appearance (data not shown). The hepatic tissues showed numerous lipid droplets in the livers of SO/CH group. However, these lipid droplets were markedly decreased in the 2FO/CH and 5FO/CH groups (Figure 2A). These results in the microscopic images of hepatic tissue were confirmed in hepatic lipid content reductions. Hepatic TG levels in both 2FO/CH and 5FO/CH group significantly decreased to about 50% of that in the SO/CH group. Hepatic TC levels also decreased by about 40% in both 2FO/CH and 5FO/CH group compared with the SO/CH group (Figure 2B). Fabbrini suggested that hepatic lipid accumulation is considered a risk factor for fatty liver and steatohepatitis, which promote the development



H&E-stained liver sections (A), total cholesterol (TC) and triglycerides (TG) (B) in mice fed SO, SO/CH, 2FO, 2FO/CH, 5FO, and 5FO/CH for 8. Values represent means  $\pm$  S.D. ( $n=5$ ). Means with different letters are different at the  $p < 0.05$  level by Fisher's protected least significant difference (PLSD) test.

Fig. 2. Liver tissue histology and lipid levels of 16-week-old male C57BL/6J mice.

of insulin resistance, dyslipidemia, and cardiovascular disease (Fabbrini, 2010). In this study, lipid deposition caused by high cholesterol feeding significantly decreased in the 2FO/CH and 5FO/CH groups, in which low dose fish oil was used as the lipid source. We confirmed that the 2 en% fish oil exerts ameliorating effects on hepatic lipid accumulation due to dietary cholesterol consumption in male C57BL/6J mice.

### 2.3 The hepatic mRNA levels of lipid metabolism-regulating genes modified with the fish oil or cholesterol feeding in male C57BL/6J mice

The hepatic mRNA levels of lipid metabolism-regulating genes are shown in Table 1. The mRNA levels of SREBP-1c, which is transcription factor of genes related to lipogenesis, were not significantly changed in all groups. Low dose fish oil or cholesterol feeding did not particularly affect in FAS mRNA. However, SCD1 mRNA levels significantly decreased in the fish oil groups regardless of the addition the addition of cholesterol, compared with the SO/CH group. Previous studies showed that fish oil feeding decreases SREBP1c mRNA expression and/or mature protein production and results in the inhibition of SREBP1 target genes, such as ACC, FAS, and SCD-1 (Kim, 1999; Nakatani, 2002). In this study, different results were observed, suggesting that the inhibitory effect of fish oil on fatty acid biosynthesis is more clear at high dose fish oil.

	SO	SO/CH	2FO	2FO/CH	5FO	5FO/CH
<b>SREBPs</b>						
SREBP-1c	1.00 ± 0.31	1.27 ± 0.45	0.96 ± 0.56	1.21 ± 0.99	0.88 ± 0.29	0.81 ± 0.55
SREBP-2	1.00 ± 0.51 <sup>c</sup>	0.95 ± 0.06 <sup>c</sup>	1.88 ± 0.19 <sup>a</sup>	0.93 ± 0.14 <sup>c</sup>	1.41 ± 0.32 <sup>b</sup>	0.71 ± 0.12 <sup>c</sup>
Insig-1	1.00 ± 0.76 <sup>bc</sup>	1.44 ± 0.75 <sup>abc</sup>	2.42 ± 1.50 <sup>a</sup>	0.89 ± 0.52 <sup>bc</sup>	1.87 ± 0.57 <sup>ab</sup>	0.61 ± 0.20 <sup>c</sup>
<b>Fatty acid biosynthesis</b>						
FAS	1.00 ± 0.84 <sup>a</sup>	0.81 ± 0.58 <sup>ab</sup>	0.98 ± 0.39 <sup>a</sup>	0.58 ± 0.24 <sup>ab</sup>	0.85 ± 0.23 <sup>ab</sup>	0.32 ± 0.19 <sup>b</sup>
SCD1	1.00 ± 0.33 <sup>ab</sup>	1.46 ± 0.73 <sup>a</sup>	0.43 ± 0.32 <sup>c</sup>	0.82 ± 0.38 <sup>bc</sup>	0.35 ± 0.08 <sup>c</sup>	0.63 ± 0.19 <sup>bc</sup>
<b>Cholesterol homeostasis</b>						
HMGCo(A) Reductase	1.00 ± 0.30 <sup>a</sup>	0.67 ± 0.15 <sup>bc</sup>	0.85 ± 0.37 <sup>ab</sup>	0.66 ± 0.03 <sup>bc</sup>	1.07 ± 0.23 <sup>a</sup>	0.54 ± 0.10 <sup>c</sup>
ABCG5	1.00 ± 0.57 <sup>c</sup>	2.61 ± 0.29 <sup>a</sup>	0.94 ± 0.15 <sup>c</sup>	1.68 ± 0.25 <sup>b</sup>	0.81 ± 0.13 <sup>c</sup>	1.41 ± 0.18 <sup>b</sup>
ABCG8	1.00 ± 0.32 <sup>d</sup>	1.54 ± 0.32 <sup>a</sup>	1.25 ± 0.29 <sup>cd</sup>	2.18 ± 0.72 <sup>ab</sup>	0.95 ± 0.15 <sup>d</sup>	1.95 ± 0.43 <sup>bc</sup>
<b>Fatty acid β-oxidation</b>						
PPARα	1.00 ± 0.30 <sup>b</sup>	1.74 ± 0.74 <sup>ab</sup>	2.51 ± 1.03 <sup>a</sup>	2.32 ± 0.68 <sup>a</sup>	1.93 ± 0.49 <sup>a</sup>	1.97 ± 0.77 <sup>a</sup>
AOX	1.00 ± 0.49 <sup>c</sup>	1.54 ± 0.61 <sup>bc</sup>	1.66 ± 0.39 <sup>b</sup>	1.79 ± 0.43 <sup>abc</sup>	1.82 ± 0.33 <sup>ab</sup>	2.42 ± 0.66 <sup>a</sup>
UCP2	1.00 ± 0.32 <sup>ab</sup>	1.54 ± 0.32 <sup>a</sup>	0.99 ± 0.74 <sup>ab</sup>	1.19 ± 0.68 <sup>ab</sup>	0.70 ± 0.13 <sup>b</sup>	1.05 ± 0.38 <sup>ab</sup>
<b>Bile acid bioynthesis</b>						
CYP7A1	1.00 ± 0.57 <sup>c</sup>	3.12 ± 0.79 <sup>ab</sup>	1.27 ± 1.17 <sup>c</sup>	1.96 ± 0.80 <sup>bc</sup>	1.43 ± 0.81 <sup>c</sup>	3.63 ± 1.01 <sup>a</sup>
CYP8B1	1.00 ± 0.45 <sup>b</sup>	1.26 ± 0.29 <sup>ab</sup>	1.12 ± 0.47 <sup>ab</sup>	1.33 ± 0.32 <sup>ab</sup>	1.01 ± 0.19 <sup>b</sup>	1.55 ± 0.58 <sup>a</sup>

The mRNA expression levels in liver of male C57BL/6J mice fed SO, SO/CH, 2FO, 2FO/CH, 5FO, and 5FO/CH for 8 weeks. Values represent means ± S.D. (n=4-5). Means with different letters are different at the p<0.05 level by Fisher's protected least significant difference (PLSD) test.

Table 1. Expression of genes associated with lipid metabolism in the liver

The mRNA levels of PPARα and AOX, genes involved in fatty acid oxidation, were significantly higher in the fish oil groups regardless of the addition of cholesterol. However, the mRNA levels of UCP-2, which is involved in heat production, were unaffected by low dose fish oil or cholesterol feeding. The mRNA levels of ABCG5 and ABCG8, genes involved in cholesterol transport into the bile, were significantly induced with the cholesterol addition. Biliary cholesterol excretion increases due to the upregulation of these genes (Repa, 2002). These increases in ABCG5 and ABCG8 mRNA by cholesterol feeding are

crucial for the maintenance of cholesterol homeostasis. Indeed, the mRNA levels of CYP7A1, the rate-limiting gene in bile acid synthesis, were significantly increased in all cholesterol-supplemented groups. However, no such increases were observed in the expression of CYP8B1 mRNA. The synthesis of bile acid in the liver is important in the catabolic pathway of cholesterol. The bile acid synthesis is controlled by CYP7A1 (Berge, 2000). In this study, the mRNA levels of CYP7A1 were not significantly affected by fish oil feeding, but its levels significantly increased when cholesterol was added. This is consistent with previous observation that CYP7A1 mRNA level is increased on the addition of cholesterol to maintain the homeostasis of cholesterol.

### 3. Conclusion

Body weight gains and adipose tissue weights did not change with the low dose fish oil feeding in male C57BL/6J mice. Hepatic triglyceride and total cholesterol levels of the SO/CH group were dramatically increased compared to the SO group. However, in 2FO/CH and 5FO/CH groups, the hepatic lipids were significantly decreased compared to the SO/CH group. Low dose fish oil or cholesterol feeding did not particularly affect in SREBP-1C and FAS mRNA levels. But, PPAR $\alpha$  and AOX mRNA levels were significantly higher in the fish oil groups regardless of the cholesterol addition. The present study indicates that low dose fish oil obviously improved the hepatic lipid accumulation by high cholesterol diet. And, this improving effect is partly due to the fatty acid degradation, which was facilitated by increased expression of fatty acid oxidation-related genes, such as AOX.


### 4. Acknowledgments

We would like to thank NOF Corporation (Tokyo, Japan) for providing FO.

### 5. References

- Berge KE, Tian H, Graf GA, Yu L, Grishin NV, Schultz J, Kwiterovich P, Shan B, Barnes R, Hobbs HH. Accumulation of dietary cholesterol in sitosterolemia caused by mutations in adjacent ABC transporters. *Science* 2000; 290: 1771-1775
- Davis HR Jr, Zhu LJ, Hoos LM, Tetzloff G, Maguire M, Liu J, Yao X, Iyer SP, Lam MH, Lund EG, Detmers PA, Graziano MP, Altmann SW. Niemann-Pick C1 Like 1 (NPC1L1) is the intestinal phytosterol and cholesterol transporter and a key modulator of whole-body cholesterol homeostasis. *J Biol Chem* 2004; 279: 33586-33592
- Fabbrini E, Sullivan S, Klein S. Obesity and nonalcoholic fatty liver disease: biochemical, metabolic, and clinical implications. *Hepatology* 2010; 51: 679-689
- Halminski MA, Marsh JB, and Harrison EH: Differential effects of fish oil, safflower oil and palm oil on fatty acid oxidation and glycerolipid synthesis in rat liver. *J Nutr*, 1991; 121: 1554-1561
- Hirako S, Kim HJ, Arai T, Chiba H, Matsumoto A. Effect of concomitantly used fish oil and cholesterol on lipid metabolism. *J Nutr Biochem* 2010; 21: 573-579
- Kim HJ, Takahashi M, Ezaki O: Fish oil feeding decreases mature sterol regulatory element-binding protein 1 (SREBP-1) by down-regulation of SREBP-1c mRNA in mouse

- liver. A possible mechanism for down-regulation of lipogenic enzyme mRNAs. *J Biol Chem*, 1999; 274: 25892-25898
- Kodama K, Sasaki H, Shimizu Y. Trend of coronary heart disease and its relationship to risk factors in a Japanese population: a 26-year follow-up, Hiroshima/Nagasaki study. *Circ J*, 1990; 54: 414-421
- Krey G, Braissant O, L'Horsset F, Kalkhoven E, Perroud M, Parker MG, Wahli W: Fatty acids, eicosanoids, and hypolipidemic agents identified as ligands of peroxisome proliferator-activated receptors by coactivator-dependent receptor ligand assay. *Mol Endocrinol*, 1997; 11: 779-791
- Latruffe N, Vamecq J: Peroxisome proliferators and peroxisome proliferator activated receptors (PPARs) as regulators of lipid metabolism. *Biochimie*, 1997; 79: 81-94
- Nakatani T, Tsuboyama-Kasaoka N, Takahashi M, Miura S, Ezaki O; Mechanism for peroxisome proliferator-activated receptor- $\alpha$  activator-induced up-regulation of UCP2 mRNA in rodent hepatocytes. *J Biol Chem*, 2002; 277: 9562-9569
- Nestel PJ. Effects of N-3 fatty acids on lipid metabolism. *Annu Rev Nutr*, 1990; 10:149-167
- Peet DJ, Turley SD, Ma W, Janowski BA, Lobaccaro JM, Hammer RE, Mangelsdorf DJ. Cholesterol and Bile Acid Metabolism Are Impaired in Mice Lacking the Nuclear Oxysterol Receptor LXR $\alpha$ . *Cell* 1998; 93: 693-704
- Repa JJ, Berge KE, Pomajzl C, Richardson JA, Hobbs H, Mangelsdorf DJ. Regulation of ATP-binding cassette sterol transporters ABCG5 and ABCG8 by the liver X receptors alpha and beta. *J Biol Chem* 2002 ; 277: 18793-18800
- Repa JJ, Liang G, Ou J, Bashmakov Y, Lobaccaro JM, Shimomura I, Shan B, Brown MS, Goldstein JL, Mangelsdorf DJ. Regulation of mouse sterol regulatory element-binding protein-1c gene (SREBP-1c) by oxysterol receptor, LXR $\alpha$  and LXR $\beta$ . *Genes Dev* 2000; 14: 2819-2830
- Rustan AC, Nossen JO, Christiansen EN, and Drevon CA: Eicosapentaenoic acid reduces hepatic synthesis and secretion of triacylglycerol by decreasing the activity of acyl-coenzyme A:1,2- diacylglycerol acyltransferase. *J Lipid Res*, 1988; 29: 1417-1426
- Schoonjans K, Staels B, Auwerx J: The peroxisome proliferator activated receptors (PPARs) and their effects on lipid metabolism and adipocyte differentiation. *Biochim Biophys Acta*, 1996; 1302: 93-109
- Schultz JR, Tu H, Luk A, Repa JJ, Medina JC, Li L, Schwendner S, Wang S, Thoolen M, Mangelsdorf DJ, Lustig KD, Shan B. Role of LXRs in control of lipogenesis. *Genes Dev* 2000; 14: 2831-2838
- Shimano H, Yahagi N, Amemiya-Kudo M, Hasty AH, Osuga J, Tamura Y, Shionoiri F, Iizuka Y, Ohashi K, Harada K, Gotoda T, Ishibashi S, Yamada N. Sterol regulatory element-binding protein-1 as a key transcription factor for nutritional induction of lipogenic enzyme genes. *J Biol Chem* 1999; 274: 35832-35829



### **Novel Strategies in Ischemic Heart Disease**

Edited by Dr. Umashankar Lakshmanadoss

ISBN 978-953-51-0184-0

Hard cover, 450 pages

**Publisher** InTech

**Published online** 29, February, 2012

**Published in print edition** February, 2012

The first edition of this book will provide a comprehensive overview of ischemic heart disease, including epidemiology, risk factors, pathogenesis, clinical presentation, diagnostic tests, differential diagnosis, treatment, complications and prognosis. Also discussed are current treatment options, protocols and diagnostic procedures, as well as the latest advances in the field. The book will serve as a cutting-edge point of reference for the basic or clinical researcher, and any clinician involved in the diagnosis and management of ischemic heart disease. This book is essentially designed to fill the vital gap existing between these practices, to provide a textbook that is substantial and readable, compact and reasonably comprehensive, and to provide an excellent blend of "basics to bedside and beyond" in the field of ischemic heart disease. The book also covers the future novel treatment strategies, focusing on the basic scientific and clinical aspects of the diagnosis and management of ischemic heart disease.

#### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Satoshi Hirako, Miki Harada, Hyoun-Ju Kim, Hiroshige Chiba and Akiyo Matsumoto (2012). Hepatic Lipid Accumulation by High Cholesterol Diet is Inhibited by the Low Dose Fish Oil in Male C57BL/6J Mice, *Novel Strategies in Ischemic Heart Disease*, Dr. Umashankar Lakshmanadoss (Ed.), ISBN: 978-953-51-0184-0, InTech, Available from: <http://www.intechopen.com/books/novel-strategies-in-ischemic-heart-disease/hepatic-lipid-accumulation-by-high-cholesterol-diet-is-inhibited-by-the-low-dose-fish-oil-in-male-c5>

**INTECH**  
open science | open minds

#### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166  
[www.intechopen.com](http://www.intechopen.com)

#### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821



© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen