

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,300

Open access books available

132,000

International authors and editors

160M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



The Genetic and Epigenetic Bases of Gastritis

Alejandro H. Corvalan, Gonzalo Carrasco and
Kathleen Saavedra

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/53131>

1. Introduction

Gastritis, the inflammation of the mucosal layer of the stomach, is a major clinical entity due to its association with gastric cancer and peptic ulcer disease. The primary cause of gastritis is the infection with the microaerophilic gram negative *Helicobacter pylori* that during the early phases elicits an acute inflammatory response which eventually evolves to a long-standing chronic gastritis (Ruggiero 2012). In the case of gastric cancer development, chronic gastritis is the first step of the so-called multistep cascade of gastric cancer. This sequence includes the non-atrophic chronic gastritis, multifocal atrophic gastritis, intestinal metaplasia, dysplasia and invasive carcinoma as were described in detail by Correa as the “human model of gastric carcinogenesis” (Fig. 1) (Correa et al. 2007). This multistep model hypothesizes that the sequence of these lesions reflects a dynamic process from a naive inflammation caused by *H. pylori* infection to a fully malignant neoplasm of the stomach (Correa et al. 1976; Cuello et al. 1976; Haenszel et al. 1976; Correa et al. 1990). Independent epidemiological studies have confirmed that non-atrophic, atrophic, intestinal metaplasia and dysplasia are all linked through a sequential cause-effect relationship, thus supporting the concept of a human model for gastric carcinogenesis (Ohata et al. 2004). However, the risk of malignant transformation of these lesions is poorly defined. Long-term follow-up studies have shown a risk from 10% to 17% in the case of dysplasia (Saraga et al. 1987; Coma del Corral et al. 1990; Koch et al. 1990; Whiting et al. 2002; Rugge et al. 2003). For intestinal metaplasia, the risk assessment has conflicting results and therefore a limited clinical value (Ramesar et al. 1987; Silva et al. 1990; Rokkas et al. 1991; Conchillo et al. 2001; Vannella et al. 2012). The recently developed Operative Link for Gastritis Assessment (OLGA) staging system (Rugge et al. 2007), through the evaluation of the extension and site of the atrophic changes, is an attempt to evaluate the risk of chronic gastritis to progress to intestinal metaplasia and gastric cancer (Rugge et al. 2008; Capelle et al. 2010; Rugge et al. 2010). In this scenario, the identifi-

cation of molecular bases in the multistep process of gastric carcinogenesis is highly relevant since it will contribute greatly to the risk assessments of the precursor lesions of gastric cancer. In this chapter we will primarily attempt to summarize and integrate our current knowledge of the genetic as well as epigenetic bases of the dynamic process of chronic gastritis, as well as the other entities of the so-called multistep cascade of gastric cancer.




Figure 1. Multistep cascade of gastric cancer. This sequence begins with the infection of *H. pylori* to sequential steps of the precancerous cascades. Taken from Correa & Piazuelo *J Dig Dis.* 2012;13:2–9.

2. The genetic bases of gastritis

Human allelic variations at single nucleotide polymorphisms (SNP) are involved at different stages of gastric carcinogenesis. Accordingly, the dynamics of chronic gastritis might be associated with specific allelic variants. These variants, recognized as polymorphisms when occur with a frequency of >1% in the normal population, affects mostly genes of the inflammatory response genes, detoxification enzymes, and cancer-related processes (Gonzalez et al. 2002).

Polymorphisms in inflammatory response genes

Two clear examples of polymorphisms associated with inflammatory response are interleukin-1 gene cluster and Toll-like receptors (TLRs). As shown in Table 1, Interleukin-1-beta (IL-1B) -31 T genotype and interleukin-1-receptor antagonist (IL1RN) IVS 86 bp VNTR, 2/2

genotype enhance the production of IL-1B, which is associated with an increased risk of hypochlorhydria induced by *H. pylori*. Because IL-1B is an important pro-inflammatory cytokine and a powerful inhibitor of gastric acid secretion, it might explain the higher risk of development of gastric cancer in chronic gastritis - *H. pylori* infected patients (El-Omar et al. 2000; Hwang et al. 2002; Yuzhalin 2011). The presence of polymorphisms in the non-coding but regulatory regions may also alter the levels of gene transcription and therefore the susceptibility to gastric cancer. In this sense, single A/T SNP at position -251 from the transcription start site in the promoter region of the IL-8 gene is one of the best examples. The presence of the -251A allele tended to be associated with an increase in IL-8 production and has been reported that this allele is associated with an increased risk for gastric cancer at cardia location (Hull et al. 2000; Xue et al. 2012). It could be extrapolated that IL-8 -251A allele may increase the risk of developing cancer through the elevation of its IL-8 expression. Mediators of the innate immune response, which provide first line of host defense against harmful pathogens are also involved in the dynamics of chronic gastritis. A good example for this type of inflammatory response are the Toll-like receptors (TLRs) (El-Omar et al. 2008). It has been reported that both, the 22-bp nucleotide deletion (-196 to -174 del) in the promoter region of the TLR-2 gene and the +896A/G and +1196C/T polymorphisms (Asp299Gly and Thr399Ile) in the coding region of the TLR-4 gene, promotes a rapidly progressive chronic gastritis (de Oliveira et al. 2012). Since it is well known that not all *H. pylori* are equally associated with the risk of development of gastric cancer, the combination of bacterial and inflammatory host genotypes have been explored. Figueiredo et al (2002) have shown that the infection with aggressive *cagA*-positive strains of *H. pylori* were associated with an increased risk of gastric cancer. In addition, the homozygous IL-1B-511*T carriers (IL-1B-511*T/*T or IL-1B-511*T/*C) and the short allele of IL-1RN (IL-1RN*2/*2) also had an increased susceptibility. However, for each combination of bacterial/host genotype, the odds of having gastric cancer were greatest in those with both bacterial and host high-risk genotypes.

Polymorphisms in genes associated with detoxification enzymes

Members of the cytochrome P-450 superfamily belong to the detoxification enzymes. One of the members of this family is the CYP2E1, a naturally ethanol-inducible enzyme involved in the metabolic activation of low molecular weight compounds such as N-nitrosamines (Boccia et al. 2007). Functional CYP2E1 polymorphisms in the 5'-flanking region (PstI, RsaI) alter the transcriptional activity of the gene. A meta-analysis by Boccia et al (2007) suggests that the CYP2E1 PstI/RsaI polymorphism may be a risk factor for gastric cancer in asiatic populations and a synergic relation with tobacco-related detoxification Glutathione S-transferase (GST) genes (see below) may account for a proportion of these cases. GSTs and the polymorphic arylamine N-acetyltransferases (NAT1 and NAT2) are another type of detoxification enzymes that metabolize tobacco-related carcinogens. Zendehdel et al (2009) reported a weak linkage between the GSTP1 Ile105Val polymorphism and the risk for gastric cancer at cardia location (OR = 1.4; 95% CI 0.9-2.1). On the other hand, Katoh et al (2000) found that the NAT1*10 allele should be considered a risk factor among heavy smokers with well-differentiated tumors (Table 1).

Polymorphisms of cancer-related genes

In cancer-related processes, polymorphisms have been reported in the promoter region of the MET gene, a crucial gene to multiple oncogenic pathways and metastatic behavior (Gherardi et al. 2012). Sequencing of the promoter region of MET revealed some alterations scattered in a proportion of clinical samples of gastric cancer. The most common substitutions were -304C>A and 206C>G. The presence of these SNPs altered the junction sites for putative transcription factors such as Sp1 and AP-1/AP-2. As a consequence, the transcription of MET was constantly being activated (Trzyna et al. 2012). Another example is the polymorphism -160 A at the promoter region of the E-cadherin gene (CDH1), which has been associated with the DNA hypermethylation (see below) of the promoter region (Borges Bdo et al. 2010). In a Japanese study, 117 cases of gastric cancer with *H. pylori*-induced chronic gastritis were compared with 116 cancer-free yet *H. pylori*-induced chronic gastritis controls. It was found that the Pro/Pro allele of CDH-1 was associated with an increased risk of developing diffuse-type of gastric cancer compared to the Arg/Arg (Hiyama et al. 2002; Zhou et al. 2007). Finally, other polymorphisms of cancer-related genes were reported to show significant associations with gastric cancer risk including EGFR, VEGF and p53 (Tahara et al. 2009). Taken together, data presented here suggest that allelic variations at regulatory or coding regions will affect gene expression patterns and modify the balance towards a more rapid progression of chronic gastritis and other preneoplastic lesions of the multistep cascade of gastric cancer (Fig. 2).




Figure 2. Allelic variations at regulatory regions will affect gene expression patterns. (A) Single nucleotide polymorphisms (SNP) in promoter region can alter the binding of transcription factors required for the expression of a gene. (B) These variations may increase or decrease the expression of the affected gene, which eventually can increase the risk of developing cancer.

Gene	Variation Allelic	SNP	rs number	Effect	Reference
IL1B	SNP	-31 T/C	rs1143627	Increased IL-1beta production and inhibit gastric acid secretion.	(Yuzhalin et al., 2011)
IL-8	SNP	-251 A/T	rs4073	Increased IL-8 production.	(Xue et al., 2012)
IL-10	SNP	-1082 A/G	rs1800896	Reported associated with the pathologically intestinal-type gastric cancer or anatomically cardia-type gastric cancer.	(Ni et al., 2012)
TLR2	Deletion	-196 to -174del	not applicable	Susceptibility to gastric cancer in the Southeastern Brazilian population.	(de Oliveira et al., 2012)
TLR4	SNP	+ 896 A/G	rs4986790		
CYP2E1	RFLP	CYP2E1 c2 allele homozygote	not applicable	Risk for gastric cancer among Asians	(Boccia et al., 2006)
GSTP1	SNP	+562 A/G	rs1695	Increased risk of esophageal squamous cell carcinoma and tended to be weakly, positively linked to cardia cancer	(Zendehdel et al., 2009)
NAT	RFLP	NAT1*10 allele	not applicable	Risk factor among the well-differentiated type of tumors of gastric cancer	(Katoh et al., 2000)
MET	SNP	-304C> A	undefined	Activate the transcription.	(Trzyna et al., 2012)
	SNP	206 C>G	undefined		
CDH1	SNP	-160 C/A	rs16260	Associated with DNA methylation linking genetic and epigenetics fields in the pathogenesis of gastritis.	(Borges Bdo et al., 2010)

Table 1. Selected single nucleotide polymorphisms (SNP) in inflammatory response genes, detoxification enzymes, and cancer-related processes associated to increase susceptibility of gastric cancer. The sign (-) indicates that the SNP is located in the direction of the promoter region, the sign (+) indicates that the SNP is located in the direction of the coding region, the number indicates the nucleotide position from TSS and the first nucleotide is substituted (/) by the second.

3. Epigenetic bases of gastritis

Epigenetic processes control the packaging and function of the human genome, and contribute to normal development and disease (Callinan et al. 2006). Epigenetic mechanisms such as DNA methylation, histone modifications and microRNAs are important events toward regulating gene expression for the biology and disease of the gastrointestinal tract. In particular, DNA methylation, a process in which cytosines acquire a methyl group in the 5' position only if they are followed by a guanine (Corvalán et al. 2010), virtually affects all of the pathways in the cellular network, such as DNA repair, cell cycle, and apoptosis (Esteller et

al. 2002). Furthermore, a growing body of evidence has shown that aberrant DNA methylation (DNA hypermethylation or DNA hypomethylation) is an early event in carcinogenesis (Jones et al. 2002; Herman et al. 2003; Kopelovich et al. 2003). In addition, aberrant DNA methylation has been recently considered to be an excellent candidate to explain how certain environmental factors may increase the risk of developing cancer (Chan et al. 2006). Accordingly, an emerging catalog of cancer-related genes inactivated by DNA hypermethylation in gastric cancer has been established (Hamilton et al. 2006; Wu et al. 2006; Bernal et al. 2008). However, there are limited reports on DNA hypermethylation analysis in precursor lesions of gastric cancer. Kang et al. (2001) tested five genes (p16, hMLH1, DAP-kinase, THBS1, and TIMP-3) in precancerous conditions to identify three different classes of hypermethylated genes. Hypermethylation of DAP-kinase was found in all premalignant conditions, whereas hMLH1 and p16 were preferentially methylated in intestinal metaplasia (6.3% and 2.1%, respectively), adenomas (9.8% and 11.5%, respectively) and gastric cancer (20.3% and 42.2%, respectively). THBS-1 and TIMP-3 were also hypermethylated at a similar frequency in all premalignant conditions, but showed a marked increase from chronic gastritis to intestinal metaplasia (10.1% & 34.7% and 14.5% & 36.7%, respectively; $P < 0.05$), as well as from adenomas to carcinomas (28.3% & 48.4% and 26.7% & 57.4% respectively). Another study including 11 genes (COX-2, DAP-kinase, E-cadherin, GSTP1, MGMT, hMLH1, p14, p16, THBS1, TIMP3, and RASSF1A) and 268 premalignant conditions (Kang et al. 2003) identified specific patterns of aberrant DNA methylation associated to aging. Specifically five genes (DAP-kinase, E-cadherin, p14, THBS1, and TIMP-3) showed a progressive increase in aberrant methylation frequency as a function of aging, whereas the other genes (COX-2, GSTP1, MGMT, hMLH1, p16, and RASSF1A) were rarely methylated. Male patients showed higher numbers of hypermethylated genes than females (3.2 vs. 2.1, respectively, $P = 0.002$) and cases with severe intestinal metaplasia also showed a higher frequency of aberrantly methylated genes. Taken together, these findings suggest that hypermethylation occurs early in the multistep gastric carcinogenesis and that accumulate during this process. In addition, age and gender are closely associated with increased frequency of aberrant DNA methylation. A further study demonstrated that DNA methylation of E-cadherin was associated with *H. pylori* infection ($p=0.002$) in an independent way of age and/or gastritis (Chan et al. 2003). Similarly, Maekita et al. (2006) carried out a more detailed analysis of the effect of *H. pylori* infection in the progression of aberrant DNA methylation in chronic gastritis. These authors collected gastric mucosa samples from 154 healthy volunteers (56 *H. pylori* negative and 98 *H. pylori* positive) and from 72 cases with gastric cancers (29 *H. pylori* negative and 43 *H. pylori* positive). Among the healthy volunteers, methylation levels were 5.4- to 303-fold higher in *H. pylori* positive subjects than in *H. pylori* negative ones ($P < 0.0001$). Particularly, methylation levels of the LOX, HAND1 and THBD at their promoter regions were identified in *H. pylori*-positive individuals. Among *H. pylori*-negative individuals, methylation levels were 2.2- to 32-fold higher in gastric cancer cases than in age-matched healthy volunteers. These findings suggest that *H. pylori* infection potently induces aberrant DNA methylation of several genes beyond E-cadherin, and that methylation levels of specific genes seemed to reflect the risk of gastric cancer in *H. pylori*-negative subjects. A recent quantitative analysis of the promoter region of a novel gene Reprimo (RPRM), per-

formed in Colombian residents from areas with high and low incidence of gastric cancer, demonstrated an association with virulent factors such as *cagA* and *vacA* s1/m1 regions of *H. pylori* strains. This data suggests that certain strains of *H. pylori* are associated with aberrant DNA methylation on specific genes (Schneider et al. 2010). Interestingly, we have shown that Reprimo is not only found in gastric mucosa but also in the plasma of gastric cancer patients (Bernal et al. 2008). Thus, hypermethylated circulating DNA offers the opportunity for non-invasive detection of gastric cancer and premalignant gastritis (Corvalán et al. 2010; Sapari et al. 2012). As mentioned before, an aberrant of DNA methylation could be considered as a candidate to explain how environmental factors increase the susceptibility of gastric cancer. One approach to explain this phenomena could be through the evaluation of the effect of the eradication of *H. pylori* in the pattern of DNA methylation of the gastric mucosa. Chen et al (2003) evaluate the presence of DNA methylation of E-cadherin promoter regions before the eradication of *H. pylori* and after 6 weeks of treatment. DNA methylation of E-cadherin was detected in 46% (19/41) and 17% (7/41) of both untreated and treated patients, respectively. Mucosal biopsy showed chronic inactive gastritis in 35 patients, intestinal metaplasia in one patient and normal mucosa in five patients after the treatment. This data suggests that *H. pylori* eradication therapy could reverse aberrant DNA methylation in patients with chronic gastritis. Similar results were reported by Leung et al. (2006), although they evaluated mucosal biopsies from the antrum and corpus of *H. pylori*-infected subjects at the baseline, and after one year of successful *H. pylori* eradication. In addition, these authors identified a significant reduction in the methylated density of the promoter region of the E-cadherin gene (Leung et al. 2006). Taken together, both reports are clear examples of how the environment affects DNA methylation and shows that the eradication of *H. pylori* infection reverses E-cadherin promoter hypermethylation. Although less explored, DNA hypomethylation can also be involved in the risk of developing cancer. The concept of DNA hypomethylation, although identified 30 years ago (Christman et al. 1977), was overlooked in preference to DNA hypermethylation. Recently, gene activation by promoter hypomethylation has been rediscovered (Ichinose et al. 1988; Cravo et al. 1996; Fang et al. 1996; Akiyama et al. 2003; Cho et al. 2003; Mesquita et al. 2003; Oshimo et al. 2003; Chalitchagorn et al. 2004; Honda et al. 2004; Kaneda et al. 2004; Yanagawa et al. 2004; Jung et al. 2005; Nishigaki et al. 2005). Emerging data suggests that global DNA hypomethylation is thought to occur during the early stages of gastric carcinogenesis (Goelz et al. 1985; Cravo et al. 1996; Narayan et al. 1998; Lin et al. 2001; Bariol et al. 2003) but hypomethylation of individual genes, such as MAGE, synuclein-alpha, MUC2, maspin, CAGE and family A melanoma antigen (Akiyama et al. 2003; Cho et al. 2003; Mesquita et al. 2003; Honda et al. 2004; Yanagawa et al. 2004; Jung et al. 2005) is thought to occur during the advanced stages. Interestingly, overexpression of well known oncogenes, such as MYC, HRAS1 and cyclin D2 has been linked to hypomethylation in their promoter region, suggesting an inverse correlation between methylation and gene expression (Fang et al. 1996; Oshimo et al. 2003). Taken together, this data suggests that promoter hypomethylation is an underlying mechanism of gene activation that should be explored in gastric carcinogenesis. In these sense, recent work from Daskalos et al (2011) has shown that the overexpression of p73, a well establish cancer-related gene associated with apoptosis and DNA repair (Zaika et al. 2011), might be associ-

ated with hypomethylation of the P2 promoter region that controls the DeltaNp73 isoform. Although this work was not performed in a gastric cancer model, our preliminary findings suggests similar results in gastric cancer cases (Corvalán et al. 2008). Based on both data, we decide to analysed the protein overexpression of p73 as well as other seven well established cancer-related genes activated by hypomethylation (BRCA1, HSP90, STAT1, FHIT, EGFR, p73, p53, p16INK4a) (Tanaka et al. 1998; Nien et al. 2007; Shutoh et al. 2009; Yu et al. 2009; Valdez et al. 2010; Daskalos et al. 2011; Rusiecki et al. 2011; Szaumkessel et al. 2011; Fornari et al. 2012). This analysis was performed in a large number of matched tumor/non-tumor adjacent mucosa of early gastric cancer as well as non-tumor cases at different stages of the multistep cascade of gastric cancer (Carrasco et al. 2010). Among 8 genes tested, only the overexpression of p73 was higher in matched tumor/non-tumor adjacent mucosa than in chronic gastritis/intestinal metaplasia (50.5% vs. 10.8%; $P < 0.0001$). We also assessed a detail histopathological analysis of the precursor lesions of gastric cancer through the Sydney and OLGA systems (Dixon et al. 1996; Corvalan et al. 2011; Rugge et al. 2011). As has been described (Rugge et al. 2010) only severe atrophy and OLGA stage IV were the most relevant features to identified high-risk gastritis (Fig. 4). However, the most interesting finding come after our attempt to integrated protein expression and histopathological assessments. This integration was performed by Significance Analysis of Microarrays (SAM), a multiple testing approach method that has been extensively applied in genomic research (Tusher et al. 2001) and confirmed by logistic regression (Nick et al. 2007). This integrative approach led us to identified that overexpression of p73 was even more significant than severe atrophy and OLGA stage IV in identifying high-risk premalignant gastritis (Carrasco et al. 2010). Therefore, we believe that the identification of the overexpression of p73 might contribute greatly to risk assessment of chronic gastritis to the development of gastric cancer.




Figure 3. The effect of the eradication of *H. pylori* in the pattern of DNA methylation as an example of environmental factors and the susceptibility of gastric cancer. (A) Before eradication of *H. pylori* (week 0), methylation was present in patients 3, 5 and 6. (B) Bisulfite sequencing confirm the methylated product. (C) After eradication of *H. pylori* (week 6), methylation was not present in any patient. (D) Bisulfite sequencing confirm the absence of methylated cytosine. MW: molecular weight marker, U: unmethylated band, M: methylated band, red color: unmethylated cytosines converted to thymidine, blue color: methylated cytosines. Taken from Chan et al., Gut 2006;55:463-8 with permission from BMJ Publishing Group Ltd.




Figure 4. Serial Analysis for Microarray from Non-tumor adjacent mucosa (NTAM) and chronic gastritis controls. NTAM group is significantly characterized by the overexpression of p73, OLGA stages III to IV, and severe atrophy (ATR-3), intestinal metaplasia (IM-3), and chronic inflammation (CI-3) according to the Sydney System. Control group cases were significantly characterized by lack of intestinal metaplasia (IM-0), atrophy (ATR-0), and chronic inflammation (CI-0). False discovery rate = 0. Taken from Carrasco et al., Clin.Cancer Res 2010;16:3253-9.

4. Conclusion

Data presented here suggests that genetic, epigenetic by means of DNA methylation might play a role in the dynamics of the progression of chronic gastritis and premalignant cascade to gastric cancer. DNA methylation works in both ways, inactivating or activating tumor-related genes through the hypermethylation or hypomethylation of the promoter regions of specific genes. The integration of these molecular bases of chronic gastritis with histopathological assessment by Sydney and OLGA systems will contribute to the better risk assessment for the development of gastric cancer.

Acknowledgements

Grant Support: Grants-in-Aid for Fondo Nacional de Ciencia y Tecnologia (FONDECYT) #1111014 and Fondo de Fomento al Desarrollo Científico y Tecnológico (FONDEF) #D09i1137 to A.H. Corvalan from the Government of Chile.

Author details

Alejandro H. Corvalan, Gonzalo Carrasco and Kathleen Saavedra

Ponfiticia Universidad Catolica de Chile, Chile

References

- [1] Akiyama Y, Maesawa C, Ogasawara S, et al. (2003). Cell-type-specific repression of the maspin gene is disrupted frequently by demethylation at the promoter region in gastric intestinal metaplasia and cancer cells. *Am J Pathol*, Vol.163; No.5: pp. 1911-1919. ISSN: 0002-9440.
- [2] Bariol C, Suter C, Cheong K, et al. (2003). The relationship between hypomethylation and CpG island methylation in colorectal neoplasia. *Am J Pathol*, Vol.162; No.4: pp. 1361-1371. ISSN: 0002-9440.
- [3] Bernal C, Aguayo FR, Villarroel C, et al. (2008). Reprimo as a potential biomarker for early detection in gastric cancer. *Clin Cancer Res*, Vol.14; No.19: pp.6264-6269. ISSN: 1078-0432..
- [4] Boccia S, De Lauretis A, Gianfagna F, et al. (2007). CYP2E1PstI/RsaI polymorphism and interaction with tobacco, alcohol and GSTs in gastric cancer susceptibility: A meta-analysis of the literature. *Carcinogenesis*, Vol.28; No.1: pp.101-106. ISSN: 0143-3334.
- [5] Borges Bdo N, Santos Eda S, Bastos CE, et al. (2010). Promoter polymorphisms and methylation of E-cadherin (CDH1) and KIT in gastric cancer patients from northern Brazil. *Anticancer Res*, Vol.30; No.6: pp.2225-2233. ISSN: 1791-7530.
- [6] Callinan PA and Feinberg AP. (2006). The emerging science of epigenomics. *Hum Mol Genet*, Vol.15 Spec No 1: pp.R95-101. ISSN: 0964-6906.
- [7] Capelle LG, de Vries AC, Haringsma J, et al. (2010). The staging of gastritis with the OLGA system by using intestinal metaplasia as an accurate alternative for atrophic gastritis. *Gastrointest Endosc*, Vol.71; No.7: pp.1150-1158. ISSN: 1097-6779.
- [8] Carrasco G, Diaz J, Valbuena JR, et al. (2010). Overexpression of p73 as a tissue marker for high-risk gastritis. *Clin Cancer Res*, Vol.16; No.12: pp.3253-3259. ISSN: 1078-0432.
- [9] Chalitchagorn K, Shuangshoti S, Hourpai N, et al. (2004). Distinctive pattern of LINE-1 methylation level in normal tissues and the association with carcinogenesis. *Oncogene*, Vol.23; No.54: pp.8841-8846. ISSN: 0950-9232.

- [10] Chan AO, Lam SK, Wong BC, et al. (2003). Promoter methylation of E-cadherin gene in gastric mucosa associated with *Helicobacter pylori* infection and in gastric cancer. *Gut*, Vol.52; No.4: pp.502-506. ISSN: 0017-5749.
- [11] Chan AO, Peng JZ, Lam SK, et al. (2006). Eradication of *Helicobacter pylori* infection reverses E-cadherin promoter hypermethylation. *Gut*, Vol.55; No.4: pp.463-468. ISSN: 0017-5749.
- [12] Cho B, Lee H, Jeong S, et al. (2003). Promoter hypomethylation of a novel cancer/testis antigen gene CAGE is correlated with its aberrant expression and is seen in premalignant stage of gastric carcinoma. *Biochem Biophys Res Commun*, Vol.307; No.1: pp.52-63. ISSN: 0006-291X.
- [13] Christman JK, Price P, Pedrinan L, et al. (1977). Correlation between hypomethylation of DNA and expression of globin genes in Friend erythroleukemia cells. *Eur J Biochem*, Vol.81; No.1: pp.53-61. ISSN: 0014-2956.
- [14] Coma del Corral MJ, Pardo-Mindan FJ, Razquin S, et al. (1990). Risk of cancer in patients with gastric dysplasia. Follow-up study of 67 patients. *Cancer*, Vol.65; No.9: pp.2078-2085. ISSN: 0008-543X.
- [15] Conchillo JM, Houben G, de Bruine A, et al. (2001). Is type III intestinal metaplasia an obligatory precancerous lesion in intestinal-type gastric carcinoma? *Eur J Cancer Prev*, Vol.10; No.4: pp.307-312. ISSN: 0959-8278.
- [16] Correa P, Cuello C, Duque E, et al. (1976). Gastric cancer in Colombia. III. Natural history of precursor lesions. *J Natl Cancer Inst*, Vol.57; No.5: pp.1027-1035. ISSN: 0027-8874.
- [17] Correa P, Haenszel W, Cuello C, et al. (1990). Gastric precancerous process in a high risk population: cross-sectional studies. *Cancer Res*, Vol.50; No.15: pp.4731-4736. ISSN:0008-5472.
- [18] Correa P and Houghton J. (2007). Carcinogenesis of *Helicobacter pylori*. *Gastroenterology*, Vol.133; No.2: pp.659-672. ISSN: 0016-5085.
- [19] Corvalán AH, Carrasco C and Olivares W (2011). Molecular Pathology of Gastritis. In: *Gastritis and Gastric Cancer - New insights in gasprotection, diagnosis and treatments*.
- [20] Rijeka, Croatia, Intech. ISBN:978-953-307-375-0.
- [21] Corvalán AH and Maturana MJ. (2010). Recent Patents of DNA Methylation Biomarkers in Gastrointestinal Oncology. *Recent Pat DNA Gene Seq*, Vol.4; No.3: pp. 202-209. ISSN: 1872-2156.
- [22] Corvalan A, Villarroel C, Ibanez P, et al. (2008). Demethylation of promoter region of p73 gene as a mechanism of overexpression of p73 protein in gastric carcinoma: Analysis by demethylation, microdissection of gastric epithelial cells and tissue mi-

- croarray. Proceedings of the 98th Annual Meeting of the American Association for Cancer Research; Washington, DC, USA. Abstract #4461
- [23] Cravo M, Pinto R, Fidalgo P, et al. (1996). Global DNA hypomethylation occurs in the early stages of intestinal type gastric carcinoma. *Gut*, Vol.39; No.3: pp.434-438. ISSN: 0017-5749.
- [24] Cuello C, Correa P, Haenszel W, et al. (1976). Gastric cancer in Colombia. I. Cancer risk and suspect environmental agents. *J Natl Cancer Inst*, Vol.57; No.5: pp. 1015-1020. ISSN: 0027-8874.
- [25] Daskalos A, Logotheti S, Markopoulou S, et al. (2011). Global DNA hypomethylation-induced DeltaNp73 transcriptional activation in non-small cell lung cancer. *Cancer Lett*, Vol.300; No.1: pp.79-86. ISSN: 1872-7980.
- [26] de Oliveira JG and Silva AE. (2012). Polymorphisms of the TLR2 and TLR4 genes are associated with risk of gastric cancer in a Brazilian population. *World J Gastroenterol*, Vol.18; No.11: pp.1235-1242. ISSN: 1007-9327.
- [27] Dixon MF, Genta RM, Yardley JH, et al. (1996). Classification and grading of gastritis. The updated Sydney System. International Workshop on the Histopathology of Gastritis, Houston 1994. *Am J Surg Pathol*, Vol.20; No.10: pp.1161-1181. ISSN:0147-5185 .
- [28] El-Omar EM, Carrington M, Chow WH, et al. (2000). Interleukin-1 polymorphisms associated with increased risk of gastric cancer. *Nature*, Vol.404; No.6776: pp.398-402. ISSN: 0028-0836.
- [29] El-Omar EM, Ng MT and Hold GL. (2008). Polymorphisms in Toll-like receptor genes and risk of cancer. *Oncogene*, Vol.27; No.2: pp.244-252. ISSN: 1476-5594.
- [30] Esteller M and Herman JG. (2002). Cancer as an epigenetic disease: DNA methylation and chromatin alterations in human tumours. *J Pathol*, Vol.196; No.1: pp.1-7. ISSN: 0022-3417.
- [31] Fang JY, Zhu SS, Xiao SD, et al. (1996). Studies on the hypomethylation of c-myc, c-Ha-ras oncogenes and histopathological changes in human gastric carcinoma. *J Gastroenterol Hepatol*, Vol.11; No.11: pp.1079-1082. ISSN: 0815-9319.
- [32] Figueiredo C, Machado JC, Pharoah P, et al. (2002). Helicobacter pylori and interleukin 1 genotyping: an opportunity to identify high-risk individuals for gastric carcinoma. *J Natl Cancer Inst*, Vol.94; No.22: pp.1680-1687. ISSN: 0027-8874.
- [33] Fornari F, Milazzo M, Chieco P, et al. (2012). In hepatocellular carcinoma miR-519d is up-regulated by p53 and DNA hypomethylation and targets CDKN1A/p21, PTEN, AKT3 and TIMP2. *J Pathol*, Vol.227; No.3: pp.275-285. ISSN: 1096-9896.
- [34] Gherardi E, Birchmeier W, Birchmeier C, et al. (2012). Targeting MET in cancer: rationale and progress. *Nat Rev Cancer*, Vol.12; No.2: pp.89-103. ISSN: 1474-1768.

- [35] Goelz SE, Vogelstein B, Hamilton SR, et al. (1985). Hypomethylation of DNA from benign and malignant human colon neoplasms. *Science*, Vol.228; No.4696: pp. 187-190. ISSN: 0036-8075.
- [36] Gonzalez CA, Sala N and Capella G. (2002). Genetic susceptibility and gastric cancer risk. *Int J Cancer*, Vol.100; No.3: pp.249-260. ISSN: 0020-7136.
- [37] Haenszel W, Correa P, Cuello C, et al. (1976). Gastric cancer in Colombia. II. Case-control epidemiologic study of precursor lesions. *J Natl Cancer Inst*, Vol.57; No.5: pp. 1021-1026. ISSN: 0027-8874.
- [38] Hamilton JP, Sato F, Jin Z, et al. (2006). Reprimo methylation is a potential biomarker of Barrett's-Associated esophageal neoplastic progression. *Clin Cancer Res*, Vol.12; No.22: pp.6637-6642. ISSN: 1078-0432.
- [39] Herman JG and Baylin SB. (2003). Gene silencing in cancer in association with promoter hypermethylation. *N Engl J Med*, Vol.349; No.21: pp.2042-2054. ISSN: 1533-4406.
- [40] Hiyama T, Tanaka S, Kitadai Y, et al. (2002). p53 Codon 72 polymorphism in gastric cancer susceptibility in patients with *Helicobacter pylori*-associated chronic gastritis. *Int J Cancer*, Vol.100; No.3: pp.304-308. ISSN: 0020-7136.
- [41] Honda T, Tamura G, Waki T, et al. (2004). Demethylation of MAGE promoters during gastric cancer progression. *Br J Cancer*, Vol.90; No.4: pp.838-843. ISSN: 0007-0920.
- [42] Hull J, Thomson A and Kwiatkowski D. (2000). Association of respiratory syncytial virus bronchiolitis with the interleukin 8 gene region in UK families. *Thorax*, Vol.55; No.12: pp.1023-1027. ISSN: 0040-6376.
- [43] Hwang IR, Kodama T, Kikuchi S, et al. (2002). Effect of interleukin 1 polymorphisms on gastric mucosal interleukin 1beta production in *Helicobacter pylori* infection. *Gastroenterology*, Vol.123; No.6: pp.1793-1803. ISSN: 0016-5085.
- [44] Ichinose M, Miki K, Tatematsu M, et al. (1988). Cell-specific hypomethylation of the pepsinogen gene in pepsinogen-producing cells. *Biochem Biophys Res Commun*, Vol.155; No.2: pp.670-677. ISSN: 0006-291X.
- [45] Jones PA and Baylin SB. (2002). The fundamental role of epigenetic events in cancer. *Nat Rev Genet*, Vol.3; No.6: pp.415-428. ISSN: 1471-0056.
- [46] Jung EJ, Kim MA, Lee HS, et al. (2005). Expression of family A melanoma antigen in human gastric carcinoma. *Anticancer Res*, Vol.25; No.3B: pp.2105-2111. ISSN: 0250-7005.
- [47] Kaneda A, Tsukamoto T, Takamura-Enya T, et al. (2004). Frequent hypomethylation in multiple promoter CpG islands is associated with global hypomethylation, but not with frequent promoter hypermethylation. *Cancer Sci*, Vol.95; No.1: pp.58-64. ISSN: 1347-9032.

- [48] Kang GH, Lee HJ, Hwang KS, et al. (2003). Aberrant CpG island hypermethylation of chronic gastritis, in relation to aging, gender, intestinal metaplasia, and chronic inflammation. *Am J Pathol*, Vol.163; No.4: pp.1551-1556. ISSN: 0002-9440.
- [49] Kang GH, Shim YH, Jung HY, et al. (2001). CpG island methylation in premalignant stages of gastric carcinoma. *Cancer Res*, Vol.61; No.7: pp.2847-2851. ISSN: 0008-5472.
- [50] Katoh T, Boissy R, Nagata N, et al. (2000). Inherited polymorphism in the N-acetyltransferase 1 (NAT1) and 2 (NAT2) genes and susceptibility to gastric and colorectal adenocarcinoma. *Int J Cancer*, Vol.85; No.1: pp.46-49. ISSN: 0020-7136.
- [51] Koch HK, Oehlert M and Oehlert W. (1990). An evaluation of gastric dysplasia in the years 1986 and 1987. *Pathol Res Pract*, Vol.186; No.1: pp.80-84. ISSN: 0344-0338.
- [52] Kopelovich L, Crowell JA and Fay JR. (2003). The epigenome as a target for cancer chemoprevention. *J Natl Cancer Inst*, Vol.95; No.23: pp.1747-1757. ISSN: 1460-2105.
- [53] Leung WK, Man EP, Yu J, et al. (2006). Effects of *Helicobacter pylori* eradication on methylation status of E-cadherin gene in noncancerous stomach. *Clin Cancer Res*, Vol.12; No.10: pp.3216-3221. ISSN: 1078-0432.
- [54] Lin CH, Hsieh SY, Sheen IS, et al. (2001). Genome-wide hypomethylation in hepatocellular carcinogenesis. *Cancer Res*, Vol.61; No.10: pp.4238-4243. ISSN: 0008-5472.
- [55] Maekita T, Nakazawa K, Mihara M, et al. (2006). High levels of aberrant DNA methylation in *Helicobacter pylori*-infected gastric mucosae and its possible association with gastric cancer risk. *Clin Cancer Res*, Vol.12; No.3 Pt 1: pp.989-995. ISSN: 1078-0432.
- [56] Mesquita P, Peixoto AJ, Seruca R, et al. (2003). Role of site-specific promoter hypomethylation in aberrant MUC2 mucin expression in mucinous gastric carcinomas. *Cancer Lett*, Vol.189; No.2: pp.129-136. ISSN: 0304-3835.
- [57] Narayan A, Ji W, Zhang XY, et al. (1998). Hypomethylation of pericentromeric DNA in breast adenocarcinomas. *Int J Cancer*, Vol.77; No.6: pp.833-838. ISSN: 0020-7136.
- [58] Nick TG and Campbell KM. (2007). Logistic regression. *Methods Mol Biol*, Vol.404: pp.273-301. ISSN: 1064-3745.
- [59] Nien WL, Dauphinee SM, Moffat LD, et al. (2007). Overexpression of the mTOR alpha4 phosphoprotein activates protein phosphatase 2A and increases Stat1alpha binding to PIAS1. *Mol Cell Endocrinol*, Vol.263; No.1-2: pp.10-17. ISSN: 0303-7207.
- [60] Nishigaki M, Aoyagi K, Danjoh I, et al. (2005). Discovery of aberrant expression of R-RAS by cancer-linked DNA hypomethylation in gastric cancer using microarrays. *Cancer Res*, Vol.65; No.6: pp.2115-2124. ISSN:0008-5472.
- [61] Ohata H, Kitauchi S, Yoshimura N, et al. (2004). Progression of chronic atrophic gastritis associated with *Helicobacter pylori* infection increases risk of gastric cancer. *Int J Cancer*, Vol.109; No.1: pp.138-143. ISSN: 0020-7136.

- [62] Oshimo Y, Nakayama H, Ito R, et al. (2003). Promoter methylation of cyclin D2 gene in gastric carcinoma. *Int J Oncol*, Vol.23; No.6: pp.1663-1670. ISSN: 1019-6439.
- [63] Ramesar KC, Sanders DS and Hopwood D. (1987). Limited value of type III intestinal metaplasia in predicting risk of gastric carcinoma. *J Clin Pathol*, Vol.40; No.11: pp. 1287-1290. ISSN: 0021-9746.
- [64] Rokkas T, Filipe MI and Sladen GE. (1991). Detection of an increased incidence of early gastric cancer in patients with intestinal metaplasia type III who are closely followed up. *Gut*, Vol.32; No.10: pp.1110-1113. ISSN: 0017-5749.
- [65] Rugge M, Cassaro M, Di Mario F, et al. (2003). The long term outcome of gastric non-invasive neoplasia. *Gut*, Vol.52; No.8: pp.1111-1116. ISSN: 0017-5749.
- [66] Rugge M, de Boni M, Pennelli G, et al. (2010). Gastritis OLGA-staging and gastric cancer risk: a twelve-year clinico-pathological follow-up study. *Aliment Pharmacol Ther*, Vol.31; No.10: pp.1104-1111. ISSN: 1365-2036.
- [67] Rugge M, Kim JG, Mahachai V, et al. (2008). OLGA gastritis staging in young adults and country-specific gastric cancer risk. *Int J Surg Pathol*, Vol.16; No.2: pp.150-154. ISSN: 1066-8969.
- [68] Rugge M, Meggio A, Pennelli G, et al. (2007). Gastritis staging in clinical practice: the OLGA staging system. *Gut*, Vol.56; No.5: pp.631-636. ISSN: 0017-5749.
- [69] Rugge M, Pennelli G, Piloizzi E, et al. (2011). Gastritis: the histology report. *Dig Liver Dis*, Vol.43 Suppl 4: pp.S373-384. ISSN: 1878-3562.
- [70] Ruggiero P. (2012). *Helicobacter pylori* infection: what's new. *Curr Opin Infect Dis*, Vol.25; No.3: pp.337-344. ISSN: 1473-6527.
- [71] Rusiecki JA, Al-Nabhani M, Tarantini L, et al. (2011). Global DNA methylation and tumor suppressor gene promoter methylation and gastric cancer risk in an Omani Arab population. *Epigenomics*, Vol.3; No.4: pp.417-429. ISSN: 1750-192X.
- [72] Sapari NS, Loh M, Vaithilingam A, et al. (2012). Clinical potential of DNA methylation in gastric cancer: a meta-analysis. *PLoS One*, Vol.7; No.4: pp.e36275. ISSN: 1932-6203.
- [73] Saraga EP, Gardiol D and Costa J. (1987). Gastric dysplasia. A histological follow-up study. *Am J Surg Pathol*, Vol.11; No.10: pp.788-796. ISSN: 0147-5185.
- [74] Schneider BG, Peng DF, Camargo MC, et al. (2010). Promoter DNA hypermethylation in gastric biopsies from subjects at high and low risk for gastric cancer. *Int J Cancer*, Vol.127; No.11: pp.2588-2597. ISSN: 1097-0215.
- [75] Shutoh Y, Takeda M, Ohtsuka R, et al. (2009). Low dose effects of dichlorodiphenyl-trichloroethane (DDT) on gene transcription and DNA methylation in the hypothalamus of young male rats: implication of hormesis-like effects. *J Toxicol Sci*, Vol.34; No. 5: pp.469-482. ISSN: 1880-3989.

- [76] Silva S, Filipe MI and Pinho A. (1990). Variants of intestinal metaplasia in the evolution of chronic atrophic gastritis and gastric ulcer. A follow up study. *Gut*, Vol.31; No.10: pp.1097-1104. ISSN: 0017-5749.
- [77] Szaumkessel M, Richter J, Giefing M, et al. (2011). Pyrosequencing-based DNA methylation profiling of Fanconi anemia/BRCA pathway genes in laryngeal squamous cell carcinoma. *Int J Oncol*, Vol.39; No.2: pp.505-514. ISSN: 1791-2423.
- [78] Tahara T, Arisawa T, Shibata T, et al. (2009). Effect of polymorphisms in the 3'-untranslated region (3'-UTR) of VEGF gene on gastric pre-malignant condition. *Anti-cancer Res*, Vol.29; No.2: pp.485-489. ISSN: 0250-7005.
- [79] Tanaka H, Shimada Y, Harada H, et al. (1998). Methylation of the 5' CpG island of the FHIT gene is closely associated with transcriptional inactivation in esophageal squamous cell carcinomas. *Cancer Res*, Vol.58; No.15: pp.3429-3434. ISSN: 0008-5472.
- [80] Trzyna E, Duleba M, Faryna M, et al. (2012). Regulation of transcription in cancer. *Front Biosci*, Vol.17: pp.316-330. ISSN: 1093-4715.
- [81] Tusher VG, Tibshirani R and Chu G. (2001). Significance analysis of microarrays applied to the ionizing radiation response. *Proc Natl Acad Sci U S A*, Vol.98; No.9: pp. 5116-5121. ISSN: 0027-8424.
- [82] Valdez BC, Li Y, Murray D, et al. (2010). 5-Aza-2'-deoxycytidine sensitizes busulfan-resistant myeloid leukemia cells by regulating expression of genes involved in cell cycle checkpoint and apoptosis. *Leuk Res*, Vol.34; No.3: pp.364-372. ISSN: 1873-5835.
- [83] Vannella L, Lahner E and Annibale B. (2012). Risk for gastric neoplasias in patients with chronic atrophic gastritis: a critical reappraisal. *World J Gastroenterol*, Vol.18; No.12: pp.1279-1285. ISSN: 1007-9327.
- [84] Whiting JL, Sigurdsson A, Rowlands DC, et al. (2002). The long term results of endoscopic surveillance of premalignant gastric lesions. *Gut*, Vol.50; No.3: pp.378-381. ISSN: 0017-5749.
- [85] Wu DL, Sui FY, Jiang XM, et al. (2006). Methylation in esophageal carcinogenesis. *World J Gastroenterol*, Vol.12; No.43: pp.6933-6940. ISSN: 1007-9327.
- [86] Xue H, Liu J, Lin B, et al. (2012). A meta-analysis of interleukin-8 -251 promoter polymorphism associated with gastric cancer risk. *PLoS One*, Vol.7; No.1: pp.e28083. ISSN: 1932-6203.
- [87] Yanagawa N, Tamura G, Honda T, et al. (2004). Demethylation of the synuclein gamma gene CpG island in primary gastric cancers and gastric cancer cell lines. *Clin Cancer Res*, Vol.10; No.7: pp.2447-2451. ISSN: 1078-0432.
- [88] Yu Y, Kanwar SS, Patel BB, et al. (2009). Elimination of Colon Cancer Stem-Like Cells by the Combination of Curcumin and FOLFOX. *Transl Oncol*, Vol.2; No.4: pp. 321-328. ISSN: 1936-5233.

- [89] Yuzhalin A. (2011). The role of interleukin DNA polymorphisms in gastric cancer. *Hum Immunol*, Vol.72; No.11: pp.1128-1136. ISSN: 1879-1166.
- [90] Zaika E, Wei J, Yin D, et al. (2011). p73 protein regulates DNA damage repair. *FASEB J*, Vol.25; No.12: pp.4406-4414. ISSN: 1530-6860.
- [91] Zendejdel K, Bahmanyar S, McCarthy S, et al. (2009). Genetic polymorphisms of glutathione S-transferase genes GSTP1, GSTM1, and GSTT1 and risk of esophageal and gastric cardia cancers. *Cancer Causes Control*, Vol.20; No.10: pp.2031-2038. ISSN: 1573-7225.
- [92] Zhou Y, Li N, Zhuang W, et al. (2007). P53 codon 72 polymorphism and gastric cancer: a meta-analysis of the literature. *Int J Cancer*, Vol.121; No.7: pp.1481-1486. ISSN: 0020-7136.

