

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,400

Open access books available

133,000

International authors and editors

165M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



---

# The Genetic Architecture of Autism and Related Conditions

---

Michael Gill, Graham Kenny and Richard Anney

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/54854>

---

## 1. Introduction

Autism is a behaviourally defined developmental disorder characterised by impairments in social communication, restricted interests and repetitive behaviours [1]. Abnormalities in these three developmental areas tend to cluster together in affected individuals. In DSM-IV, Autism is part of a larger continuum of disorders collectively called Pervasive Developmental Disorders. Autism spectrum disorders (ASD) refer to Autism, Pervasive developmental disorder, not otherwise specified, and Asperger syndrome. All individuals with ASDs have qualitative abnormalities of social development in combination with disorders of communication and/or stereotyped repetitive interests and behaviors. The social skills that develop naturally in typically-developing children do not do so in children with ASD. In addition, there are several behaviors and co-morbid symptoms that relate to each of the three classical impairments. Recent studies have reported rates of co-occurring intellectual disability in the range of 25-50%. Neither developmental delay nor cognitive impairment are required for an ASD diagnosis.

Fombonne and colleagues recently estimated the prevalence of strictly-defined autism at approximately 15-20 per 10,000 people [2]. When the definition of autism is relaxed to include Autism Spectrum Disorders, the prevalence estimated expands to approximately 60 in 10,000 children [2, 3].

Little is known of the biological basis of ASD and the future development of rational knowledge based treatments will depend on a comprehensive understanding of innate biological predisposition and its interaction with environmental factors. The identification and characterisation of the genetic variation and genes involved in ASD is a route towards this goal. This chapter outlines the various approaches that have been applied to this task, in the context of rapidly evolving technology and human genome resources, and summarises the state of

knowledge at this time, anticipating future developments, and outlining the implications for clinical management.

## 2. Autism is a heritable disorder

There is unequivocal evidence to support the role of genetic liability as the basis of familiarity in the aetiology of ASD. In their seminal work, Folstein and Rutter [4] observed that 4 of 11 (36%) pairs of identical twins were concordant for strictly defined autism – whilst none of 10 pairs of fraternal twins were concordant. In a large follow-up study, which included these original twins, Bailey and colleagues observed a more striking concordance rate in identical twins (60%) with no concordance in the fraternal twins [5]. This highly cited study underlined ASD as the most heritable of the neurodevelopmental and psychiatric disorders with heritability estimates of 91 - 93%. A recent review of over 30 twin studies of ASD [6] noted that the early twin studies, which relied upon a strict diagnosis of autism, show a median identical and fraternal concordance of 76% and 0% respectively. The expansion of these and other studies to include a broader definition of ASD revealed a median concordance rate of 88% and 31%. In both cases, these data reveal that ASD is underpinned by a strong genetic component. More recently, Hallmayer and colleagues [7] using more stringent clinical diagnostic tools, such as the Autism Diagnostic Interview (ADI; [8]) and the Autism Diagnostic Observation Schedule (ADOS; [9]) in a group of 202 twin pairs from the California Twin Registry, identified similar concordance rate of ASD as previously attained; with identical twin boys showing a concordance rate of 77% and fraternal twin boys 31%.

## 3. Genetic linkage and association

The heritability estimates above calculated from twin studies imply that there is unknown genetic variation in children with ASD that confers risk. Several models may fit the observed family and twin data including, at the extremes, a ‘common disorder-common-variant’ model, with many DNA variants of small effect combining together to create risk; and a multiple rare variant model where ASD is in reality a large collection of different and individually rare disorders. The true picture is only slowly emerging, and as with many other common disorders, common low penetrant and rare higher penetrant DNA variants combine to jointly contribute to risk. Molecular genetic methods have developed exponentially over the last few decades with ever increasing knowledge of the structure and function of the human genome. For disorders with largely unknown aetiology, such as ASD, using human genetic studies to identify risk genes is a gateway to an understanding of underlying biology. Such studies may focus on individuals or families each using different but related genetic strategies to track down risk or causative genetic variation. These methods include genetic linkage, genetic association, and direct examination of structural and copy number variation.

The basis of genetic linkage is the physical co-location of genes or other DNA variation. Such DNA variation is more likely to be transmitted together (or linked) than those that are further

apart. Linkage studies take advantage of this non-random assortment of genetic variation. A linkage study calculates whether a known genetic variant and a disease mutation (represented by the disease trait) are linked and if so, roughly localises the causative mutation. Following the successes of using these approaches in the discovery of multiple loci implicate in Mendelian disorders, researchers were encouraged to apply linkage methodology to more complex traits, such as ASD where Mendelian principles may apply, at least in a proportion of families. However, in ASD, only a few scans have highlighted loci with *significant* linkage, highlighting loci including chromosome 7q, 2q and 3q.

In 1998, the International Molecular Genetic Study of Autism Consortium (IMGSAC) [10] reported modest evidence for linkage. This included *significant* linkage arising on chromosome 7q32-q34. Supplemented analyses of IMGSAC families provided additional support for linkage at 7q22, 16p13, and 2q31 [11, 12]. The long arm of chromosome 7 has received particular attention with additional support reported for 7q21 [13], 7q22 [14], 7q31 [15, 16], 7q32 [17] and 7q36 [16].

Few individual families exist where ASD segregates in an obvious Mendelian fashion that is large enough to provide significant evidence for linkage by themselves. Most linkage studies required the assumption that a significant proportion of the families in the sample might be linked to a given locus and few were sufficiently large to accommodate even modest locus heterogeneity. Under a 'common disorder - common variant' model, multiply affected families will occur but linkage methods would be considerably underpowered. The effects of DNA variation with low penetrance are more easily identified using a genetic association study design in a sample of cases drawn from a population.

Fine-mapping and candidate gene association studies at implicated regions on 7q have implicated a number of potential susceptibility genes including *RELN*, *MET*, *CNTNAP2* and *EN2*. Persico and colleagues [18] studied five DNA variants or polymorphisms across the *RELN* gene locus, including a GGC repeat variant located close to the *RELN* gene translation initiator codon. Located at 7q22, *RELN* encodes an extracellular matrix protein Reelin, which plays a pivotal role in the development of laminar structures including cerebral cortex, cerebellum and hippocampus. Using a genetic association approach, Persico and colleagues identified a nominally significant association with this 5'-UTR GCC-triplet-repeat polymorphism [18]. This finding was further supported by some studies [19-25] but not others [26-31]. *MET*, located at 7q31 received considerable attention following a high-profile association reported by Campbell and colleagues. The *MET* gene encodes a protein involved in MET (Mesenchymal epithelial transition factor) receptor tyrosine kinase signalling which has been implicated in brain growth and maturation – offering biological plausibility to its candidature. As with other candidate genes in ASD, the original findings have been supported in some [32], but not other studies [33] [34]. A similar scenario played out for the *EN2* homeobox gene located at 7q36 [35, 36]. Arking and colleagues [37] observed an association at *CNTNAP2* in the NIHM/AGRE collection. The main association observed was for rs7794745 located in intron 2 of the gene (*Discovery P* = 0.00002, *Validation P* = 0.005). The *CNTNAP2* (7q35) gene encodes the contactin-associated protein-like 2 protein, which is a member of the neurexin family and thought to play a role in axonal differentiation and guidance. Li and colleagues also found mild support for *CNTNAP2*

where they observed a weak association with a haplotype containing the SNP rs7794745 [38]. More recently, a large GWAS of 2705 families identified a strong association for the SNP rs1718101 in the *CNTNAP2* gene with a subset of individuals without intellectual disability and of European ancestry ( $P = 7.78 \times 10^{-9}$ ) [39].

The Paris Autism Research International Sibpair (PARIS) study [15], that supported the findings on 7q, noted additional loci at 2q31-q32, 16p13 and 19p13, that overlapped to some extent with regions identified in the IMGSAC analyses. Fine mapping of the 2q31-q32 led to additional evidence implicating the genes *SLC25A12*, *STK39* and *ITGA4* as putative risk genes for ASD. As with most candidate genes examined in small samples, these associations were validated by some [40-42] but not all studies [43-46]. The positional candidature of the 2q31-q32 was further supported by a case study of a young Irish male with high-functioning autism with a complex translocation traversing chromosome 2q32 (46,XY,t(9;2)(q31.1;q32.2q31.3)) [47]. Fine mapping and mutation analysis identified an association with a polymorphism within the splice donor sequence of exon 16 of the *ITGA4* gene and ASD (rs12690517;  $P = 0.008$ ) [48].

In a study of Finnish families, Auranen and colleagues [49, 50], who included individuals with autism, infantile autism, Asperger syndrome (AS) and developmental dysphasia reported a *significant* linkage to 3q25-q27. This was supported by *suggestive* evidence at 3q25-q27 in a study of AS also in the Finnish population [51] as well as linkage at 3q25-q27 in a single large extended Utah pedigree of Northern European ancestry [52].

Interestingly, the advent of large collaborative studies by the Autism Genome Research Exchange (AGRE), the Autism Genome Project (AGP) and the National Institute of Mental Health (NIMH) has not yielded stronger evidence in favour of specific loci. Liu and colleagues using data from 110 multiplex families from the AGRE collection [16] observed only *suggestive* linkage on chromosomes 5p13, Xq26-qtel, and 19q12 alongside modest support for previously reported linkage on 7q (7q31 and 7q36) and 16p13. A follow-up analysis including 235 additional AGRE multiplex families was again limited to only *suggestive* loci at chromosomes 17q11, 5p13, 11p11-p13, 4q21-q22 and 8q24 [53]. In a much larger collection of 1168 multiply-affected families from the AGP (including families previously included in the AGRE, CPEA, IMGSAC, PARIS and Seaver linkage studies), Szatmari and colleagues identified *suggestive* linkage to chromosome 11p12-p13 and a large region on chromosome 15q23-q25 [14]. Of the regions that were featured prominently in previous linkage analyses, there was only modest support for previously highlighted linkage regions on chromosome 2q31 (female autism-probands) and 7q22 (male ASD-probands) from families of European ancestry [14]. A similarly large linkage study of 1031 families, including 1553 affected offspring, Weiss and colleagues identified *suggestive* linkage at 6p27 and *significant* linkage at 20p13 [54].

Early association studies examined whether specific variation in genes was associated with the disease and focused on candidate loci identified from linkage and cytogenetic studies (positional candidate genes) as well as in genes within biological processes that we perceived as having a role in ASD. In the mid-1990's Risch and Merikangas [55] demonstrated that where genetic variants have only small effect on risk the association study is a more powerful ap-

proach than linkage to identify genetic risk. However, the transition from candidate gene to genome-wide association studies was not realised until technological advances firstly enriched the maps of common variation across the genome and secondly enabled the interrogation of these variants *en masse* as part of genome-wide SNP arrays.

Wang and colleagues [56] performed one of the first GWAS on individuals in European ancestry individuals from the AGRE collection, the Autism Case Control Collection and unaffected controls from the Children's Hospital of Philadelphia control collection. Neither family-based nor case-control analysis alone yielded genome-wide significant findings. However, in a combined analyses the authors identified genome-wide significant association on chromosome 5p14.1 (rs4307059;  $P = 3.4 \times 10^{-8}$ ) and a number of additional association signals on chromosome 13q33.3, 14q21.1 and Xp22.32. The 5p14.1 association was validated in the Collaborative Autism Project (CAP) and Centre for Autism Research and Treatment (CART) study. The authors found a modest to strong replication of the association signal on chromosome 5p14.1. In a reciprocal study with the CAP and CART study as the discovery sample followed by validation using the AGRE dataset was published in parallel by Ma and colleagues [57]. The authors examined approximately 500k SNPs, more than in the Wang report and albeit not genome-wide significant, they retained the association signal on chromosome 5p14.1.

A second independent GWAS was reported by Weiss and colleagues [54]. In the initial scan the authors did not find any GW-significant associations. However, as with the previous GWAS, additional supplementation of their family-based studies with a case-control set derived from 90 probands without parental data garnered some additional signal for the top hits. A replication consortium of greater than 2000 trios was genotyped for 45 SNPs across all of the top associated regions. The only marker that showed evidence of replication resides on the short arm of chromosome 5 at 5p15. Although, like Ma and colleagues, this report has considerable overlap with the AGRE families reported by Wang and colleagues, the authors did not observe strong association at 5p14.1. The chromosome 5p association lies in close proximity to *TAS2R1*. The *TAS2R1* gene encodes a G-protein coupled receptor that is involved in bitter taste recognition. The authors highlight a more biologically plausible ASD candidate gene approximately 80 Kb telomeric, *SEMA5A*. *SEMA5A* encodes a gene important in axonal guidance and has been shown to be down regulated in the occipital lobe cortex, lymphoblast cell lines and lymphocytes from individuals with autism.

Two additional GWAS followed by the Autism Genome Project (AGP) [39, 58]. In the first report of 1369 families, Anney and colleagues identified a single GW-significant finding on chromosome 20 at position 20p12 within the *MACROD2* (MACRO-domain containing 2) gene locus (rs4141463;  $P = 2.1 \times 10^{-8}$ ; OR = 0.56). Weak statistical support was observed for *MACROD2* in an AGRE validation sample, albeit showing the same direction of effect for the risk allele. In a follow-up study using an additional 1301 families the authors showed little if any signal of association (rs4141463;  $P = 0.206$ , OR=0.91). In a combined analyses the association at *MACROD2* was less compelling (rs4141463;  $P = 1.2 \times 10^{-6}$ ; OR=0.77). The role of *MACROD2* is largely unknown although structural variation in this gene and specifically the region harbouring the ASD association signal have been implicated in schizophrenia [59, 60] and epilepsy [61]. However, this same region has been identified as a hotspot for deletions in the genome

identified as a region of large rare deletions [62]. MACROD-like proteins are highly conserved across evolutionary time, which may indicate an essential role. The MACRO-domain is an ADP-ribose binding module [63] and has been implicated in the ADP-ribosylation of proteins, an important post-translational modification that occurs in a variety of biological processes such as DNA repair [64], heterochromatin formation, histone modification and sirtuin biology [65-67] as well as long-term memory formation [68]. The association signal observed at rs4141463, albeit tagged to the *MACROD2* gene, resides in an intronic region near an intragenic non-protein-coding-RNA, *NCRNA00186 (MACROD2-AS1)*. Two *MACROD2-AS1* transcripts have been reported of 673bp and 1230bp in length, located on the reverse strand between exon 5 and 6 of *MACROD2*. Anti-sense RNAs typically interact with mRNA, resulting in transcriptional or post-transcriptional effects and have been linked to brain development and plasticity [69]. However, unlike *MSNP1AS* described for the 5p14.1 association observed by Wang and colleagues, a function for this non-coding RNA has not yet been reported.

The strongest association signal observed by the AGP combined analyses was rs1718101 ( $P = 7.8 \times 10^{-9}$ ; OR=2.13 (1.63-2.80)), a SNP within the *CNTNAP2* gene which was previously implicated in ASD through linkage analyses. This association was observed in a secondary analysis which was restricted to ASD individuals of European ancestry with a higher IQ. Anney and colleagues suggest that with the current data “few if any common variants have an impact on risk exceeding (an effect size of) 1.2 (or below its inverse).” In an attempt to seek evidence for or against common variants having an impact on risk, the authors constructed an allele-score. The allele-score method, as previously described by Purcell and colleagues [70], calculates a score for each individual based on number of risk associated alleles that the individual possesses. This score is then used to either calculate the predictive value of the score between cases and controls or estimate the amount of variance that this score predicts for the disease. Allele-scores derived from the transmission of common alleles from the families described in the AGP stage 1 GWAS [58] could significantly predict case-status in the independent Stage 2 sample. The authors concluded however that despite the limited findings for individual loci from GWAS studies to date, *en masse* the top results exert a detectable impact suggesting that as the sample sizes increase, additional significant loci will emerge. Putting together samples of the size seen in successful GWAS studies in other disorders ( $n > 20,000$ ) is challenging for a disorder with a complex phenotype such as ASD.

#### 4. Structural variation: Chromosomal abnormalities in autism

Autism has been frequently associated with chromosome abnormalities, such as deletions, duplications, inversions, or translocations; with abnormalities on the long arm of Chromosome 15 and with numerical and structural abnormalities of the sex chromosomes. Reddy in a survey of chromosomal abnormality in autism found one in 14 of 421 individuals (3.33%) [71]. These fourteen cases broke down into 4 supernumerary chromosome markers, 4 deletions, 3 inversions and 3 duplications. Several reviews have since confirmed that such abnormalities can be identified in ~3-5% of these patients [72, 73]. The regions commonly reported include 2q37,

5p14, 5p15, multiple locations on chromosome 7, 11q25, 15q11-q13, 16q22.3, 17p11.2, 18q21.1, 18q23, 22q11.2, 22q13.3 and Xp22.2–p22.3.

Many studies have converged on particular chromosomal abnormalities in autism, the most common of which are maternally inherited duplications at 15q11-13. These duplications are found in as many as 1–3% of patients diagnosed with autism.

Another relatively common chromosomal variant is the 22q11.2 microdeletion syndrome, or velocardiofacial syndrome (VCFS) that occurs in ~1/4000 live births. This syndrome is also identified in the context of learning disability or in schizophrenia and has a complex phenotypic expression affecting multiple organs. The physical features include a typical facial appearance, (long face, narrow palpebral fissures, flattened malar eminences, prominent nose and small mouth); anatomical and/or functional abnormalities of the palatal shelves such as cleft palate and velopharyngeal insufficiency; lymphoid tissue hypoplasia and heart defects. A wide range of childhood onset developmental symptoms and disorders are described in association with the VCFS mutation [74] including attention deficit hyperactivity disorder (ADHD), oppositional defiant disorder, phobias, anxiety, obsessive compulsive disorder and autism spectrum disorders.

## 5. Copy number variation

The human genome has both sequence and structural variation, the majority of which has no functional consequences and is unrelated to any disorder. Structural variation may be balanced, such as is the case for inversions and balanced translocations or may alter DNA copy number. The latter, referred to as copy number variants (CNVs) extend from the duplication or deletion of a single base pair to whole chromosome abnormalities. The term CNV is generally used to indicate the larger changes. Initially, large CNVs were identified with classical chromosomal staining and light microscopy. As many of these abnormalities were initially identified in sub-telomeric regions, there was interest in knowing if such deletions, duplications and other re-arrangements might occur throughout the genome. This hypothesis was confirmed with the development of high resolution array based comparative genomic hybridization (aCGH) [75].

In the arrays, comparative hybridization is performed with DNA immobilized on a platform such as a glass slide. Initially the DNA arrays consisted of human DNA cloned into bacterial artificial chromosomes (BAC arrays) representing the human genome at approximately 1Mb intervals. The present arrays consist of 25-base pair oligonucleotides (probes). In the last decade the resolution of the arrays have improved to the extent that the number of smaller CNVs identified in the genome that were previously invisible to microscopy has increased enormously. Deletions and duplications at least as small as 10kb are now known to occur throughout the genome.

Interestingly, some CNVs appear to differ from single nucleotide polymorphisms (SNPs) in terms of locus specific mutation rates. Rates for genomic re-arrangements range from  $10^{-4}$  to

$10^{-5}$ , a rate that is considerably more frequent than point mutations [76]. This high mutation rate, coupled with reduced fecundity in some carriers, and the fact that comparatively, they affect a larger proportion of the genome, make CNVs a potentially important source of new and recent mutation in neurodevelopmental disorders.

## 6. CNVs in autism

As discussed above, autism is a phenotypic feature of many genomic disorders. Betancur [77], in her review, lists 103 disease genes and 44 genomic loci where autism or autistic like behaviors have been described. ASD is diagnosed in ~30% of males with Fragile X syndrome and in reverse, Fragile X mutations are found in as many as 7–8% of individuals with ASD [78]. Similarly, mutations in *MECP2*, the Rett Syndrome gene, have been found among cases of autism that do not have the classical Rett phenotype and autism patients have an increased risk for neurofibromatosis and other rare monogenic diseases like tuberous sclerosis and Joubert's Syndrome, again inversely patients with these disorders have an increased risk for having autism [79, 80].

The genes and loci listed in the Betancur study are all causally implicated in learning disability (LD), indicating that these two neurodevelopmental disorders share some genetic risk factors. Early use of aCGH in non-syndromic autism suggested the method had promise in detecting hitherto unrecognized CNVs. For example Jacquemont et al [81] identified 6 deletions and 2 duplications in 29 patients presenting with syndromic ASD where previous high resolution karyotyping was reported as normal. Another study showing the potential for CNV analysis was a large linkage study [14] using a 10k SNP array where intensity data was used to determine copy number. The authors highlight some individual findings including a family with two sisters with ASD, both of whom had a ~300kb deletion on Ch. 2p16 that included the coding exons of the *neurexin 1* gene (*NRXN1*). A second finding was a recurrent 1.1Mb duplication at Ch. 1q2.1 in four affected individuals from three families, a third was a ~900kb de novo duplication at 17p12 in an affected sib-pair, and with the same region appearing as a maternally inherited deletion in two male siblings, with a paternally inherited deletion in a further female. Duplications in this region cause Charcot-Marie-Tooth 1A (*CMT1A*) and hereditary neuropathy with pressure palsies when deleted, and overlapping deletions are seen in Smith-Magenis syndrome that includes autism symptoms in many cases [82].

A key development was the report of de novo copy number variants in autism [83] using aCGH. These authors showed that individually rare CNVs, and in particular ones that affect neurodevelopmental genes, were enriched in cases. They further suggested that the rate of de novo CNVs differed between simplex cases, where they occurred in 10% of families in the sample, and familial cases where they occurred in 3% of families, suggesting that sporadic and familial cases of ASD might have different underlying genetic mechanisms, although not all studies since then have found this distinction. Several studies of CNVs in large autism case and family series have followed. Marshall et al. [84] examined 427 ASD families using a 500k

SNP array and karyotyping by standard clinical diagnostic method. A de novo rate of 7.1% and 2.0% in simplex and multiplex families respectively was observed, supporting the previous findings [83]. Families occasionally showed more than a single de-novo event where both may combine to produce risk. A further set of loci were identified in two or more unrelated families, increasing the evidence supporting a pathogenic role. As with the LD literature, at some loci, both deletions and duplication were found suggesting a more complex mechanism than simple over or under-expression of gene products. Of the 196 inherited CNVs confirmed experimentally, 90 were of maternal and 106 of paternal origin. The authors list numerous potential ASD candidate genes where a structural change was either de-novo, found in two or more unrelated ASD cases, or, in the case of the X-chromosome, transmitted from an unaffected mother. Given their rarity, very few individual CNVs in this study provided statistical evidence to support their role in autism. For example, 4 CNVs from 427 cases were found at the *DPP6* – a subunit that affects the function of Kv4.2 channels at the same site of expression as *SHANK3* and *NLGN* gene products. Only one similar CNV was found in 1652 controls (Fischer's exact test  $p = 0.016$ ). In keeping with previous cytogenetic findings and the emerging overlap in disorders involved, CNVs were found in ASD cases that involved known loci or genes in disorders such as Waardenburg Type IIa, Speech and language disorder, learning disability and VCFS. A further study [85] found a total of 51 CNVs in 46 cases and not in the controls. 42 of these were familial and 9 de novo with recurrence in two or more cases at three loci. In total, case specific CNVs were found in 11.6% of cases, although, in keeping with the Marshall et al [84] study, none were individually associated with case status with the majority being observed in only a single case. Pinto et al [86] compared CNVs in 996 ASD cases of European ancestry to 1,287 matched controls, using the Illumina 1M SNP array. Cases were found to carry a higher global burden of rare, genic CNVs, especially so for loci previously implicated in either ASD and/or intellectual disability. Nearly 6% of the cases had de-novo mutations with some having two or more events. Novel candidate genes were identified that were de-novo in cases and not controls, including *SHANK2*, *SYNGAP1* and *DLGAP2*. In keeping with previous studies only one novel CNV (maternally inherited X-linked deletions at *PTCHD1*) occurred statistically more frequently in cases compared to controls (7 vs 0). *PTCHD1* involvement in autism and LD was further extended and confirmed in a focused examination of the *PTCHD1* locus in cohorts of autism and LD cases and extending the study of CNVs to sequence data, identifying additional maternally inherited missense mutations in 8 probands not seen in controls [87]. In the Pinto et al study [86], certain gene sets were found to be enriched for case deletions but not duplications. These included sets involved in cell and neuronal development, projection, motility and proliferation; GTPase/Ras signaling known to be involved in regulating dendrite and spine plasticity; and kinase activity/regulation. There was additional overlap with gene sets thought to be involved in LD including microtubule cytoskeleton, glycosylation and CNS development/adhesion. More recently, Salyakina and colleagues [88] have shown the value of extended multiply affected families in a CNV study of 42 families. They found 5 deletions and 7 duplications that co-segregated with ASD, two overlapping with known autism CNVs on 7p21.3 and 15q24.1 and two near regions on 3p26.3 and 12q24.32 previously associated with schizophrenia.

As the resolution of the probe arrays improves, smaller CNVs will be detected, and the boundaries of previously identified CNVs will become more refined [89]. Nord et al. [90], examined genomic DNA of 41 children with autism and 367 healthy controls for rare CNVs using a very high-resolution aCGH platform. They found that cases were more likely than controls to have CNVs as small as ~10 kb, likely to affect genes involved in transcription, nervous system development, and receptor activity. They found that expression of *CNTNAP2*, *ZNF214*, *PRODH* and *ARID1B* genes affected by CNVs were decreased in probands compared with controls suggesting reduced expression as a potentially aetiological factor during development.

Larger samples, particularly those based on families, will also enable the improved estimation of the overall effects of de-novo mutations and the assessment of rare recurrent events as disease associated mutations [91]. These authors studied 1124 autism families containing probands, unaffected parents and an unaffected sibling using the Illumina 1M SNP array. In a related paper [92] they were able to confirm de novo CNVs identified using the SNP array with those detected using a Nimblegen 2.1M aCGH platform. A combined total of 58 rare de novo CNVs were identified across the two studies with each array type identifying 95% of the total. However, the sensitivity for smaller CNVs was low for both arrays. Overall, the burden of rare de novo CNVs in the Sanders et al [91] study was greater in probands than in siblings for total number, size and gene content. Using the rate in siblings as a control to evaluate findings in the cases, there was strong individual statistical support for recurrent de novo duplications at 7q11.23, the locus at which deletions cause Williams-Beuren syndrome; deletions at 16p11.2 and duplications at 16p11.2. In addition the authors observed 8 loci at which rare transmitted CNVs, present only in probands, overlapped with one of the 51 regions in probands containing one or more rare *de novo* CNVs. However, the rare transmitted CNVs were not more likely to be in cases than in unaffected siblings, even when subdivided into genic, exonic, brain-expressed or previously identified as ASD related. This suggested that the excess burden in their sample was due to rare de novo events, although when the gene sets were applied to gene pathway analysis, more pathways showed enrichment in the case set compared to the sibling set. To date, the number of definitive replicated findings for ASD from all studies has been small, with the data suggesting an extreme heterogeneity model with no single risk variant occurring in more than 1% of cases.

## 7. Mutation rates and models of risk in autism

Given the replicated finding that de-novo mutations are more frequent in simplex cases compared to familial cases, Zhao et al [93] have suggested a model of autism risk in which families fall into two groups; those in which the overall risk for autism is low, representing the majority of families, and those in which the risk is higher due to a disease mutation with a dominant mode of transmission with greater penetrance in males compared to females. Under this model, sporadic cases of autism occur in low-risk families due to a de novo mutation of relatively high penetrance, whereas familial autism occurs due to the inheritance of an existing mutation from a clinically unaffected or asymptomatic parent. In another model, Girirajan et al. [94] proposed the necessity in some families for a second mutation to lead to severe neurodeve-

developmental disorder. In this study, individuals with childhood developmental delay are enriched approximately fourfold for a rare 520-kb 16p12 deletion. In nearly all cases examined (22/23), the deletion was inherited. Thus, 16p12 deletions appear to be an example of inherited predisposition to neurodevelopmental disorder with dominant transmission. However, these individuals were more likely to carry a second large (>500 kb) CNV compared to matched controls, and clinical features of those with a second large CNV were typically more severe than those with the 16p12 deletion alone. Itsara et al. [95] suggest that multiply affected autism pedigrees segregate an existing inherited mutation of low penetrance which by itself is rarely sufficient to cause disease. Secondary mutations, such as de-novo mutations are required to manifest as disorder. Whether or not these second hits are disease specific remains to be examined. The authors propose that the excess of *de novo* CNVs among cases may be due to a depletion of second-hits in the unaffected sibling due to the initial low penetrant mutation segregating in the family. The abundance of inherited low penetrance mutations and the high rate of de novo CNVs in the population enable multiple cases to appear within families with apparently unusual patterns of inheritance.

## 8. Next-generation sequencing studies

There is an increasing literature describing the occurrence of single nucleotide mutation in ASD. Unlike the common polymorphisms described earlier, these variants are often rare or private to an individual and considered to have a large impact on the gene's function. Early studies have focused on candidate genes with strong *a priori* evidence to suggest causation, such as the Neuroligin and Shank encoding genes. Sanger sequencing of the exons of families with ASD have identified many putatively causative mutation in these synaptic proteins [96, 97]. As with the transition from candidate gene association studies to GWAS, the advent of more sophisticated high-throughput next-generation sequencing approaches have enabled many genes to be screened in a single experiment. This has extended to the systematic scanning of the whole exome of individuals with ASD to identify putative risk mutations.

In April 2012, three key manuscripts were published in Nature describing a considerable exome sequencing effort of 622 family trios and 250 unaffected siblings [98-100]. In these studies the focus was on de novo mutation in the exome of individuals with ASD. *De novo* mutations are those changes that occurred in the child which are not inherited from the parent. Given the natural mutation rate, the number of families studied and the number of genes under investigation it was estimated that three independent mutations would provide strong evidence implicating a gene in ASD, whilst two independent mutations would be suggestive but not definitive. Unfortunately in these studies no gene was burdened with three independent mutations, whilst several revealed two independent de novo mutations. Follow-up analyses across the studies of the two-hit genes in a cohort of over 2600 individuals with ASD and over 1600 typically developing controls identified additional non-synonymous mutations in *KATNAL2* (katanin p60 subunit A-like 2), *CHD8* (Chromodomain-helicase-DNA-binding protein 8), *GRIN2B* (Glutamate [NMDA] receptor subunit epsilon-2), *LAMC3* (Laminin subunit gamma-3), *SCN1A* (sodium channel, voltage-gated, type I, alpha subunit) and *SCN2A* (sodium

channel, voltage-gated, type II, alpha subunit). Despite the early caution regarding the number of de novo events observed per gene, the burgeoning evidence from focused re-sequencing of these suggests that other families with ASD have damaging loss-of-function mutation in these genes.

## 9. Implications of genomic studies for clinical practice

Genetic findings to date in autism include known genetic syndromes such as Rett and Fragile X; chromosomal structural variation such as duplications on ch15 and deletions on ch22 and increasing numbers of CNVs, many de novo, in a small proportion of cases, especially those with co-morbid developmental abnormalities. Common variants are only slowly emerging from genome-wide studies but sample sizes are still relatively modest compared to other disorders. The diagnosis remains as a clinical one, but genetic testing has diagnostic, prognostic and family planning implications, and in the coming years may also be of importance with respect to potential treatments for specific conditions such as Rett and Fragile X. Therefore, a clinical genetics evaluation should be considered in ASD children in order to identify syndromic forms of autism, identify familial cases, and drive diagnostic testing. Genetic testing using aCGH has been recommended by the American Academy of Pediatrics [101] and the American College of Medical Genetics [102, 103], particularly to evaluate aetiological heterogeneity and identify syndromes. For the vast majority of cases with an identified genetic cause, there is no specific benefit in terms of treatment, but the diagnosis itself may be of value to parents seeking an explanation for the symptoms they see in their child. Acceptance of the disorder may be easier in the presence of a known cause as echoes of social causation and blame can still be heard. Family investigation may also be warranted, particularly where the proband has an identified genetic causation as recurrence risk in siblings may have implications for family planning. If a genetic cause of pathogenic significance is identified, the recurrence risk for sibs may be established according to the particular genetic diagnosis. In the majority of cases, no genetic alteration is found, and an empirical risk for siblings of 10-20% can be provided. Establishing a clear de novo origin for a given mutation may enable a low risk estimate in siblings although there are caveats relating to the exact origin of the mutation. Finally, and re-stating the main goal of genetic research in ASD, a better understanding of underlying biology may lead to novel and unexpected treatment possibilities. Early signs of this are emerging in human trials for treatments aimed at the specific mutations found in Rett and Fragile X syndromes.

### Author details

Michael Gill, Graham Kenny and Richard Anney

Department of Psychiatry, School of Medicine, Trinity College Dublin, Ireland

## References

- [1] Lord C. Epidemiology: How common is autism? *Nature*. 2011 Jun 9;474(7350):166-8. PubMed PMID: 21654793.
- [2] Fombonne E. Epidemiology of pervasive developmental disorders. *Pediatr Res*. 2009 Jun;65(6):591-8. PubMed PMID: 19218885. Epub 2009/02/17. eng.
- [3] Fernell E, Gillberg C. Autism spectrum disorder diagnoses in Stockholm preschoolers. *Res Dev Disabil*. 2010 May-Jun;31(3):680-5. PubMed PMID: 20149593. Epub 2010/02/13. eng.
- [4] Folstein S, Rutter M. Genetic influences and infantile autism. *Nature*. 1977 Feb 24;265(5596):726-8. PubMed PMID: 558516. Epub 1977/02/24. eng.
- [5] Bailey A, Le Couteur A, Gottesman I, Bolton P, Simonoff E, Yuzda E, et al. Autism as a strongly genetic disorder: evidence from a British twin study. *Psychol Med*. 1995 Jan; 25(1):63-77. PubMed PMID: 7792363. Epub 1995/01/01. eng.
- [6] Ronald A, Hoekstra RA. Autism spectrum disorders and autistic traits: a decade of new twin studies. *Am J Med Genet B Neuropsychiatr Genet*. 2011 Apr;156B(3):255-74. PubMed PMID: 21438136. Epub 2011/03/26. eng.
- [7] Hallmayer J, Cleveland S, Torres A, Phillips J, Cohen B, Torigoe T, et al. Genetic Heritability and Shared Environmental Factors Among Twin Pairs With Autism. *Arch Gen Psychiatry*. 2011 Jul 4. PubMed PMID: 21727249. Epub 2011/07/06. Eng.
- [8] Le Couteur A, Rutter M, Lord C, Rios P, Robertson S, Holdgrafer M, et al. Autism diagnostic interview: a standardized investigator-based instrument. *J Autism Dev Disord*. 1989 Sep;19(3):363-87. PubMed PMID: 2793783. Epub 1989/09/01. eng.
- [9] Lord C, Rutter M, Goode S, Heemsbergen J, Jordan H, Mawhood L, et al. Autism diagnostic observation schedule: a standardized observation of communicative and social behavior. *J Autism Dev Disord*. 1989 Jun;19(2):185-212. PubMed PMID: 2745388. Epub 1989/06/01. eng.
- [10] International Molecular Genetic Study of Autism Consortium (IMGSAC). A full genome screen for autism with evidence for linkage to a region on chromosome 7q. International Molecular Genetic Study of Autism Consortium. *Hum Mol Genet*. 1998 Mar;7(3):571-8. PubMed PMID: 9546821. Epub 1998/04/18. eng.
- [11] International Molecular Genetic Study of Autism Consortium (IMGSAC), Bailey A. Further characterization of the autism susceptibility locus AUTS1 on chromosome 7q. *Hum Mol Genet*. 2001 Apr 15;10(9):973-82. PubMed PMID: 11392322. Epub 2001/06/08. eng.
- [12] International Molecular Genetic Study of Autism Consortium (IMGSAC), Monaco AP. A genomewide screen for autism: strong evidence for linkage to chromosomes 2q, 7q,

- and 16p. *Am J Hum Genet.* 2001 Sep;69(3):570-81. PubMed PMID: 11481586. Epub 2001/08/02. eng.
- [13] Barrett S, Beck JC, Bernier R, Bisson E, Braun TA, Casavant TL, et al. An autosomal genomic screen for autism. Collaborative linkage study of autism. *Am J Med Genet.* 1999 Dec 15;88(6):609-15. PubMed PMID: 10581478. Epub 1999/12/03. eng.
- [14] Szatmari P, Paterson AD, Zwaigenbaum L, Roberts W, Brian J, Liu XQ, et al. Mapping autism risk loci using genetic linkage and chromosomal rearrangements. *Nat Genet.* 2007 Mar;39(3):319-28. PubMed PMID: 17322880. Epub 2007/02/27. eng.
- [15] Philippe A, Martinez M, Guilloud-Bataille M, Gillberg C, Rastam M, Sponheim E, et al. Genome-wide scan for autism susceptibility genes. Paris Autism Research International Sibpair Study. *Hum Mol Genet.* 1999 May;8(5):805-12. PubMed PMID: 10196369. Epub 1999/04/10. eng.
- [16] Liu J, Nyholt DR, Magnussen P, Parano E, Pavone P, Geschwind D, et al. A genomewide screen for autism susceptibility loci. *Am J Hum Genet.* 2001 Aug;69(2):327-40. PubMed PMID: 11452361. Epub 2001/07/14. eng.
- [17] Schellenberg GD, Dawson G, Sung YJ, Estes A, Munson J, Rosenthal E, et al. Evidence for multiple loci from a genome scan of autism kindreds. *Mol Psychiatry.* 2006 Nov; 11(11):1049-60, 979. PubMed PMID: 16880825. Epub 2006/08/02. eng.
- [18] Persico AM, D'Agruma L, Maiorano N, Totaro A, Militerni R, Bravaccio C, et al. Reelin gene alleles and haplotypes as a factor predisposing to autistic disorder. *Mol Psychiatry.* 2001 Mar;6(2):150-9. PubMed PMID: 11317216. Epub 2001/04/24. eng.
- [19] Skaar DA, Shao Y, Haines JL, Stenger JE, Jaworski J, Martin ER, et al. Analysis of the RELN gene as a genetic risk factor for autism. *Mol Psychiatry.* 2005 Jun;10(6):563-71. PubMed PMID: 15558079. Epub 2004/11/24. eng.
- [20] Zhang H, Liu X, Zhang C, Mundo E, Macciardi F, Grayson DR, et al. Reelin gene alleles and susceptibility to autism spectrum disorders. *Mol Psychiatry.* 2002;7(9):1012-7. PubMed PMID: 12399956. Epub 2002/10/26. eng.
- [21] Serajee FJ, Zhong H, Mahbulul Huq AH. Association of Reelin gene polymorphisms with autism. *Genomics.* 2006 Jan;87(1):75-83. PubMed PMID: 16311013. Epub 2005/11/29. eng.
- [22] Ashley-Koch AE, Jaworski J, Ma de Q, Mei H, Ritchie MD, Skaar DA, et al. Investigation of potential gene-gene interactions between APOE and RELN contributing to autism risk. *Psychiatr Genet.* 2007 Aug;17(4):221-6. PubMed PMID: 17621165. Epub 2007/07/11. eng.
- [23] Li H, Li Y, Shao J, Li R, Qin Y, Xie C, et al. The association analysis of RELN and GRM8 genes with autistic spectrum disorder in Chinese Han population. *Am J Med Genet B Neuropsychiatr Genet.* 2008 Mar 5;147B(2):194-200. PubMed PMID: 17955477. Epub 2007/10/24. eng.

- [24] Holt R, Barnby G, Maestrini E, Bacchelli E, Brocklebank D, Sousa I, et al. Linkage and candidate gene studies of autism spectrum disorders in European populations. *Eur J Hum Genet.* 2010 Sep;18(9):1013-9. PubMed PMID: 20442744. Pubmed Central PMCID: 2987412. Epub 2010/05/06. eng.
- [25] Kelemenova S, Schmidtova E, Ficek A, Celec P, Kubranska A, Ostatnikova D. Polymorphisms of candidate genes in Slovak autistic patients. *Psychiatr Genet.* 2010 Aug; 20(4):137-9. PubMed PMID: 20436377. Epub 2010/05/04. eng.
- [26] Krebs MO, Betancur C, Leroy S, Bourdel MC, Gillberg C, Leboyer M. Absence of association between a polymorphic GGC repeat in the 5' untranslated region of the reelin gene and autism. *Mol Psychiatry.* 2002;7(7):801-4. PubMed PMID: 12192627. Pubmed Central PMCID: 1913931. Epub 2002/08/23. eng.
- [27] Bonora E, Beyer KS, Lamb JA, Parr JR, Klauck SM, Benner A, et al. Analysis of reelin as a candidate gene for autism. *Mol Psychiatry.* 2003 Oct;8(10):885-92. PubMed PMID: 14515139. Epub 2003/09/30. eng.
- [28] Devlin B, Bennett P, Dawson G, Figlewicz DA, Grigorenko EL, McMahon W, et al. Alleles of a reelin CGG repeat do not convey liability to autism in a sample from the CPEA network. *Am J Med Genet B Neuropsychiatr Genet.* 2004 Apr 1;126B(1):46-50. PubMed PMID: 15048647. Epub 2004/03/30. eng.
- [29] Li J, Nguyen L, Gleason C, Lotspeich L, Spiker D, Risch N, et al. Lack of evidence for an association between WNT2 and RELN polymorphisms and autism. *Am J Med Genet B Neuropsychiatr Genet.* 2004 Apr 1;126B(1):51-7. PubMed PMID: 15048648. Epub 2004/03/30. eng.
- [30] Dutta S, Sinha S, Ghosh S, Chatterjee A, Ahmed S, Usha R. Genetic analysis of reelin gene (RELN) SNPs: no association with autism spectrum disorder in the Indian population. *Neurosci Lett.* 2008 Aug 15;441(1):56-60. PubMed PMID: 18597938. Epub 2008/07/04. eng.
- [31] He Y, Xun G, Xia K, Hu Z, Lv L, Deng Z, et al. No significant association between RELN polymorphism and autism in case-control and family-based association study in Chinese Han population. *Psychiatry Res.* 2011 May 30;187(3):462-4. PubMed PMID: 20554015. Epub 2010/06/18. eng.
- [32] Jackson PB, Boccuto L, Skinner C, Collins JS, Neri G, Gurrieri F, et al. Further evidence that the rs1858830 C variant in the promoter region of the MET gene is associated with autistic disorder. *Autism Res.* 2009 Aug;2(4):232-6. PubMed PMID: 19681062. Epub 2009/08/15. eng.
- [33] Sousa I, Clark TG, Toma C, Kobayashi K, Choma M, Holt R, et al. MET and autism susceptibility: family and case-control studies. *Eur J Hum Genet.* 2009 Jun;17(6):749-58. PubMed PMID: 19002214. Epub 2008/11/13. eng.

- [34] Thanseem I, Nakamura K, Miyachi T, Toyota T, Yamada S, Tsujii M, et al. Further evidence for the role of MET in autism susceptibility. *Neurosci Res.* 2010 Oct;68(2):137-41. PubMed PMID: 20615438. Epub 2010/07/10. eng.
- [35] Benayed R, Gharani N, Rossman I, Mancuso V, Lazar G, Kamdar S, et al. Support for the homeobox transcription factor gene ENGRAILED 2 as an autism spectrum disorder susceptibility locus. *Am J Hum Genet.* 2005 Nov;77(5):851-68. PubMed PMID: 16252243. Epub 2005/10/28. eng.
- [36] Zhong H, Serajee FJ, Nabi R, Huq AH. No association between the EN2 gene and autistic disorder. *J Med Genet.* 2003 Jan;40(1):e4. PubMed PMID: 12525552. Pubmed Central PMCID: 1735256. Epub 2003/01/15. eng.
- [37] Arking DE, Cutler DJ, Brune CW, Teslovich TM, West K, Ikeda M, et al. A common genetic variant in the neurexin superfamily member CNTNAP2 increases familial risk of autism. *Am J Hum Genet.* 2008 Jan;82(1):160-4. PubMed PMID: 18179894. Epub 2008/01/09. eng.
- [38] Li X, Hu Z, He Y, Xiong Z, Long Z, Peng Y, et al. Association analysis of CNTNAP2 polymorphisms with autism in the Chinese Han population. *Psychiatr Genet.* 2010 Jun;20(3):113-7. PubMed PMID: 20414140. Epub 2010/04/24. eng.
- [39] Anney R, Klei L, Pinto D, Almeida J, Bacchelli E, Baird G, et al. Individual common variants exert weak effects on the risk for autism spectrum disorders. *Hum Mol Genet.* 2012 Aug 8. PubMed PMID: 22843504. Epub 2012/07/31. Eng.
- [40] Segurado R, Conroy J, Meally E, Fitzgerald M, Gill M, Gallagher L. Confirmation of association between autism and the mitochondrial aspartate/glutamate carrier SLC25A12 gene on chromosome 2q31. *Am J Psychiatry.* 2005 Nov;162(11):2182-4. PubMed PMID: 16263864. Epub 2005/11/03. eng.
- [41] Kim SJ, Silva RM, Flores CG, Jacob S, Guter S, Valcante G, et al. A quantitative association study of SLC25A12 and restricted repetitive behavior traits in autism spectrum disorders. *Mol Autism.* 2011;2(1):8. PubMed PMID: 21609426. Epub 2011/05/26. eng.
- [42] Turunen JA, Rehnstrom K, Kilpinen H, Kuokkanen M, Kempas E, Ylisaukko-Oja T. Mitochondrial aspartate/glutamate carrier SLC25A12 gene is associated with autism. *Autism Res.* 2008 Jun;1(3):189-92. PubMed PMID: 19360665. Epub 2009/04/11. eng.
- [43] Correia C, Coutinho AM, Diogo L, Grazina M, Marques C, Miguel T, et al. Brief report: High frequency of biochemical markers for mitochondrial dysfunction in autism: no association with the mitochondrial aspartate/glutamate carrier SLC25A12 gene. *J Autism Dev Disord.* 2006 Nov;36(8):1137-40. PubMed PMID: 17151801. Epub 2006/12/08. eng.
- [44] Rabionet R, McCauley JL, Jaworski JM, Ashley-Koch AE, Martin ER, Sutcliffe JS, et al. Lack of association between autism and SLC25A12. *Am J Psychiatry.* 2006 May;163(5):929-31. PubMed PMID: 16648338. Epub 2006/05/02. eng.

- [45] Blasi F, Bacchelli E, Carone S, Toma C, Monaco AP, Bailey AJ, et al. SLC25A12 and CMYA3 gene variants are not associated with autism in the IMGSAC multiplex family sample. *Eur J Hum Genet.* 2006 Jan;14(1):123-6. PubMed PMID: 16205742. Epub 2005/10/06. eng.
- [46] Chien WH, Wu YY, Gau SS, Huang YS, Soong WT, Chiu YN, et al. Association study of the SLC25A12 gene and autism in Han Chinese in Taiwan. *Prog Neuropsychopharmacol Biol Psychiatry.* 2010 Feb 1;34(1):189-92. PubMed PMID: 19913066. Epub 2009/11/17. eng.
- [47] Gallagher L, Becker K, Kearney G, Dunlop A, Stallings R, Green A, et al. Brief report: A case of autism associated with del(2)(q32.1q32.2) or (q32.2q32.3). *J Autism Dev Disord.* 2003 Feb;33(1):105-8. PubMed PMID: 12708586. Epub 2003/04/24. eng.
- [48] Conroy J, Cochrane L, Anney RJ, Sutcliffe JS, Carthy P, Dunlop A, et al. Fine mapping and association studies in a candidate region for autism on chromosome 2q31-q32. *Am J Med Genet B Neuropsychiatr Genet.* 2009 Jun 5;150B(4):535-44. PubMed PMID: 18846500. Epub 2008/10/11. eng.
- [49] Auranen M, Vanhala R, Varilo T, Ayers K, Kempas E, Ylisaukko-Oja T, et al. A genomewide screen for autism-spectrum disorders: evidence for a major susceptibility locus on chromosome 3q25-27. *Am J Hum Genet.* 2002 Oct;71(4):777-90. PubMed PMID: 12192642. Epub 2002/08/23. eng.
- [50] Auranen M, Varilo T, Alen R, Vanhala R, Ayers K, Kempas E, et al. Evidence for allelic association on chromosome 3q25-27 in families with autism spectrum disorders originating from a subisolate of Finland. *Mol Psychiatry.* 2003 Oct;8(10):879-84. PubMed PMID: 14515138. Epub 2003/09/30. eng.
- [51] Ylisaukko-oja T, Nieminen-von Wendt T, Kempas E, Sarenus S, Varilo T, von Wendt L, et al. Genome-wide scan for loci of Asperger syndrome. *Mol Psychiatry.* 2004 Feb; 9(2):161-8. PubMed PMID: 14966474. Epub 2004/02/18. eng.
- [52] Coon H, Matsunami N, Stevens J, Miller J, Pingree C, Camp NJ, et al. Evidence for linkage on chromosome 3q25-27 in a large autism extended pedigree. *Hum Hered.* 2005;60(4):220-6. PubMed PMID: 16391490. Epub 2006/01/05. eng.
- [53] Yonan AL, Alarcon M, Cheng R, Magnusson PK, Spence SJ, Palmer AA, et al. A genomewide screen of 345 families for autism-susceptibility loci. *Am J Hum Genet.* 2003 Oct;73(4):886-97. PubMed PMID: 13680528. Epub 2003/09/19. eng.
- [54] Weiss LA, Arking DE, Daly MJ, Chakravarti A. A genome-wide linkage and association scan reveals novel loci for autism. *Nature.* 2009 Oct 8;461(7265):802-8. PubMed PMID: 19812673. Epub 2009/10/09. eng.
- [55] Risch N, Merikangas K. The future of genetic studies of complex human diseases. *Science.* 1996 Sep 13;273(5281):1516-7. PubMed PMID: 8801636. Epub 1996/09/13. eng.

- [56] Wang K, Zhang H, Ma D, Bucan M, Glessner JT, Abrahams BS, et al. Common genetic variants on 5p14.1 associate with autism spectrum disorders. *Nature*. 2009 May 28;459(7246):528-33. PubMed PMID: 19404256. Epub 2009/05/01. eng.
- [57] Ma D, Salyakina D, Jaworski JM, Konidari I, Whitehead PL, Andersen AN, et al. A genome-wide association study of autism reveals a common novel risk locus at 5p14.1. *Ann Hum Genet*. 2009 May;73(Pt 3):263-73. PubMed PMID: 19456320. Epub 2009/05/22. eng.
- [58] Anney R, Klei L, Pinto D, Regan R, Conroy J, Magalhaes TR, et al. A genome-wide scan for common alleles affecting risk for autism. *Hum Mol Genet*. 2010 Oct 15;19(20):4072-82. PubMed PMID: 20663923. Epub 2010/07/29. eng.
- [59] Xu B, Woodroffe A, Rodriguez-Murillo L, Roos JL, van Rensburg EJ, Abecasis GR, et al. Elucidating the genetic architecture of familial schizophrenia using rare copy number variant and linkage scans. *Proc Natl Acad Sci U S A*. 2009 Sep 29;106(39):16746-51. PubMed PMID: 19805367. Epub 2009/10/07. eng.
- [60] Kirov G, Pocklington AJ, Holmans P, Ivanov D, Ikeda M, Ruderfer D, et al. De novo CNV analysis implicates specific abnormalities of postsynaptic signalling complexes in the pathogenesis of schizophrenia. *Mol Psychiatry*. 2012 Feb;17(2):142-53. PubMed PMID: 22083728. Epub 2011/11/16. eng.
- [61] Lesca G, Rudolf G, Labalme A, Hirsch E, Arzimanoglou A, Genton P, et al. Epileptic encephalopathies of the Landau-Kleffner and continuous spike and waves during slow-wave sleep types: Genomic dissection makes the link with autism. *Epilepsia*. 2012 Sep;53(9):1526-38. PubMed PMID: 22738016. Epub 2012/06/29. Eng.
- [62] Bradley WE, Raelson JV, Dubois DY, Godin E, Fournier H, Prive C, et al. Hotspots of large rare deletions in the human genome. *PLoS ONE*. 2010;5(2):e9401. PubMed PMID: 20195527. Pubmed Central PMCID: 2828468. Epub 2010/03/03. eng.
- [63] Karras GI, Kustatscher G, Buhecha HR, Allen MD, Pugieux C, Sait F, et al. The macro domain is an ADP-ribose binding module. *EMBO J*. 2005 Jun 1;24(11):1911-20. PubMed PMID: 15902274. Epub 2005/05/20. eng.
- [64] Timinszky G, Till S, Hassa PO, Hothorn M, Kustatscher G, Nijmeijer B, et al. A macrodomain-containing histone rearranges chromatin upon sensing PARP1 activation. *Nat Struct Mol Biol*. 2009 Sep;16(9):923-9. PubMed PMID: 19680243. Epub 2009/08/15. eng.
- [65] Hoff KG, Wolberger C. Getting a grip on O-acetyl-ADP-ribose. *Nat Struct Mol Biol*. 2005 Jul;12(7):560-1. PubMed PMID: 15999106. Epub 2005/07/07. eng.
- [66] Liou GG, Tanny JC, Kruger RG, Walz T, Moazed D. Assembly of the SIR complex and its regulation by O-acetyl-ADP-ribose, a product of NAD-dependent histone deacetylation. *Cell*. 2005 May 20;121(4):515-27. PubMed PMID: 15907466. Epub 2005/05/24. eng.
- [67] Chen D, Vollmar M, Rossi MN, Phillips C, Kraehenbuehl R, Slade D, et al. Identification of macrodomain proteins as novel O-acetyl-ADP-ribose deacetylases. *J Biol Chem*. 2011 Apr 15;286(15):13261-71. PubMed PMID: 21257746. Epub 2011/01/25. eng.

- [68] Cohen-Armon M, Visochek L, Katzoff A, Levitan D, Susswein AJ, Klein R, et al. Long-term memory requires polyADP-ribosylation. *Science*. 2004 Jun 18;304(5678):1820-2. PubMed PMID: 15205535. Epub 2004/06/19. eng.
- [69] Qureshi IA, Mehler MF. Emerging roles of non-coding RNAs in brain evolution, development, plasticity and disease. *Nature reviews Neuroscience*. 2012 Aug;13(8):528-41. PubMed PMID: 22814587. Epub 2012/07/21. eng.
- [70] Purcell SM, Wray NR, Stone JL, Visscher PM, O'Donovan MC, Sullivan PF, et al. Common polygenic variation contributes to risk of schizophrenia and bipolar disorder. *Nature*. 2009 Aug 6;460(7256):748-52. PubMed PMID: 19571811. Epub 2009/07/03. eng.
- [71] Reddy KS. Cytogenetic abnormalities and fragile-X syndrome in Autism Spectrum Disorder. *BMC medical genetics*. 2005 Jan 18;6:3. PubMed PMID: 15655077. Pubmed Central PMCID: 548305.
- [72] Gillberg C. Chromosomal disorders and autism. *J Autism Dev Disord*. 1998 Oct;28(5):415-25. PubMed PMID: 9813777. Epub 1998/11/14. eng.
- [73] Vorstman JA, Staal WG, van Daalen E, van Engeland H, Hochstenbach PF, Franke L. Identification of novel autism candidate regions through analysis of reported cytogenetic abnormalities associated with autism. *Mol Psychiatry*. 2006 Jan;11(1):1, 18-28. PubMed PMID: 16205736. Epub 2005/10/06. eng.
- [74] Gothelf D, Schaer M, Eliez S. Genes, brain development and psychiatric phenotypes in velo-cardio-facial syndrome. *Dev Disabil Res Rev*. 2008;14(1):59-68. PubMed PMID: 18636637.
- [75] Zhang F, Gu W, Hurles ME, Lupski JR. Copy number variation in human health, disease, and evolution. *Annual review of genomics and human genetics*. 2009;10:451-81. PubMed PMID: 19715442. Epub 2009/09/01. eng.
- [76] Lupski JR. Structural variation in the human genome. *The New England journal of medicine*. 2007 Mar 15;356(11):1169-71. PubMed PMID: 17360997. Epub 2007/03/16. eng.
- [77] Betancur C. Etiological heterogeneity in autism spectrum disorders: more than 100 genetic and genomic disorders and still counting. *Brain Res*. 2011 Mar 22;1380:42-77. PubMed PMID: 21129364. Epub 2010/12/07. eng.
- [78] Muhle R, Trentacoste SV, Rapin I. The genetics of autism. *Pediatrics*. 2004 May;113(5):e472-86. PubMed PMID: 15121991. Epub 2004/05/04. eng.
- [79] Marui T, Hashimoto O, Nanba E, Kato C, Tochigi M, Umekage T, et al. Association between the neurofibromatosis-1 (NF1) locus and autism in the Japanese population. *American journal of medical genetics Part B, Neuropsychiatric genetics : the official publication of the International Society of Psychiatric Genetics*. 2004 Nov 15;131B(1):43-7. PubMed PMID: 15389774. Epub 2004/09/25. eng.

- [80] Zafeiriou DI, Ververi A, Vargiami E. Childhood autism and associated comorbidities. *Brain & development*. 2007 Jun;29(5):257-72. PubMed PMID: 17084999. Epub 2006/11/07. eng.
- [81] Jacquemont ML, Sanlaville D, Redon R, Raoul O, Cormier-Daire V, Lyonnet S, et al. Array-based comparative genomic hybridisation identifies high frequency of cryptic chromosomal rearrangements in patients with syndromic autism spectrum disorders. *Journal of medical genetics*. 2006 Nov;43(11):843-9. PubMed PMID: 16840569. Pubmed Central PMCID: 2563185. Epub 2006/07/15. eng.
- [82] Smith ACM, Boyd K, Elsea SH, Finucane BM, Haas-Givler B, Gropman A, et al. Smith-Magenis Syndrome. In: Pagon RA, Bird TD, Dolan CR, Stephens K, editors. *GeneReviews*. Seattle (WA)1993.
- [83] Sebat J, Lakshmi B, Malhotra D, Troge J, Lese-Martin C, Walsh T, et al. Strong association of de novo copy number mutations with autism. *Science*. 2007 Apr 20;316(5823):445-9. PubMed PMID: 17363630. eng.
- [84] Marshall CR, Noor A, Vincent JB, Lionel AC, Feuk L, Skaug J, et al. Structural variation of chromosomes in autism spectrum disorder. *Am J Hum Genet*. 2008 Feb;82(2):477-88. PubMed PMID: 18252227. Epub 2008/02/07. eng.
- [85] Christian SL, Brune CW, Sudi J, Kumar RA, Liu S, Karamohamed S, et al. Novel sub-microscopic chromosomal abnormalities detected in autism spectrum disorder. *Biol Psychiatry*. 2008 Jun 15;63(12):1111-7. PubMed PMID: 18374305. Epub 2008/04/01. eng.
- [86] Pinto D, Pagnamenta AT, Klei L, Anney R, Merico D, Regan R, et al. Functional impact of global rare copy number variation in autism spectrum disorders. *Nature*. 2010 Jul 15;466(7304):368-72. PubMed PMID: 20531469. Pubmed Central PMCID: 3021798. Epub 2010/06/10. eng.
- [87] Noor A, Whibley A, Marshall CR, Gianakopoulos PJ, Piton A, Carson AR, et al. Disruption at the PTCHD1 Locus on Xp22.11 in Autism spectrum disorder and intellectual disability. *Sci Transl Med*. 2010 Sep 15;2(49):49ra68. PubMed PMID: 20844286. Pubmed Central PMCID: 2987731. Epub 2010/09/17. eng.
- [88] Salyakina D, Cukier HN, Lee JM, Sacharow S, Nations LD, Ma D, et al. Copy number variants in extended autism spectrum disorder families reveal candidates potentially involved in autism risk. *PloS one*. 2011;6(10):e26049. PubMed PMID: 22016809. Pubmed Central PMCID: 3189231. Epub 2011/10/22. eng.
- [89] Celestino-Soper PB, Shaw CA, Sanders SJ, Li J, Murtha MT, Ercan-Sencicek AG, et al. Use of array CGH to detect exonic copy number variants throughout the genome in autism families detects a novel deletion in TMLHE. *Human molecular genetics*. 2011 Nov 15;20(22):4360-70. PubMed PMID: 21865298. Pubmed Central PMCID: 3196886. Epub 2011/08/26. eng.
- [90] Nord AS, Roeb W, Dickel DE, Walsh T, Kusenda M, O'Connor KL, et al. Reduced transcript expression of genes affected by inherited and de novo CNVs in autism. *European*

- journal of human genetics : EJHG. 2011 Jun;19(6):727-31. PubMed PMID: 21448237. Pubmed Central PMCID: 3110052. Epub 2011/03/31. eng.
- [91] Sanders SJ, Ercan-Sencicek AG, Hus V, Luo R, Murtha MT, Moreno-De-Luca D, et al. Multiple recurrent de novo CNVs, including duplications of the 7q11.23 Williams syndrome region, are strongly associated with autism. *Neuron*. 2011 Jun 9;70(5):863-85. PubMed PMID: 21658581. Epub 2011/06/11. eng.
- [92] Levy D, Ronemus M, Yamrom B, Lee YH, Leotta A, Kendall J, et al. Rare de novo and transmitted copy-number variation in autistic spectrum disorders. *Neuron*. 2011 Jun 9;70(5):886-97. PubMed PMID: 21658582. Epub 2011/06/11. eng.
- [93] Zhao X, Leotta A, Kustanovich V, Lajonchere C, Geschwind DH, Law K, et al. A unified genetic theory for sporadic and inherited autism. *Proceedings of the National Academy of Sciences of the United States of America*. 2007 Jul 31;104(31):12831-6. PubMed PMID: 17652511. Pubmed Central PMCID: 1933261. Epub 2007/07/27. eng.
- [94] Girirajan S, Eichler EE. Phenotypic variability and genetic susceptibility to genomic disorders. *Human molecular genetics*. 2010 Oct 15;19(R2):R176-87. PubMed PMID: 20807775. Pubmed Central PMCID: 2953748. Epub 2010/09/03. eng.
- [95] Itsara A, Wu H, Smith JD, Nickerson DA, Romieu I, London SJ, et al. De novo rates and selection of large copy number variation. *Genome Res*. 2010 Nov;20(11):1469-81. PubMed PMID: 20841430. Pubmed Central PMCID: 2963811. Epub 2010/09/16. eng.
- [96] Jamain S, Quach H, Betancur C, Rastam M, Colineaux C, Gillberg IC, et al. Mutations of the X-linked genes encoding neuroligins NLGN3 and NLGN4 are associated with autism. *Nat Genet*. 2003 May;34(1):27-9. PubMed PMID: 12669065. Epub 2003/04/02. eng.
- [97] Durand CM, Betancur C, Boeckers TM, Bockmann J, Chaste P, Fauchereau F, et al. Mutations in the gene encoding the synaptic scaffolding protein SHANK3 are associated with autism spectrum disorders. *Nat Genet*. 2007 Jan;39(1):25-7. PubMed PMID: 17173049. Epub 2006/12/19. eng.
- [98] O'Roak BJ, Vives L, Girirajan S, Karakoc E, Krumm N, Coe BP, et al. Sporadic autism exomes reveal a highly interconnected protein network of de novo mutations. *Nature*. 2012 Apr 4. PubMed PMID: 22495309. Epub 2012/04/13. Eng.
- [99] Neale BM, Kou Y, Liu L, Ma'ayan A, Samocha KE, Sabo A, et al. Patterns and rates of exonic de novo mutations in autism spectrum disorders. *Nature*. 2012 Apr 4. PubMed PMID: 22495311. Epub 2012/04/13. Eng.
- [100] Sanders SJ, Murtha MT, Gupta AR, Murdoch JD, Raubeson MJ, Willsey AJ, et al. De novo mutations revealed by whole-exome sequencing are strongly associated with autism. *Nature*. 2012 Apr 4. PubMed PMID: 22495306. Epub 2012/04/13. Eng.
- [101] Johnson CP, Myers SM. Identification and evaluation of children with autism spectrum disorders. *Pediatrics*. 2007 Nov;120(5):1183-215. PubMed PMID: 17967920. Epub 2007/10/31. eng.

- [102] Schaefer GB, Mendelsohn NJ. Genetics evaluation for the etiologic diagnosis of autism spectrum disorders. *Genetics in medicine : official journal of the American College of Medical Genetics*. 2008 Jan;10(1):4-12. PubMed PMID: 18197051. Epub 2008/01/17. eng.
- [103] Roesser J. Diagnostic yield of genetic testing in children diagnosed with autism spectrum disorders at a regional referral center. *Clinical pediatrics*. 2011 Sep;50(9):834-43. PubMed PMID: 21525079. Epub 2011/04/29. eng.

IntechOpen

IntechOpen