

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,600

Open access books available

138,000

International authors and editors

170M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Pesticides: Environmental Impacts and Management Strategies

Harsimran Kaur Gill and Harsh Garg

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/57399>

1. Introduction

Increase in food production is the prime-most objective of all countries, as world population is expected to grow to nearly 10 billion by 2050. Based on evidence, world population is increasing by an estimated 97 million per year (Saravi and Shokrzadeh, 2011). The Food and Agricultural Organization (FAO) of the United Nations has in-fact issued a sobering forecast that world food production needs to increase by 70%, in order to keep pace with the demand of growing population. However, increase in food production is faced with the ever-growing challenges especially the new area that can be increased for cultivation purposes is very limited (Saravi and Shokrzadeh, 2011). The increasing world population has therefore put a tremendous amount of pressure on the existing agricultural system so that food needs can be met from the same current resources like land, water etc. In the process of increasing crop production, herbicides, insecticides, fungicides, nematicides, fertilizers and soil amendments are now being used in higher quantities than in the past. These chemicals have mainly come into the picture since the introduction of synthetic insecticides in 1940, when organochlorine (OCl) insecticides were first used for pest management. Before this introduction, most weeds, pests, insects and diseases were controlled using sustainable practices such as cultural, mechanical, and physical control strategies.

Pesticides have now become an integral part of our modern life and are used to protect agricultural land, stored grain, flower gardens as well as to eradicate the pests transmitting dangerous infectious diseases. It has been estimated that globally nearly \$38 billion are spent on pesticides each year (Pan-Germany, 2012). Manufacturers and researchers are designing new formulations of pesticides to meet the global demand. Ideally, the applied pesticides should only be toxic to the target organisms, should be biodegradable and eco-friendly to some extent (Rosell et al., 2008). Unfortunately, this is rarely the case as most of the pesticides are

non-specific and may kill the organisms that are harmless or useful to the ecosystem. In general, it has been estimated that only about 0.1% of the pesticides reach the target organisms and the remaining bulk contaminates the surrounding environment (Carriger et al., 2006). The repeated use of persistent and non-biodegradable pesticides has polluted various components of water, air and soil ecosystem. Pesticides have also entered into the food chain and have bioaccumulated in the higher trophic level. More recently, several human acute and chronic illnesses have been associated with pesticides exposure (Mostafalou and Abdollahi, 2012). Below, we have detailed the effect of pesticides on target and non-target organisms including earthworms, predators, pollinators, humans, fishes, amphibians, and birds. Additionally, impact of pesticides on soil, water and air ecosystems is also discussed. Furthermore, an eco-friendly practice (Integrated Pest Management (IPM) approach) has been detailed as a strategy that could minimize the use of pesticides.

2. Effects of pesticides on target organisms

Over the past era there has been an increase in the development of pesticides to target a broad spectrum of pests. The increased quantity and frequency of pesticide applications have posed a major challenge to the targeted pests causing them to either disperse to new environment and/or adapt to the novel conditions (Meyers and Bull, 2002; Cothran et al., 2013). The adaptation of the pest to the new environment could be attributed to the several mechanisms such as gene mutation, change in population growth rates, and increase in number of generations etc. This has ultimately resulted in increased incidence of pest resurgence and appearance of pest species that are resistant to pesticides.

2.1. Pesticide resistance

“Resistance may be defined as a heritable change in the sensitivity of a pest population that is reflected in the repeated failure of a product to achieve the expected level of control when used according to the label recommendation for that pest species” (IRAC, 2013). Resistant individuals tend to be rare in a normal population, but indiscriminate use of chemicals can eliminate normal susceptible populations and thereby providing the resistant individuals a selective advantage in the presence of a pesticide. Resistant individuals continue to multiply in the absence of competition and eventually become the dominant portion of the population over generations. As majority of the individuals of a population are resistant, the insecticide is no longer effective thus causing the appearance or development of insecticide resistance.

Resistance is the most serious bottleneck in the successful use of pesticides these days. The intensive use of pesticides has led to the development of resistance in many targeted pest species around the globe (Tabashnik et al., 2009). Number of resistant insects and mite species had risen to 600 by the end of 1990, and increased to over 700 by the end of 2001. This trend is likely to be continued in 21st century as well. Resistance has been found in different insecticides groups e.g., 291 species have developed cyclodiene resistance, followed by DDT (263 species), organophosphates (260 species), carbamates (85 species), pyreth-

roids (48 species), fumigants (12 species), and other (40 species) (Dhaliwal et al., 2006). Important crop pests, parasites of livestock, common urban pests and disease vectors in some cases have developed resistance to such an extent that their control has become exceedingly challenging (Van Leeuwen et al., 2010; Gondhalekar et al., 2011). However, many factors such as genetics, biology/ecology and control operations influence the development of pesticide resistance (Georghiou and Taylor, 1977).

Insecticide bioassays using whole insects continue to be one of the most widely used approaches for detecting resistance (Brown and Brogdon, 1987; Gondhalekar et al., 2013) despite some associated drawbacks. In the past two decades, however, several new methods employing advanced biochemical and molecular techniques, and combination of insecticide bioassays have been developed for detecting insecticide resistance (Symondson and Hemingway, 1997; Scharf et al., 1999; Zhou et al., 2002). Some examples of these techniques are enzyme electrophoresis, enzyme assays, immuno-assays, allele-specific polymerase chain reaction (PCR) etc.

2.2. Pest resurgence

Pest resurgence is defined as the rapid reappearance of a pest population in injurious numbers following pesticide application. Use of persistent and broad spectrum pesticides that kills the beneficial natural enemies is thought to be the leading cause of pest resurgence. However, resurgence is known to occur due to several reasons, for example, increase in feeding and reproductive rates of insect pests, due to application of sub-lethal doses of pesticides, and sometimes elimination of a primary pest provides favorable conditions for the secondary pests to become primary/key pests (Dhaliwal et al., 2006). There are many pesticide-induced pest outbreaks reported in walnut (*Juglans regia*) (Bartlett and Ewart, 1951), hemlock (*Conium maculatum*) (McClure, 1977), soybeans (*Glycine max*) (Shepard et al., 1977), and cotton (*Gossypium hirsutum*) (Bottrell and Rummel, 1978). Among these, brown plant hopper (BPH) (*Nilaparvata lugens* (Stal)) in rice (*Oryza sativa* L.) cultivation has gained a major importance in Asian countries (Chelliah and Heinrichs, 1984). In general, natural BPH populations were kept under check by natural enemies including mirid bugs, ladybird beetles, spiders and various pathogens. However, pesticides have not only destroyed the natural enemies (Fabellar and Heinrichs, 1986), but have influenced the fecundity of BPH females (Wang et al., 2010) further enhancing their resurgence. Additionally, the resurgence of bed bug, *Cimex lectularius* (Davies et al., 2012) and cotton bollworm *Helicoverpa armigera* (Mironidis et al., 2013) have been reported due to insecticide resistance and indiscriminate use of pesticides.

3. Effects of pesticides on non-target organisms

The effect of pesticides on non-target organisms has been a source of worldwide attention and concern for decades. Adverse effects of applied pesticides on non-target arthropods have been widely reported (Ware, 1980). Unfortunately, natural insect enemies e.g., parasitoids and predators are most susceptible to insecticides and are severely affected (Aveling, 1977; Vickerman, 1988). The destruction of natural enemies can exacerbate pest problems as they

play an important role in regulating pest population levels. Usually, if natural enemies are absent, additional insecticide sprays are required to control the target pest. In some cases, natural enemies that normally keep minor pests under check are also affected and this can result in secondary pest outbreaks. Along with natural enemies, population of soil arthropods is also drastically disturbed because of indiscriminate pesticide application in agricultural systems. Soil invertebrates including nematodes, springtails, mites, micro-arthropods, earthworms, spiders, insects and other small organisms make up the soil food web and enable decomposition of organic compounds such as leaves, manure, plant residues etc. They are essential for the maintenance of soil structure, transformation and mineralization of organic matter. Pesticide effects on above mentioned soil arthropods therefore negatively impact several links in the food web. The following are the examples of non-target organisms that are adversely impacted by pesticides.

3.1. Earthworms

Earthworms represent the greatest proportion of terrestrial invertebrates (>80%) (Yasmin and D'Souza, 2010) and play a significant role in improving soil fertility by decomposing the organic matter into humus. Earthworms also play a major role in improving and maintaining soil structure, by creating channels in soil that enable the process of soil aeration and drainage. However, their diversity, density and biomass are strongly influenced by soil management. They are considered as an important indicator of soil quality in agricultural ecosystems (Paoletti, 1999). Earthworms are affected by various agricultural practices and indiscriminate use of pesticides is one of the leading practices affecting them (Pelosi et al., 2013).

Pesticide applications can cause decline in earthworm populations. For example, carbamate insecticides are very toxic to earthworms and some organophosphates have been shown to reduce earthworm populations (Edwards, 1987). Similarly, a field study conducted in South Africa has also reported that earthworms were influenced detrimentally due to chronic and intermittent exposures to chlorpyrifos and azinphos methyl, respectively (Reinecke and Reinecke, 2007). Various scientific studies reported that pesticides influence earthworm growth, reproduction (cocoon production, number of hatchlings per cocoon, and incubation period) in a dose-dependent manner (Yasmin and D'Souza, 2010). Earthworms exposed to different kind of pesticides showed rupturing of cuticle, oozing out of coelomic fluid, swelling, and paling of body that led to softening of body tissues (Solaimalai et al., 2004). Similarly a study carried out in France showed that the combination of insecticides and fungicides at different concentrations caused neurotoxic effects in earthworms (Schreck et al., 2008). Increased exposure period and higher dose of insecticides can also cause physiological damage (cellular dysfunction and protein catabolism) to earthworms (Schreck et al., 2008).

3.2. Predators

Predators are organisms that live by preying on other organisms and they play a very crucial role in keeping pest populations under control. Predators (beneficial organisms) are also an important part of the "biological control" approach which is one component of the integrated

pest management strategy discussed later. In some of the examples cited below, pesticides were the main cause for decline in predator population:

- In brinjal (*Solanum melongena* L.) ecosystem, spraying with cypermethrin and imidacloprid caused higher mortality of coccinellids, braconid wasps and predatory spiders compared to when sprayed with bio-pesticides and neem (*Azadirachta indica*) based insecticides (Ghananand et al., 2011).
- Species diversity, richness and evenness of collembola, and numbers of spiders were found to be lower in chlorpyrifos treated plots compared with control, in grassland pastures in UK (Fountain et al., 2007).
- Studies were carried out to investigate the effects of chemicals on soil arthropods in agricultural area near Everglades National Park, USA. It was found that higher number of arthropods (including predators such as coccinellids and spiders) were present in non-sprayed fields compared to fields sprayed with insecticides and herbicides (Amlin et al., 2009).
- In foliar application, all the systemic neonicotinoids such as imidacloprid, clothianidin, admire, thiamethoxam and acetamiprid were found highly toxic to natural enemies in comparison with spirotetramat, buprofezin and fipronil (Kumar et al., 2012).

Additionally, pesticides can also affect predator behavior and their life-history parameters including growth rate, development time and other reproductive functions. For example, in the eastern USA, glyphosate-based herbicides affected behavior and survival of spiders and ground beetles, apart from affecting arthropod community dynamics that can also influence biological control in an agroecosystem (Evans et al., 2010). Similarly, dimethoate was shown to significantly decrease the body size, haemocyte counts and reduction of morphometric parameters on carabid beetle (*Pterostichus melas italicus*), in Calabria, Italy (Giglio et al., 2011).

3.3. Pollinators

Pollinators are biotic agents that play a very important role in pollination process. Some of the recognized pollinators are different species of bees, bumble bees (*Bombus* spp.), honey bees (*Apis* spp.), fruit flies, some beetles, and birds (e.g., hummingbirds, honeyeaters, and sunbirds etc.). Pollinators can be used as bioindicators of ecosystemic processes (process by which physical, chemical, biological events help connecting organisms with their environment) in many ways as their activities are affected by environmental stress caused by parasites, competitors, diseases, predators, pesticides and habitat modifications (Kevan, 1999). However, using pesticides causes direct loss of insect pollinators and indirect loss to crops because of the lack of adequate populations of pollinators (Fishel, 2011).

Pesticide application also affects various activities of pollinators including foraging behaviour, colony mortality and pollen collecting efficiency. Most of our current knowledge about effects of pesticides on change in pollinator behaviour has come from various bee studies as they comprise 80% of the insect pollinator population. For instance, many laboratory studies have demonstrated the lethal and sub-lethal effects of neonicotinoid insecticides (imidacloprid,

acetamiprid, clothianidin, thiamethoxam, thiacloprid, dinotefuran and nitenpyram) on foraging behavior, learning and memory abilities of bees (Blacque`re et al., 2012). Worker bee (female bees that lack full reproductive capacity and play many other roles in bee colony) mortality, decreased pollen collecting efficiency and eventually colony collapse occur due to pesticides (neonicotinoid and pyrethroid) application (Gill et al., 2012). In addition to this, non-lethal exposure of honey bees to neonicotinoid insecticide (thiamethoxam) causes high mortality due to homing failure at a level that could put a risk of colony collapse (Henry et al., 2012). Sub-lethal doses of imidacloprid (the most commonly used pesticide worldwide) affected longevity and foraging in honey bees (*A. mellifera*). *Nosema ceranae* (*Nosema* invades the intestinal tracts of adult bees causing colony collapse disorder (CCD) and nosema disease/nosemosis, which consequently lead to decrease in honey production). Microsporidial infections increased significantly in gut of bees from imidacloprid treated hives. It has been anticipated that interactions between pathogens and imidacloprid pesticide could be a main reason for worldwide honey bee colony mortality, including CCD (Pettis et al., 2012; Wu et al., 2012). There are also reports that imidacloprid reduced brood production due to decline in the fecundity of bumble bees (*B. terrestris*) (Laycock et al., 2012; Whitehorn et al., 2012). On the other hand, little work has been done on the impact of pesticides on wild pollinators. For example, a field study carried out in Italy on an agricultural field found lower bumblebee and butterfly species richness associated with pesticide application. They also found that bees (insect pollinators) were at higher risk from pesticide use (Brittain et al., 2010).

3.4. Humans

The deleterious effects of pesticides on human health have started to grow due to their toxicity and persistence in environment and ability to enter into the food chain. Pesticides can enter the human body by direct contact with chemicals, through food especially fruits and vegetables, contaminated water or polluted air. Both acute and chronic diseases can result from pesticide exposure and these are summarized below:

3.4.1. Acute illness

Acute illness generally appears a short time after contact or exposure to the pesticide. Pesticide drift from agricultural fields, exposure to pesticides during application and intentional or unintentional poisoning generally leads to the acute illness in humans (Dawson et al., 2010; Lee et al., 2011b). Several symptoms such as headaches, body aches, skin rashes, poor concentration, nausea, dizziness, impaired vision, cramps, panic attacks and in severe cases coma and death could occur due to pesticide poisoning (Pan-Germany, 2012). The severity of these risks is normally associated with toxicity and quantity of the agents used, mode of action, mode of application, length and frequency of contact with pesticides and person that is exposed during application (Richter, 2002). About 3 million cases are reported worldwide every year that occur due to acute pesticides poisoning. Out of these 3 million pesticide poisoning cases, 2 million are suicide attempts and the rest of these are occupational or accidental poisoning cases (Singh and Mandal, 2013). Suicide attempts due to acute pesticide poisoning are mainly the result of widespread availability of pesticides in rural areas (Richter, 2002; Dawson et al., 2010). Several

strategies have been proposed to reduce the incidences that occur due to acute pesticide poisoning such as restricting the availability of pesticides, substituting the pesticide with a less toxic but with an equally effective alternative and by promoting use of personal protection equipment (Murray and Taylor, 2000; Konradsen et al., 2003). Strict laws regulating pesticide sales along with preventive health programs and community development efforts are needed to enforce such strategies.

3.4.2. Chronic illness

Continued exposure to sub-lethal quantities of pesticides for a prolonged period of time (years to decades), results in chronic illness in humans (Pan-Germany, 2012). Symptoms are not immediately apparent and manifest at a later stage. Agricultural workers are at a higher risk to get affected, however general population is also affected especially due to contaminated food and water or pesticides drift from the fields (Pan-Germany, 2012). Incidences of chronic diseases have started to grow as pesticides have become an increasing part of our ecosystem. There is mounting evidence that establish a link between pesticides exposure and the incidences of human chronic diseases affecting nervous, reproductive, renal, cardiovascular, and respiratory systems (Mostafalou and Abdollahi, 2012). The list of chronic diseases that are linked to prolonged pesticide exposure by various studies is summarized in Table 1.

Diseases	References
Cancer (Childhood and adult brain cancer; Renal cell cancer; lymphocytic leukaemia (CLL); Prostate Cancer)	Lee et al., 2005; Shim et al., 2009; Heck et al., 2010; Xu et al., 2010; Band et al., 2011; Cocco et al., 2013
Neuro degenerative diseases including Parkinson disease, Alzheimer disease	Elbaz et al., 2009; Hayden et al., 2010; Tanner et al., 2011
Cardio-vascular disease including artery disease	Abdullah et al., 2011; Andersen et al., 2012
Diabetes (Type 2 Diabetes)	Son et al., 2010; Lee et al., 2011a
Reproductive disorders	Petrelli and Mantovani, 2002; Greenlee et al., 2003
Birth defects	Winchester et al., 2009; Mesnage et al., 2010
Hormonal imbalances including infertility and breast pain	Xavier et al., 2004
Respiratory diseases (Asthma, Chronic obstructive pulmonary disease (COPD))	Chakraborty et al., 2009; Hoppin et al., 2009

Table 1. The List of chronic diseases that are linked to the exposure to pesticides

Several mechanisms have been illustrated that link development of chronic diseases with pesticide exposure. Direct interaction of pesticides with genetic material resulting in DNA damages and chromosomal aberration is considered to be one of the primary mechanisms that lead to the chronic diseases such as cancer etc (Mostafalou and Abdollahi, 2012). In this context, several studies report an increase in frequency of chromosomal aberration, sister chromatid exchange, and breakage in DNA strand in pesticide applicators who worked in agricultural

fields (Grover et al., 2003; Santovito et al., 2012). Similar to this, pesticides are also known to induce epigenetic changes (heritable changes without any alteration in DNA sequences) through DNA methylation, histone modifications and expression of non-coding RNAs. For example, neurotoxic pesticide paraquat has been implicated to induce the Parkinson's disease (PD) through epigenetic changes by promoting histone acetylation (Song et al., 2010). Pesticides may also induce oxidative stress by increasing reactive oxygen species (ROS) through altering levels of antioxidant enzymes such as superoxide dismutase, glutathione reductase and catalase (Agrawal and Sharma, 2010). Several health problems such as Parkinson disease, disruption of glucose homeostasis have been linked with pesticides induced oxidative stress (Mostafalou and Abdollahi, 2012).

4. Pesticides and soil environment

A major fraction of the pesticides that are used for agriculture and other purposes accumulates in the soil. The indiscriminate and repeated use of pesticides further aggravates this soil accumulation problem. Several factors such as soil properties and soil micro-flora determine the fate of applied pesticides, owing to which it undergoes a variety of degradation, transport, and adsorption/desorption processes (Weber et al., 2004; Laabs et al., 2007; Hussain et al., 2009). The degraded pesticides interact with the soil and with its indigenous microorganisms, thus altering its microbial diversity, biochemical reactions and enzymatic activity (Hussain et al., 2009; Munoz-Leoz et al., 2011). A summary of the effects of pesticides on its various components are given below:

1. Pesticides that reach the soil can alter the soil microbial diversity and microbial biomass. Any alteration in the activities of soil microorganisms due to applied pesticides eventually leads to the disturbance in soil ecosystem and loss of soil fertility (Handa et al., 1999). Numerous studies have been undertaken which highlight these adverse impacts of pesticides on soil microorganisms and soil respiration (Dutta et al., 2010; Sofo et al., 2012). In addition to this, exogenous applications of pesticides could also influence the function of beneficial root-colonizing microbes such as bacteria and arbuscular mycorrhiza (AM), fungi and algae in soil by influencing their growth, colonization and metabolic activities etc (Debenest et al., 2010; Menendez et al., 2010; Tien and Chen, 2012).

The pesticides that reach the soil can interact with soil microflora in several ways:

- a. It can adversely affect the growth, microbial diversity or microbial biomass of the soil microflora. For example, sulfonyleurea herbicides- metsulfuron methyl, chlorsulfuron and thifensulfuron methyl were reported to reduce the growth of the fluorescent bacteria *Pseudomonas* strains that were isolated from an agricultural soil (Boldt and Jacobson, 1998). The *Pseudomonas* spp. is known to play an important ecological role in the soil habitat (Boldt and Jacobson, 1998), and hence its reduction can adversely affect soil fertility. Similarly, benomyl, captan and chlorothalonil were reported to suppress the peak soil respiration (an indicator of microbial biomass) in an unamended soil by 30–50% (Chen et al., 2001b).

al., 1997; Romero et al., 2010). In fact, it has generally been assumed that measuring the change in enzyme activity is an earlier indicator of soil degradation as compared to the chemical or physical parameters (Dick et al., 1994). Several studies have already been undertaken which indicate both increase and decrease in activities of soil enzymes such as hydrolases, oxidoreductases, and dehydrogenase (Ismail et al., 1998; Megharaj et al., 1999). A description of pesticides interactions with soil enzymes has been summarized in Table 2.

Enzyme (Function in soil)	Examples of the pesticides applied	Comments
Nitrogenase (An enzyme used by organisms to fix atmospheric nitrogen gas).	Carbendazim, Imazetapir, Thiram, Captan, 2,4-D, Quinalphos, Monocrotophos, Endosulfan, γ -HCH, Butachlors	Pesticide reduced or inhibited the nitrogenase activity in laboratory or field conditions (Chalam et al., 1996; Martinez-Toledo et al., 1998; Niewiadomska, 2004; Niewiadomska and Klama, 2005; Prasad et al., 2011)/ Pesticides stimulated the nitrogenase activity (Patnaik et al., 1995)
Phosphatase (hydrolyzes organic P compounds to inorganic P)	2,4-D, Nitrapyrin, Monocrotophos, Chlorpyrifos, Mancozeb and Carbendazim	Inhibited (Tu, 1981); Activity increased, but higher concentration or increasing incubation period has inhibitory effects (Madhuri and Rangaswamy, 2002; Srinivasulu et al., 2012b)
Urease (catalyzes the hydrolysis of urea into CO ₂ and NH ₃ and is a key component in the nitrogen cycle in soils)	Isoproturon, Benomyl, Captan, Diazinon, Profenofos	Increase in urease activity (Chen et al., 2001a; Nowak et al., 2004), Pesticide reduced/inhibited urease activity (Abdel-Mallek et al., 1994; Ingram et al., 2005)
Dehydrogenase (DHA): (an oxidoreductase enzyme that catalyzes the removal of hydrogen)	Azadirachtin, Acetamiprid, Quinalphos, Glyphosate	Positive/stimulatory influence on the DHA (Singh and Kumar, 2008; Kizilkaya et al., 2012)/Initially inhibited but later on activity was restored (Andrea et al., 2000; Mayanglambam et al., 2005)
Invertase (hydrolyzes sucrose to fructose and glucose)	Atrazine, Carbaryl, Paraquat	Inhibited invertase activity (Gianfreda et al., 1995; Sannino and Gianfreda, 2001)
β -glucosidase (hydrolyzes disaccharides in soil to form β -glucose)	Metalaxyl, Ridomil gold plus copper	Enzyme activity increased and then decreased (Sukul, 2006) or inhibited (Demanou et al., 2004)
Cellulase (hydrolyzes cellulose to D-glucose)	Benlate, Captan, Brominal	Inhibited enzyme activity (Arinze and Yubedee, 2000; Omar and Abdel-Sater, 2001)

Enzyme (Function in soil)	Examples of the pesticides applied	Comments
Arylsulphatase (an enzyme that hydrolyzes aryl sulfates)	Cinosulfuron, Prosulfuron, Thifensulfuron methyl, Triasulfuron	Decreased enzyme activity (Sofo et al., 2012)

Table 2. A summary of the effects of pesticides on different soil enzymes

Several environmental factors control the bioavailability, degradation and effect of pesticides on soil microorganisms in addition to the persistence, concentration and toxicity of the applied pesticides. These include soil texture, presence of organic matter, vegetation and cultural practices (Murage et al., 2007). For instance, a mixture of compost and straw was found to have the capability of bio-degrading different mixtures of fungicides that are usually applied in vineyards when tested under laboratory conditions (Coppola et al., 2011). Similarly, persistence of the herbicide imazapyr was reported to be different in three Argentinean soils (Tandil, Anguil, and Cerro Azul sites) and its half-life was negatively associated with soil pH, iron and aluminum content, and positively related with clay content (Gianelli et al., 2013). Additionally, level of soil moisture is also one of the most important factors that regulates pesticide bioavailability and degradation, as water acts as solvent for pesticide movement and diffusion, and is essential for microbial functioning (Pal and Tah, 2012). For example, degradation of herbicide saflufenacil was found to be faster at field capacity for Nada, Crowley and Gilbert soils as compared to the saturated soil conditions (Camargo et al., 2013).

It is important to monitor the response of soil microbial communities and various enzymatic activities to pesticide exposure in order to reduce their deleterious effects. A combination of both cultivation-dependent (e.g., community-level physiological profiling (CLPP), measuring overall rates of microbial activity) and cultivation-independent (e.g., DNA sequence information, proteomics of environmental samples) methods can be applied to measure and interpret the effects of pesticide exposure (Imfeld and Vuilleumier, 2012). With the advent of efficient new sequencing techniques and metagenomics, the scope of deploying cultivation independent methods for measuring bacterial diversity and function in soil ecosystem has been further increased. Metagenomics approach has been applied already to measure microbial diversity for a range of soil systems including contaminated sites (Ono et al., 2007) and land managed with different cultural practices (Souza et al., 2013). Such high-tech approaches hold the key for future methods to measure the mode of adaptation ecosystem to different pesticides and in development of new methods to better manage pesticide applications.

A careful screening of pesticide effects on soil microflora should be done in laboratory before their field applications. This is because pesticides tend to accumulate in soil due to repeated applications over time and can pose adverse effects on soil microflora even though they are applied at recommended doses (Ahemad et al., 2009). For instance, Ahemad and Khan (2011) reported the highest toxicity to plant growth promoting characteristics of the *Bradyrhizobium* sp. when its strain MRM6 was grown with three times the recommended field rates of glyphosate, imidacloprid and hexaconazole. Similarly, Dunfield et al. (2000) assessed the

effects of the fungicides captan and thiram at rates of 0.25-2 g a.i. kg⁻¹ on the survival and phenotypic characteristics of bacteria *Rhizobium leguminosarum* bv. *viceae*, strain C1. They found that even though both captan and thiram significantly reduced the numbers of rhizobia recovered from seed and altered the FAME (fatty acid methyl ester) and biological profiles of recovered rhizobia, it was only the highest concentrations of captan that affected nodulation and plant growth. Similarly, herbicide mesotrione affected soil microbial communities, but the effects were only detected at doses far exceeding the recommended field rates (Crouzet et al., 2010). Overall, it is crucial to comprehend the role of pesticides in perturbing soil environment, so that the risk of pesticide contamination and its consequent adverse impacts on soil environment can be evaluated.

5. Pesticides in water and air ecosystem

Pesticide residues in water are a major concern as they pose a serious threat to biological communities including humans. There are different ways by which pesticides can get into water such as accidental spillage, industrial effluent, surface run off and transport from pesticide treated soils, washing of spray equipments after spray operation, drift into ponds, lakes, streams and river water, aerial spray to control water-inhibiting pests (Carter and Heather, 1995; Singh and Mandal, 2013). Pesticides generally move from fields to various water reservoirs by runoff or in drainage induced by rain or irrigation (Larson et al., 2010). Similarly, the presence of pesticides in air can be caused by number of factors including spray drift, volatilization from the treated surfaces, and aerial application of pesticides. Extent of drift depends on: droplet size and wind speed. The rate of volatilization is dependent on time after pesticide treatment, the surface on which the pesticide settles, the ambient temperature, humidity and wind speed and the vapor pressure of the ingredients (Kips, 1985). The volatility or semi-volatility nature of the pesticide compounds similarly constitutes an important risk of atmospheric pollution of large cities (Trajkovska et al., 2009). For instance, organophosphorus (OP) pesticides were identified from environmental samples of air and surface following agricultural spray applications in California and Washington (USA) (Armstrong et al., 2013). In Italian forests, indiscriminate use of pesticides and its active metabolites has led to the contamination of water bodies and ambient air, possibly affecting the health of aquatic biota fishes, amphibians and birds (Trevisan et al., 1993). The following section describes the effect of pesticides on fishes, amphibians and birds.

5.1. Fishes

Fishes are an important part of marine ecosystem as they interact closely with physical, biological and chemical environment. Fishes provide food source for other animals such as sea birds and marine mammals and thus fishes form an integral part of the marine food web. A lot of research has been carried out to examine the impact of pesticides on decline in fish population (Scholz et al., 2012). Pesticides have been directly linked to causing fish mortality worldwide. For example, 27 freshwater fish species are found to be affected by “plant protection products” (PPP) in Europe (Ibrahim et al., 2013). Another pesticide pentachlorophenol

(NaPCP) is reported to cause large numbers of fish mortality in the rice fields of Surinam (Vermeer et al., 1970). Pesticides not only impact the fish but also food webs related to them. The persistent pesticides (organochlorine pesticides and polychlorinated biphenyls) have already been found in the major Arctic Ocean food webs (Hargrave et al., 1992). A survey was conducted to examine the influence of pesticides on aquatic community in West Bengal, India. Many body tissues of the fish such as gills, alimentary canal, liver and brain of carp and catfish were found drastically damaged by pesticides. It was reported that such level of pesticides in fish could harm the fish consumers as well (Konar, 2011).

Several examples are available where pesticides impacted the vital fish organs and behavior. Organophosphate pesticide "Abate" has the potential to alter the vitellogenesis (the process whereby yolky eggs are produced) of catfish (*Heteropneustes fossilis* (Bloch.)), which can severely affect catfish farming (Kumari, 2012). Another major effect of toxic contaminants is on olfaction in fishes since it can affect activities such as mating, locating food, avoiding predators, discriminating kin and homing etc (Tierney et al., 2010). Simultaneous exposure of trematode parasite (*Telogaster opisthorchis*), freshwater fish (*Galaxias anomalus*) and snails to high glyphosate concentrations significantly reduced their survival and development. Within 24 hrs of exposure to higher glyphosate concentrations, 100% mortality of individuals was found (Kelly et al., 2010).

The impact of pesticides within an aquatic environment is influenced by their water solubility and uptake ability within an organism (Pereira et al., 2013). For example, Clomazone, a popular herbicide, is particularly water soluble; a property that increases its likelihood of contaminating surface and groundwater. The hydrophilic (water-loving) or lipophobic (fat-hating) nature of this pesticide makes it less available in the fatty tissues of an organism (Pereira et al., 2013). Further to this, the toxicity of chemical (e.g., endosulfan in this case) in juvenile rainbow trout (*Oncorhynchus mykiss*) was affected by alkalinity, temperature of water and size of the fish (Capkin et al., 2006).

Pesticides in natural water within the acceptable concentration range can still pose harmful effects. Kock-Schulmeyer et al. (2012) found that even if the pesticide levels found in Llobregat River basin of Spain were within the European Union Environmental Quality Standards, they still accounted for a low to high ecotoxicological risk for aquatic organisms, especially algae and macro-invertebrates. Proper measures should be taken while disposing of expired pesticides, so that their discharge into the water bodies does not danger the aquatic life. This is because the alteration in water pH by expired insecticides can lead to acute toxicity of different fish (Satyavani et al., 2011).

5.2. Amphibians

Amphibians are ectothermic, tetrapod vertebrates of class Amphibia. They inhabit a wide variety of habitats, with most species living within fossorial, arboreal, terrestrial, and freshwater aquatic ecosystems. The global decline in the amphibian population has become an environmental concern worldwide. Many amphibian species are on the brink of extinction with 7.4% listed as critically endangered, and at least 43.2% experiencing some sort of population decrease (Stuart et al., 2004). There could be multitude of reasons for

decline in amphibian species diversity, but pesticides appear to be playing an important role. Global warming and climate change are leading to more variable and warmer temperatures which may have increased the impact of pesticides on amphibian populations (Relyea, 2003; Johnson et al., 2013).

Many studies showed that amphibians are susceptible to environmental contaminants due to their permeable skin, dual aquatic-terrestrial cycle and relatively rudimentary immune system (Kerby et al., 2010). Several studies showing the impact of pesticides on amphibians are being mentioned here. It has been reported that the world's most commonly used herbicide (Roundup (Glyphosate)) may have far reaching effects on non-target amphibians (Relyea, 2012). Roundup, a globally used herbicide caused high mortality of larval tadpoles (3 different species in North America) and juvenile frogs under natural conditions in an outdoor pond mesocosm (Relyea, 2005a). Most of the evidence supported the toxic effects of pesticides on juvenile European common frogs (*Rana temporaria*) in an agricultural field that was over sprayed. Mortality of frogs ranged from 100% after 1 hour to 40% after 7 days at the recommended concentrations of pesticides (Bruhl et al., 2013). It was found that population of the wood frog (*Lithobates sylvaticus*) near an agricultural area was more resistant to common insecticide (chlorpyrifos), but not to the common herbicide (Roundup). However, no evidence was reported that resistance carried a performance cost when facing competition and the fear of predation (Cothran et al., 2013).

Further to this, pesticides indirectly affect amphibian populations by influencing growth of aquatic communities such as fungi, zooplankton, and phytoplankton as they are one of their prime energy resources. Malathion is the most commonly used broad-spectrum insecticide in United States. It is legal to spray malathion over aquatic habitats to control mosquitoes (Family: Culicidae), that vector malaria and West Nile Virus. A study found that even low concentration of malathion caused direct and indirect effects on aquatic communities (Relyea, 2012). For example, indirect effect of malathion led to decrease in zooplankton diversity, that led to increase in phytoplankton, a decrease in periphyton, and finally decrease in growth of frog tadpoles (Relyea and Hoverman, 2008). Moreover, it was found that repeated applications of low doses had largest impacts than single high dose application of malathion on an aquatic system (Relyea and Diecks, 2008). A comprehensive study was conducted to examine the effect of globally used pesticides including insecticides (carbaryl, malathion, and herbicides (glyphosate, 2, 4-D)) on aquatic communities (algae, 25 animal species). Species richness reduced differentially, 15% with carbaryl, 30% with malathion, and 22% with roundup, whereas 0% with 2, 4-D. It was found that Roundup completely eliminated two species of tadpoles and led to 70% decline in tadpole species (Relyea, 2005b). Another study demonstrated that frogs (*Rana pipiens*) living in agricultural area, where they experienced higher exposure to chemicals were smaller in size and weight than frogs living in area exposed to low-levels of chemicals. It suggests that frogs living in agricultural areas might have more vulnerability to infections and diseases due to their smaller size and alternation in their immune system (decrease in number of splenocytes and phagocytic activity) (Christin et al., 2013).

5.3. Birds

Birds are a diverse group, and apart from their distinct songs and calls, showy displays and bright colors adding enjoyment to lives of humans, they play a very critical role in food chains and webs in our ecosystems. Birds are also called “aerial acrobats” consuming different kinds of insects such as mosquitoes, European corn borer moth (*Ostrinia nubilalis*), Japanese beetles (*Popillia japonica*), and many other insect species that are considered as some of the most serious agricultural and health pests. Birds are important biotic components of an ecosystem and help in maintaining a natural equilibrium of insect populations by preying on them. In absence of birds, outbreaks of insect pest populations would become more common, ultimately leading to increased pesticide use. Pesticides exposure by different means such as direct ingestion of pesticide granules and treated seeds, treated crops, direct exposure to sprays, contaminated water, or feeding on contaminated prey, and baits cause birds mortality (Fishel, 2011; Guerrero et al., 2012). In USA, almost 50 pesticides are known for killing song birds, game birds, seabirds, shorebirds, and raptors (BLI, 2004).

Pesticides have a potential to alter behavior and reproduction of birds. Some of the examples cited here, using different synthetic chemicals including carbamates, organochlorines, and organophosphates can cause a decline in the populations of raptorial birds by altering their feeding behavior and reproduction (Mitra et al., 2011). A large area in the world is under rice and therefore cultivation and volume of pesticides applied in rice field is quite significant. Many different kinds of organochlorines, cholinesterase-inhibiting insecticides including carbofuran, monocrotophos, phorate, diazinon, fenthion, phosphamidon, methyl parathion and azinphos-methyl along with fungicides, herbicides and molluscicides are being used in rice fields. Some of these chemicals are highly toxic to birds causing mortality and some chemicals even have the potential to affect their reproductive systems (Parsons et al., 2010). Indirect effects of pesticides, through food chain have been proposed as a possible factor in decline of farmland bird species. Insecticides applied in breeding season can affect breeding performance of corn bunting (*Miliaria calandra*) and yellowhammer (*Emberiza citrinella*) (Boatman et al., 2004).

Pesticides, especially insecticides such as carbamates and organophosphates have the potential to cause bird mortality due to their high toxicity (Hunter, 1995). Further to this, insecticides and fungicides pose a most prominent threat to ground-nesting farmland birds as compared to other agricultural practices. The decline of US grassland birds is attributed to acute pesticide toxicity and not agricultural intensification as previously thought (Mineau and Whiteside, 2013). An estimate suggests that 672 million birds are directly exposed to pesticides every year on farmlands, and 10% of these birds die due to acute toxic effects of pesticides (Williams, 1997). A study was conducted in rice fields of Surinam to examine the effects of pesticides, pentachlorophenol (NaPCP) on birds. NaPCP was sprayed for the purpose of killing *Pomacea* snails. Large numbers of dead sick/dead egrets, herons and jacana birds were found during the period of pesticide application. Pentachlorophenol and endrin levels in these birds suggested that ingestion of contaminated food was the probable cause of sickness and mortality (Vermeer et al., 1970).

6. Pesticides and biomagnification

The increase in concentration of pesticides due to its persistent and non-biodegradable nature in the tissues of organisms at each successive level of food chain is known as biomagnification. Due to this phenomenon, organisms at the higher levels of food chain experience greater harm as compared to those at lower levels. Several studies have been undertaken that demonstrate enhanced amount of toxic compounds with increase in trophic levels. For example, out of 36 species collected from three lakes of northeastern Louisiana (USA) that were found to contain residues of 13 organochlorines, tertiary consumers such as green-backed heron (*Butorides striatus*), and snakes etc., contained the highest residues as compared to secondary consumers (bluegill (*Lepomis macrochirus*), blacktail shiner (*Notomis venustus*)) (Niethammer et al., 1984). Similarly, significantly higher concentrations of dichlorodiphenyltrichloroethane (4,4'-DDE) were found in the top consumer fish in Lake Ziway, catfish (*Clarias gariepinus*) than in lower consumers, Nile tilapia (*Oreochromis niloticus*), tilapia (*Tilapia zillii*) and goldfish (*Carassius auratus*) (Deribe et al., 2013). Some of the adverse effects of pesticides on non-target organisms such as fish, amphibians and humans discussed in the above section have also occurred as a result of biomagnifications of the toxic compounds. For example, reproductive failure and population decline in the fish-eating birds (e.g., gulls, terns, herons etc.) was observed as a result of DDE induced eggshell thinning (Grasman et al., 1998). The extent of biomagnifications increases with increase in persistence and lipophilic (fat-loving) characteristics of the particular pesticide. As a result of this, organochlorines are known to have higher biomagnification rate and are more persistent in a wider range of organisms as compared to organophosphates (Favari et al., 2002). It is important to do the risk assessments associated with the pesticides on the basis of their bioaccumulation and biomagnifications before considering them for agricultural purposes.

7. Strategies for pesticide management

There are a relatively few pesticide resistance management tactics that have been proposed risk-free and have a reasonable chance of success under a variety of different circumstances. Headmost among these are: monitoring of pest population in field before any pesticide application, alteration of pesticides with different modes of action, restricting number of applications over time and space, creating or exploiting refugia, avoiding unnecessary persistence, targeting pesticide applications against the most vulnerable stages of pest life cycle, using synergists which can enhance the toxicity of given pesticides by inhibiting the detoxification mechanisms. The most difficult challenge in managing resistance is not the unavailability of appropriate methods but ensuring their adoption by growers and pest control operators (Denholm et al., 1998; Dhaliwal et al., 2006).

Pest resurgence is a dose-dependent process and there are ways to tackle this problem using correct dosage of effective and recommended pesticides. Resurgence problem occurs due to a number of reasons. One of them is due to farmers' tendency to apply low-dose insecticides

due to economic constraints that lead to inadequate and ineffective control of pests. Pest resurgence also occurs due to reduced biological control (most common with insects), reduced competition (most common with weeds; monocots vs. dicots), direct stimulation of pest (due to sub-lethal dose), and improved crop growth.

In the current scenario, optimized use of pesticides is important to reduce environmental contamination while increasing their effectiveness against target pest. This way we can reduce pesticide resistance as well as pest resurgence problems. This has led to the consideration of rational use of pesticides, and the physiological and ecological selectivity of pesticides. Physiological selectivity is characterized by differential toxicity between taxa for a given insecticide. However, ecological selectivity refers to the modification of operational procedure in order to reduce unnecessary destruction to non-target organisms (Dent, 2000). Farmers should focus to use insecticides that are more toxic to target species than their natural enemies which could help to reduce resurgence to some extent (Dhaliwal et al., 2006).

One should consider adopting an Integrated Pest Management (IPM) approach for controlling pests, as these practices are designed to have minimal environment disturbance. The aim of IPM is not only to reduce indiscriminate pesticide use but also to substitute hazardous chemicals with safe chemistries. IPM is a process of achieving long-term, environmentally safe pest control using wide variety of technology and other potential pest management practices. According to National Academy of Science, "IPM refers to an ecological approach in pest management in which all available necessary techniques are consolidated in a unified program so that populations can be managed in such a manner that economic damage is avoided and adverse side effects are minimized" (NAS, 1969). In European arable systems, applied multi-disciplinary research and farmer incentives to encourage the adoption of innovative IPM strategies are essential for development of sustainable maize-based cropping systems. These IPM strategies can contribute immensely to address the European strategic commitment to the environmentally sustainable use of pesticides (Vasileiadis et al., 2011). The added cost and time to do an IPM approach is sometimes a difficult task for growers, but government and extension services can help in convincing and encouraging growers to go for IPM strategy for eco-friendly and long term pest control. We have already discussed earlier that continuous use of pesticides leads to pesticide resistance and pest resurgence problem. To avoid these issues we can always go for other potential management options that include cultural and physical control, host plant resistance, biocontrol, and the use of biopesticides etc.

7.1. Cultural control

Historically, cultural control methods were the farmer's most important tool of preventing crop losses. Cultural control for pest management has been adopted by growers throughout the world for a long time due to its environmentally friendly nature and minimal costs (Gill et al., 2013). Cultural control practices are regular farm operations, which are used to destroy the pests or to prevent them from causing plant damage. Several methods of cultural control have been practiced, such as crop rotation, sanitation, soil solarization, timed planting and harvest, use of resistant varieties, certified seeds, allelopathy, intercropping or "companion planting", use of farmyard manure, and living and organic mulches (Altieri et al., 1978; Dent,

2000; Dhaliwal et al., 2006). Soil solarization (McSorley and Gill, 2010; Gill and McSorley, 2011b) and organic mulches (Gill and McSorley, 2011a) alone and their integration (Gill and McSorley, 2010) were reported as economical and eco-friendly technique for controlling soil-surface arthropods (various insects, and nematodes) (Gill et al., 2010; Gill et al., 2011) and weeds (Gill et al., 2009; Gill and McSorley, 2011b). More effective cultural control can be achieved by synchronizing existing practices with life cycles of pests. This way the weakest link in their life cycle is subjected to adverse climatic conditions.

Large insect populations are killed automatically by farmers when they expose them to adverse climatic conditions through agricultural practices like weeding, ploughing, and hoeing. Ploughing of agricultural field allows turnover of the upper layer of soil while burying the weeds and residues from last year. For example, in South Africa, about 70% of overwintering populations of spotted stalk borer (*Chilo partellus*) and maize stalk borer (*Busseola fusca*) in grain sorghum (*Sorghum bicolor* L.) and maize (*Zea mays* L.) fields were destroyed by slashing the plants. Ploughing and discing of plant residues after slashing further destroyed 24% population on grain sorghum and 19% on maize (Kfir, 1990). Planting dates (Goyal and Kanta, 2005a), and barrier crops (teosinte (*Zea* spp.) and pearl millet (*Pennisetum glaucum* (L.)) (Goyal and Kanta, 2005b) were found to be effective against maize stem borer (*Chilo partellus*) in India. The brown seaweeds *Spatoglossum asperum* and *Sargassum swartzii* can be used as manure to protect plants (tomato (*Solanum lycopersicum* L.) in this case) from root rotting fungi, (*Macrophomina phaseolina*, *Rhizoctonia solani* and *Fusarium solani*) and root-knot nematode (*Meloidogyne javanica*) and for providing necessary nutrients to plants (Sultana et al., 2012). In India, rodents are pests in agriculture, horticulture, forestry, animal husbandry as well as in human dwellings and rural and urban storage facilities. Cultural methods, such as clean cultivation, proper soil tillage and crop scheduling, barriers, repellents and proofing that reduce the rodent harbourage, food sources and immigration may have long lasting effects (Parshad, 1999).

7.2. Physical and mechanical control

Managing pest populations using devices which affect them physically or alter their physical environment is called physical control. Exposure to sun rays, steaming, moisture management especially for stored grain pests, and light traps for attracting various kinds of moths, beetles and other pests are different methods used in physical control. For example steaming woolen winter clothes help in eliminating population of the woolly bear moth, *Antherenus vorax* (waterhouse) (Dhaliwal et al., 2006). Hot water treatment of plant storage products like corns, and bulbs helps to kill many concealed pests such as eelworms and bulb flies. Superheating of empty grain storage godowns to a temperature of 50°C for 10-12 hours helps killing hibernating stored grain pests. Exposure of cotton seeds to sun's heat in thin layers for 2-3 days during summer helps in killing diapausing larvae of pink bollworm (*Pectinophora gossypiella* Saunders) (Dhaliwal et al., 2006).

Mechanical control refers to suppression of pest population by manual devices. It includes various practices such as hand picking, trapping and suction devices, clipping, pruning and crushing of infested shoots and floral parts, and exclusion by screens and barriers to keep away house flies (*Musca domestica*), mosquitoes and other pests. In south-eastern Australia, the

common starling (*Sturnus vulgaris*) is an established invasive avian pest that is now making incursions into Western Australia which is currently free of this species. Trapping with live-lure birds is suggested to be the most cost-effective and widely implemented starling control technique (Campbell et al., 2012). Numerous wildlife species such as coyotes (*Canis latrans* Say), squirrels (Sciuridae family), and birds are known pests of California agriculture in the United States. For these pests, different non-lethal control options including habitat modification, exclusionary devices, and baiting are generally preferred (Baldwin et al., 2013). Mechanical weed control is mainly associated with tillage practices which are performed with special tools such as harrows, hoes, and brushes in growing crops. Increased knowledge about side effects of herbicides has further driven the interest in adoption of mechanical weed control thus increasing the prevalence of organic farming (Rueda-Ayala et al., 2010; Jat et al., 2011). Trapping using yellow colored sticky traps is an effective way for controlling tephritid flies (Dhaliwal et al., 2006).

7.3. Host plant resistance

Host plant resistance (HPR) is the genetic ability of the plant to improve its survival and reproduction by a range of adaptations as compared to the other cultivars when exposed to the same level of pest infestation. HPR offers the most effective, economical and eco-friendly method of pest control (Sharma and Ortiz, 2002), and is considered to be a key element of the IPM strategy. Due to this, identifying and developing HPR has always been a major thrust area of plant breeding, and a number of breeding programs aiming to develop pest resistant crops have been deployed in almost all the cultivated crop species. For example, identification and/or development of resistant varieties in maize against European corn borer (*Ostrinia nubilalis* (Hubner)) (Dhaliwal et al., 2006), brassica against cabbage butterfly (*Pieris brassicae* Linn.) (Chahil and Kular, 2013), wheat (*Triticum aestivum* L.) and rye (*Secale cereale* L.) against *Fusarium* diseases (Miedaner, 1997) *Brassica* sp. against Sclerotinia disease (Garg et al., 2008), and in rice against bacterial blight (Khush et al., 1989). Additionally, availability and access to various germplasm collections have increased the scope of widening the gene-pool of cultivated crops and further identifying and developing HPR. Wild species are especially known to possess a rich repository of genes against various defense traits as they have evolved under different geographic locations. Considerable progress has been made where identification and/or transfer of resistance gene from wild to cultivated species against various pest species has been achieved such as in potato (*Solanum tuberosum* L.) against late blight (*Phytophthora infestans*) from ten wild *Solanum* sp. (Colon and Budding, 1988), wheat against powdery mildew (*Erysiphe graminis*) from wild emmer wheat (*Triticum dicoccoides*) (Reader and Miller, 1991) and mustard (*Brassica juncea*) against *Sclerotinia sclerotiorum* from *Erucastrum cardaminoides* (Garg et al., 2010).

7.3.1. Use of biotechnology and molecular approaches for developing resistant genotype

The advent of new biotechnological and molecular approaches has opened the way to develop resistant genotype that could not only reduce the pesticides application, but it also has a potential to be a part of IPM. Development of resistant genotypes in classical breeding is met

with several challenges such as it is time consuming, desired traits are linked with the undesirable traits (linkage drag) and most importantly lack of resistant genotypes in the gene pool. On the other hand, use of biotechnology in crop improvement ensures the development of pest-resistant genotypes in a comparatively short period of time and minimizes the effects of linkage drag. One of the classic examples where biotechnology was successfully deployed to develop resistant genotype is by the synthesis of transgenic plants which involves modifying plant traits by inserting foreign DNA from a different species (De la Pena et al., 1987). A number of different crops including cotton, rice, mustard, and maize have been modified up to now to engineer the genotypes against various biotic stresses (Ahmad et al., 2012). One of the most successful examples of synthesis of transgenic genotype against pest resistance is in cotton where the gene coding for Bt toxin from the bacterium *Bacillus thuringiensis* (Bt) was inserted leading cotton genotypes to produce Bt toxin in its tissue (Pray et al., 2002; Wu et al., 2008). The lepidopteran larvae that fed on the transgenic plants were killed due to Bt toxin eventually decreasing the amount of pesticide applied to the field. Examples of transgenic crops that have been developed with a potential to reduce pesticides use are abundant and few of them include potato lines against potato tuber moth (*Phthorimaea operculella*) expressing Cry1Ab (Kumar et al., 2010), rice against yellow stem borer (*Scirpophaga incertulas*) expressing potato proteinase inhibitor 2 (Bhutani et al., 2006) and oilseed rape lines resistant to various fungal attack over-expressing tomato chitinase gene (Grison et al., 1996).

Another strategy where biotechnology and molecular approaches have been deployed to combat biotic stresses involves the use of RNA interference (RNAi) technique. This technique primarily uses transgenic plants expressing double stranded RNA (dsRNA) and that reduces the messenger RNA (mRNA) levels (with a high specificity and fidelity) of a crucial gene in the target pest upon feeding (Price and Gatehouse, 2008; Kos et al., 2009). This ultimately interferes with the development and survival of the target pest. RNAi has emerged as a powerful functional genomics approach and it has been used to engineer several crops against number of insect-pests. For example, RNAi technique was used in tobacco genotype that targeted the gene “integrase splicing factor” in root knot nematode, *Meloidogyne incognita* nematode eventually leading to the decrease in the number of nematodes 6-7 weeks post inoculation (Yadav et al., 2006). When such an advanced and effective approach is combined with IPM, it has a great potential to decrease chemical use in agricultural and other ecosystems.

7.4. Biological control

The process of using natural enemies of particular pests to reduce their populations to such a level where economic losses are either eliminated or suppressed is called biological control. Traditionally the most important biocontrol agents are parasitoids, predators and pathogens. Biological control involves three major techniques, *viz.*, introduction, conservation, and augmentation of natural enemies. Biocontrol agents include vertebrates, nemathelminthes (flatworms, and roundworms), arthropods (spiders, mites, and insects), pathogens like viruses, bacteria, protozoa, fungi and rickettsiae all of which play a dynamic role in natural regulation of insect and mite populations (Dhaliwal et al., 2006). In 1762, the Indian Mynah, *Acridotheres tristis* (Linnaeus), was introduced to control red locust in Mauritius. First signifi-

cant success in controlling a pest was achieved on the suggestion of C. V. Riley of California (USA) in 1888. The Vedalia beetle (*Rodolia cardinalis* (Mulsant)), was introduced from Australia into California (USA) for the control of cottony-cushion scale (*Icerya purchasi* maskell) on citrus plants. This scale insect had been accidentally introduced earlier from Australia (Dhaliwal et al., 2006).

Biological control of weeds has been very successful worldwide. There are about 41 species of weeds which have been successfully controlled using insects and pathogens as biocontrol agents. Also, 3 weed species have been controlled using native fungi as mycoherbicides (Mcfadyen, 2000). A total of 12 insects were released in Australia against prickly pear (*Opuntia stricta*), out of these, *Dactylopius opuntiae* and *Cactoblastis cactorum* were responsible for the successful control of prickly pear weed (Julien and Griffiths, 1998). In the past decade, Australia has released 43 species of arthropods and pathogens in 19 different projects for successful biological control of many exotic weeds. Effective biological control was achieved in several projects and outstanding success was achieved in the control of rubber vine (*Cryptostegia grandiflora*), and bridal creeper (*Asparagus asparagoides*) (Palmer et al., 2010).

Examples of biological control are available for other organisms like helminthes, nematodes, fungi, bacteria etc. A nematophagous fungus (*Monacrosporium thaumasium*) was found to be effective in controlling cyathostomin, one of the most important helminthes in tropical region of southeastern Brazil (Tavela et al., 2011). *Trichoderma* species are free-living fungi that have been used to control a broad range of plant pathogenic fungi, viruses, bacteria and nematodes especially root-knot nematodes (*Meloidogyne javanica* and *M. incognita*) (Sharon et al., 2011).

7.4.1. Biorational pesticides

Biorational pesticides/ biopesticides are considered as third-generation pesticides that are rapidly gaining popularity. The word biorational is derived from two words, "biological" and "rational", which means pesticides of natural origin that have limited or no adverse effects on the environment or beneficial organisms. Biopesticides encompass a broad array of microbial pesticides, plant pesticides and biochemical pesticides which are derived from micro-organisms and other natural sources, and processes involving the genetic incorporation of DNA into agricultural commodities. The most commonly used biopesticides include biofungicides (e.g., *Trichoderma* spp.), bioherbicides (*Phytophthora* spp.), bioinsecticides (spore forming bacteria, *Bacillus thuringiensis*, and *B. popilliae*, Actinomycetes), naturally occurring fungi (*Beauveria bassiana*), microscopic roundworms (Entomopathogenic nematodes), Spinosad, insect hormones and insect growth regulators (Gupta and Dikshit, 2010; Singh et al., 2013).

Applications of microbial insecticide, *Chromobacterium subtsugae* for suppression of pecan weevil (*Curculio caryae* (Horn)), and combination of eucalyptus extract and microbial insecticide, *Isaria fumosorosea* (Wize) for control of black pecan aphid (*Melanocallis caryaefoliae* (Davis)) were found promising as alternative insecticides (Shapiro-Ilan et al., 2013). Entomopathogenic nematodes (EPNs) belonging to the families Heterorhabditidae and Steinernematidae are potentially used in South Africa as biocontrol agents against vine mealybug (*Planococcus ficus* (Signoret)) (le Vieux and Malan, 2013). Spinosad was found effective in controlling Colorado potato beetle (*Leptinotarsa decemlineata*) in Iran, and is recommended for use in IPM program for Colorado potato beetle (Soltani and Agricultural, 2011). In China, entomopatho-

genic fungus (*Beauveria bassiana*) has shown great potential for the management of some bark beetle species including red turpentine beetle (RTB) (*Dendroctonus valens* LeConte), a destructive invasive pest (Zhang et al., 2011).

The allelopathic properties of plants can be exploited successfully as a tool for weed and pathogen reduction. In a rice field, application of allelopathic plant material @ 1-2 tonne/ha reduced weed diversity by 70% and increased yield by 20%. Numerous growth inhibitors identified from these allelopathic plants are responsible for their allelopathic properties and may be a useful source for the future development of bio-herbicides and pesticides (Xuan et al., 2005). A combination of coleopteran-active toxin, *Bacillus thuringiensis* Cry3Aa protoxin and protease inhibitors, especially a potato carboxypeptidase inhibitor, have efficiency in preventing damage to stored products and grains by stored grain coleopteran pests (Oppert et al., 2011).

7.5. Chemical control

Sometimes cultural and other agro-technical practices are not sufficient to keep pest population below economic injury level (lowest pest population density that will cause economic crop damage). Therefore, the chemical control agents are resorted to both as preventive and curative measures to minimize the insect pest damage. A good pesticide should be potent against pests, should not endanger the health of humans and non-target organisms, and should ultimately break down into harmless compounds so that it does not persist in environment. Both relative and specific toxicities of the pesticide need to be estimated in order to determine its potency.

It is very important to know spray droplet size and density chemical dosage, application timing, which can provide adequate pest control. There is also a need for research into the development of suitable packaging and disposal procedures, as well as refining of the application equipment. All of these shall rationalize the use of pesticides, so that they can be used in an acceptable way.

Very strict laws should be enacted to protect wildlife and other non-target organisms. Following directions on the pesticide label can prevent injury to non-target organisms. However, when these directions are not followed, benefits from pesticides can be outweighed by the harm and risk associated with pesticides (Fishel, 2011). During pesticide application, things that need to be considered are timing of insecticide application, dosage and persistence, and selective placement of insecticides as discussed below.

7.5.1. Timing of pesticide application

The timing of pesticide application is an important factor to consider before doing any pesticide application. Appropriate application time can ensure not only maximum impact on the target organisms but also least impact on beneficial organisms. Pesticide application timing mainly depends on availability of weather window, time at which pests can be best controlled, and when least damage will be caused to non-target organisms and environment. Flowering period in crops and middle of the day are the times when bees are more prone to insecticides. Hence, insecticide application should not happen at those times to avoid decline in bee populations. Time of insecticide application should coincide with the most vulnerable stage of insect life cycle. Monitoring of insects in the field is thus extremely important for knowing the stage of

insect pest in the field. Monitoring systems are available for most of the insect pests, but spray regime or experiments need to be carried out to determine the most appropriate time for insecticide application for insects for which monitoring systems are not available (Hull and Starner, 1983; Richter and Fuxa, 1984).

Time of the day and season of the year are also important to consider when making pesticide applications. The early morning and evening hours are often the best times for pesticide application because windy conditions are more likely to occur around midday when the temperature warms near the ground level. This causes hot air to rise quickly and mix rapidly with the cooler air above it, favoring drift. During stable conditions, a layer of warm air can stay overhead and not promote mixing with colder air that stays below and closer to the ground. Inversions tend to dissipate during the middle of the day when wind currents mix the air layers. It is very important that applicators recognize thermal inversions and do not spray under those conditions. A temperature or thermal inversion is a condition that occurs naturally and exists when the air at ground level is cooler than the temperature of the air above it. Wind speed is the most important weather factor influencing drift. High wind speeds will move droplets downwind and deposit them off the target. On the other hand, dead calm conditions are never recommended due to likelihood of temperature inversions (Fishel and Ferrell, 2013). Drifting of pesticides increases the possibility of injury to pollinators, humans, domestic animals and wildlife. It is recommended not to spray in wind speed above 2.5 miles/second which otherwise can cause excessive drift and eventually contamination of adjacent areas (Matthews, 1981). Pesticide application should not be made just before rain because pesticides can be washed off by the rain without any impact on the target pest.

7.5.2. Dosage and persistence

Pesticide dose should be sufficient but no greater than the level required for best results. The pesticide manufacturer sets the dose to ensure an acceptable level of control, producing acceptable residue levels, and maximizing returns per unit of formulated insecticide. Persistent pesticides have their benefit of longer persistence on the target and therefore requires less frequent spraying compared to non-persistent pesticides. But care should be taken while using persistent pesticides since these might diminish benefits from natural enemies even at lower doses. If an insecticide is persistent in nature, chances of insecticide residues being harmful to natural enemies are greatly increased (Dent, 2000).

7.5.3. Selective placement

Distribution of pesticides in the field should be such that maximum target cover is achieved. Usually only about 1% of the applied pesticides is able to reach its target, while a large amount of it is wasted. Understanding the pest biology and behavior is critical as it can provide information on pest's habitat, fecundity, feeding etc., which can be important considerations before applying pesticides. Most of the pesticides are applied in liquid form and thus the droplet size is very important in determining their effectiveness. Small droplets provide better coverage and greater likelihood of coming in contact with the target compared to larger droplets that can bounce off the plant surface very easily. The disadvantage with smaller and

bigger droplets is the increased chance of drift and therefore a balance has to be considered between smaller droplets to obtain the maximum effectiveness and reduced drift.

In situations where crops are grown on beds covered with plastic mulch, pesticides should be injected into soil at the time the plastic is laid or injected afterward through drip irrigation system to achieve maximum pesticide effectiveness. For termite (Order: Isoptera) treatments, sometimes perimeter application of insecticides is required around structures/buildings. Additionally, liquids that form foams following injections can be injected into small spaces that are or might be inhabited by termites or other small creatures.

8. Conclusion

Although, pesticides were used initially to benefit human life through increase in agricultural productivity and by controlling infectious disease, their adverse effects have outweighed the benefits associated with their use. The above discussion clearly highlights the severe consequences of indiscriminate pesticide use on different environmental components. Some of the adverse effects associated with pesticide application have emerged in the form of increase in resistant pest population, decline in beneficial organisms such as predators, pollinators and earthworms, change in soil microbial diversity, and contamination of water and air ecosystem. The persistent nature of pesticides has impacted our ecosystem to such an extent that pesticides have entered into various food chains and into the higher trophic levels such as that of humans and other large mammals. Some of the acute and chronic human illnesses have now emerged as a consequence of intake of polluted water, air or food.

This is the time that necessitates the proper use of pesticides to protect our environment and eventually health hazards associated with it. Alternative pest control strategies such as IPM that deploys a combination of different control measures such as cultural control, use of resistant genotype, physical and mechanical control, and rational use of pesticide could reduce the number and amount of pesticide applications. Further, advanced approaches such as biotechnology and nanotechnology could facilitate in developing resistant genotype or pesticides with fewer adverse effects. Community development and various extension programs that could educate and encourage farmers to adopt the innovative IPM strategies hold the key to reduce the deleterious impact of pesticides on our environment.

Acknowledgements

We would like to say special thanks to Drs. Gaurav Goyal (Territory Agronomist, Monsanto), Ameya D. Gondhalekar (Research Assistant Professor, Purdue University), Siddharth Tiwari (Entomologist, BASF), and Matthew R. Tarver (Research Entomologist, USDA) for their valuable suggestions and comments for improving this manuscript.

Author details

Harsimran Kaur Gill^{1*} and Harsh Garg²

*Address all correspondence to: harsimrangill.pau@gmail.com

1 University of Florida, Gainesville, FL, USA

2 The University of Sydney, NSW, Australia

References

- [1] Abdel-Mallek AY, Moharram AM, Abdel-Kader MI, Omar SA. Effect of soil treatment with the organophosphorus insecticide profenofos on the fungal flora and some microbial activities. *Microbiological Research* 1994;149:167-171.
- [2] Abdullah NZ, Ishaka A, Samsuddin N, Mohd Rus R, Mohamed AH. Chronic organophosphate pesticide exposure and coronary artery disease: finding a bridge. *IJUM Research, Invention and Innovation Exhibition (IRIIE)* 2011;223.
- [3] Agrawal A, Sharma B. Pesticides induced oxidative stress in mammalian systems: a review. *International Journal of Biological and Medical Research (IJBMR)* 2010;1:90-104.
- [4] Ahemad M, Khan MS, Zaidi A, Wani PA. Remediation of herbicides contaminated soil using microbes. In: Khan MS, Zaidi A, Musarrat J (eds.) *Microbes in sustainable agriculture*. Nova Science Publishers Inc., New York;2009. p261-284.
- [5] Ahemad M, Khan MS. Effect of pesticides on plant growth promoting traits of green-gram-symbiont, *Bradyrhizobium* sp. strain MRM6. *Bulletin of Environmental Contamination and Toxicology* 2011;86:384-388.
- [6] Ahmad P, Ashraf M, Younis M, Hu X, Kumar A, Akram NA, Al-Qurainy F. Role of transgenic plants in agriculture and biopharming. *Biotechnology Advances* 2012;30:524-540.
- [7] Altieri MA, Francis CA, Schoonhoven AV, Doll JD. A review of insect prevalence in maize (*Zea mays* L.) and bean (*Phaseolus vulgaris* L.) polycultural systems. *Field Crops Research* 1978;1:33-49.
- [8] Amalin DM, Peña JE, Duncan R, Leavengood J, Koptur S. Effects of pesticides on the arthropod community in the agricultural areas near the Everglades National Park. *Proceedings of Florida State Horticultural Society* 2009;122:429-437.
- [9] Andersen HR, Wohlfahrt-Veje C, Dalgård C, Christiansen L, Main KM, Nellemann C, Murata K, Jensen TK, Skakkebaek NE, Grandjean P. Paraoxonase 1 polymorphism

and prenatal pesticide exposure associated with adverse cardiovascular risk profiles at school age. *PloS One* 2012;7:e36830.

- [10] Andrea MM, Peres TB, Luchini LC, Pettinelli Junior A. Impact of long-term pesticide application on some soil biological parameters. *Journal of Environmental Science and Health B* 2000;35:297-307.
- [11] Arinze AE, Yubedee AG. Effect of fungicides on *Fusarium* grain rot and enzyme production in maize (*Zea mays* L.). *Global Journal of Pure and Applied Sciences* 2000;6:629-634.
- [12] Armstrong JL, Fenske RA, Yost MG, Galvin K, Tchong-French M, Yu J. Presence of organophosphorus pesticide oxygen analogs in air samples. *Atmospheric Environment* 2013;66:145-150.
- [13] Aveling C. The biology of Anthocorids (Heteroptera: Anthocoridae) and their role in the integrated control of the damson-hop aphid (*Phorodon humili* Schrank). PhD Thesis, University of London;1977.
- [14] Baldwin RA, Salmon TP, Schmidt RH, Timm RM. Wildlife pests of California agriculture: regional variability and subsequent impacts on management. *Crop Protection* 2013;46:29-37.
- [15] Band PR, Abanto Z, Bert J, Lang B, Fang R, Gallagher RP, Le ND. Prostate cancer risk and exposure to pesticides in British Columbia farmers. *Prostate* 2011;71:168-183.
- [16] Bartlett B, Ewart WH. Effect of parathion on parasites of *Coccus hesperidum*. *Journal of Economic Entomology* 1951;44:344-347.
- [17] Bhagobaty RK, Malik A. Utilization of chlorpyrifos as a sole source of carbon by bacteria isolated from wastewater irrigated agricultural soils in an industrial area of western Uttar Pradesh, India. *Research Journal of Microbiology* 2010;3:293-307.
- [18] Bhutani S, Kumar R, Chauhan R, Singh R, Choudhury VK, Choudhury JB. Development of transgenic indica rice plants containing potato proteinase inhibitor 2 gene with improved defense against yellow stem borer. *Physiological and Molecular Biology of Plants* 2006;12:43-52.
- [19] Blacquiere T, Smagghe G, Cornelis AM, Gestel van, Mommaerts V. Neonicotinoids in bees: a review on concentrations, side-effects and risk assessment. *Ecotoxicology* 2012;21:973-992.
- [20] BLI. State of the world's birds. Birdlife International 2004. (<http://www.biodiversityinfo.org/default.php>).
- [21] Boatman ND, Brickle NW, Hart JD, Milsom TP, Morris AJ, Murray AWA, Murray KA, Robertson PA. Evidence for the indirect effects of pesticides on farmland birds. *Ibis* 2004;146:131-143.

- [22] Boldt TS, Jacobsen CS. Different toxic effects of the sulphonylurea herbicides metsulfuron methyl, chlorsulfuron and thifensulfuron methyl on fluorescent pseudomonads isolated from an agricultural soil. *FEMS Microbiology Letters* 1998;161:29-35.
- [23] Bottrell DG, Rummel DR. Response of *Heliothis* populations to insecticides applied in an area-wide reproduction diapause boll weevil suppression program. *Journal of Economic Entomology* 1978;71:87-92.
- [24] Brittain CA, Vighi M, Bommarco R, Settele J, Potts SG. Impacts of a pesticide on pollinator species richness at different spatial scales. *Basic and Applied Ecology* 2010;11:106-115.
- [25] Brown TM, Brogdon WG. Improved detection of insecticides resistance through conventional and molecular techniques. *Annual Review of Entomology* 1987;32:145-162.
- [26] Bruhl CA, Schmidt T, Pieper S, Alscher A. Terrestrial pesticide exposure of amphibians: an underestimated cause of global decline? *Scientific Reports* 2013;3. (Article number 1135). DOI:10.1038/srep01135.
- [27] Camargo ER, Senseman SA, Haney RL, Guice JB, McCauley GN. Soil residue analysis and degradation of saflufenacil as affected by moisture content and soil characteristics. *Pest Management Science* 2013;69:1291-1297. DOI:10.1002/ps.3494.
- [28] Campbell S, Cook S, Mortimer L, Palmer G, Sinclair R, Woolnough AP. To catch a starling: testing the effectiveness of different trap and lure types. *Wildlife Research* 2012;39:183-191.
- [29] Capkin E, Altinok I, Karahan S. Water quality and fish size affect toxicity of endosulfan, an organochlorine pesticide, to rainbow trout. *Chemosphere* 2006;64:1793-1800.
- [30] Carriger JF, Rand GM, Gardinali PR, Perry WB, Tompkins MS, Fernandez AM. Pesticides of potential ecological concern in sediment from South Florida canals: an ecological risk prioritization for aquatic arthropods. *Soil and Sediment Contamination* 2006;15:21-45.
- [31] Carter AD, Heather AIJ. Pesticides in groundwater: In: Best GA, Ruthven AD. (eds.) *Pesticides- developments, impacts, and controls*. The Royal Society of Chemistry, London, UK;1995. p113-123.
- [32] Chahil GS, Kular JS. Biology of *Pieris brassicae* (Linn.) on different *brassica* species in the plains of Punjab. *Journal of Plant Protection Research* 2013;53:53-59.
- [33] Chakraborty S, Mukherjee S, Roychoudhury S, Siddique S, Lahiri T, Ray MR. Chronic exposures to cholinesterase-inhibiting pesticides adversely affect respiratory health of agricultural workers in India. *Journal of Occupational Health* 2009;51:488-497.

- [34] Chalam AV, Sasikala C, Ramana CV, Rao PR. Effect of pesticides on hydrogen metabolism of *Rhodobacter sphaeroides* and *Rhodospseudomonas palustris*. FEMS Microbiology Ecology 1996;19:1-4.
- [35] Chelliah S, Heinrichs EA. Factors contributing to brown planthopper resurgence. Resurgence of the brown planthopper, *Nilaparvata lugens* (Stål) following insecticide application. Paper presented at the 9th Annual Conference of the Pest Control Council of the Philippines, Philippine International Convention Center, Manila, May 3-6;1984. p36 (mimeo).
- [36] Chen SK, Edwards CA, Subler S. A microcosm approach for evaluating the effects of the fungicides benomyl and captan on soil ecological processes and plant growth. Applied Soil Ecology 2001a;18:69-82.
- [37] Chen SK, Edwards CA, Subler S. Effect of fungicides benomyl, captan and chlorothalonil on soil microbial activity and nitrogen dynamics in laboratory incubations. Soil Biology and Biochemistry 2001b;33:1971-1980.
- [38] Christin MS, Menard L, Giroux I, Marcogliese DJ, Ruby S, Cyr D, Fournier M, Brousseau P. Effects of agricultural pesticides on the health of *Rana pipiens* frogs sampled from the field. Environmental Science and Pollution Research 2013;20:601-611.
- [39] Cocco P, Satta G, Dubois S, Pili C, Pilleri M, Zucca M, Mannetje AM, Becker N, Benavente Y, Sanjose SDe, Foretova L, Staines A, Maynadie M, Nieters A, Brennan P, Miliigi L, Ennas MG, Boffetta P. Lymphoma risk and occupational exposure to pesticides: results of the Epilymph study. Occupational and Environmental Medicine 2013;70:91-98.
- [40] Colon LT, Budding DJ. Resistance to late blight (*Phytophthora infestans*) in ten wild *Solanum* species. Euphytica 1988;39:77-86.
- [41] Coppola L, Comitini F, Casucci C, Milanovic V, Monaci E, Marinozzi M, Taccari M, Ciani M, Vischetti, C. Fungicides degradation in an organic biomixture: impact on microbial diversity. New Biotechnology 2011;29:99-106.
- [42] Cothran RD, Brown JM, Relyea RA. Proximity to agriculture is correlated with pesticide tolerance: evidence for the evolution of amphibian resistance to modern pesticides. Evolutionary Applications 2013;6:832-841.
- [43] Crouzet O, Batisson I, Besse-Hoggan P, Bonnemoy F, Bardot C, Poly F, Bohatier J, Mallet C. Response of soil microbial communities to the herbicide mesotrione: a dose-effect microcosm approach. Soil Biology and Biochemistry 2010;42:193-202.
- [44] Davies TGE, Field LM, Williamson MS. The re-emergence of the bed bug as a nuisance pest: implications of resistance to the pyrethroid insecticides. Medical and Veterinary Entomology 2012;26:241-254.

- [45] Dawson AH, Eddleston M, Senarathna L, Mohamed F, Gawarammana I, Bowe SJ, Manuweera G, Buckley NA. Acute human lethal toxicity of agricultural pesticides: a prospective cohort study. *PLoS Medicine* 2010;7:e1000357.
- [46] De la Pena A, Lörz H, Schell J. Transgenic rye plants obtained by injecting DNA into young floral tillers. *Nature* 1987;325:274-276.
- [47] Debenest T, Silvestre J, Coste M, Pinelli E. Effects of pesticides on freshwater diatoms. In: *Reviews of environmental contamination and toxicology*, Springer, New York 2010;203:87-103.
- [48] Demanou J, Monkiedje A, Njine T, Foto SM, Nola M, Serges H, Togouet Z, Kemka N. Changes in soil chemical properties and microbial activities in response to the fungicide Ridomil gold plus copper. *International Journal of Environmental Research and Public Health* 2004;1:26-34.
- [49] Denholm I, Birnie LC, Kennedy PJ, Shaw KE, Perry JN, Powell W. The complementary roles of laboratory and field testing in ecotoxicological risk assessment. *The 1998 Brighton Conferences: Pest and Diseases Volume 2*. British Crop Protection Council, Farham, UK;1998. p583-590.
- [50] Dent D. Cultural and interference methods. In: *Insect pest management*, 2nd edition. CABI publishing, Cambridge, MA, USA;2000. p235-266.
- [51] Deribe E, Rosseland BO, Borgstrom R, Salbu B, Gebremariam Z, Dadebo E, Skipperud L, Eklo OM. Biomagnification of DDT and its metabolites in four fish species of a tropical lake. *Ecotoxicology and Environmental Safety* 2013;95:10-18.
- [52] Dhaliwal GS, Singh R, Chhillar BS. In: *Essentials of agricultural entomology*. Kalyani Publishers, New Delhi, India;2006.
- [53] Dick RP. Soil enzyme activities as indicators of soil quality. In: Doran JW, Coleman DC, Bezdicek DF, Stewart BA. (eds.) *Defining soil quality for a sustainable environment*, Soil Science Society of America, Madison, WI, USA. 1994;p107-124.
- [54] Dunfield KE, Siciliano SD, Germida JJ. The fungicides thiram and captan affect the phenotypic characteristics of *Rhizobium leguminosarum* strain C1 as determined by FAME and Biolog analyses. *Biology and Fertility of Soils* 2000;31:303-309.
- [55] Dutta M, Sardar D, Pal R, Kole RK. Effect of chlorpyrifos on microbial biomass and activities in tropical clay loam soil. *Environmental Monitoring and Assessment* 2010;160:385-391.
- [56] Edwards CA. The environmental impact of insecticides. In: Delucchi V (ed.) *Integrated pest management, Protection Integree Quo vadis? An International Perspective*. Parasitis 86, Geneva, Switzerland;1987. p309-329.

- [57] Elbaz A, Clavel J, Rathouz PJ, Moisan F, Galanaud JP, Delemotte B, Alperovitch A, Tzourio C. Professional exposure to pesticides and Parkinson disease. *Annals of Neurology* 2009;66:494-504.
- [58] Evans SC, Shaw EM, Rypstra AL. Exposure to a glyphosate-based herbicide affects agrobiont predatory arthropod behavior and long-term survival. *Ecotoxicology* 2010;19:1249-1257.
- [59] Fabellar LT, Heinrichs EA. Relative toxicity of insecticides to rice planthopper and leafhoppers and their predators. *Crop Protection* 1986;5:254-258.
- [60] Favari L, Lopez E, Martinez-Tabche L, Diaz-Pardo E. Effect of insecticides on plankton and fish of Ignacio Ramirez reservoir (Mexico): a biochemical and biomagnification study. *Ecotoxicology and Environmental Safety* 2002;51:177-186.
- [61] Fishel FM, Ferrell JA. Managing pesticide drift. Agronomy department. PI232 University of Florida, Gainesville, FL, USA;2013 (<http://edis.ifas.ufl.edu/pi232>) (accessed 14 October 2013).
- [62] Fishel FM. Pesticides effects on nontarget organisms. PI-85. Pesticide information office, Florida Cooperative Extension Service, IFAS, University of Florida, Gainesville, FL, USA; 2011. (<http://edis.ifas.ufl.edu/pi122>) (accessed 14 October 2013).
- [63] Floch C, Chevremont AC, Joanico K, Capowiez Y, Criquet S. Indicators of pesticide contamination: Soil enzyme compared to functional diversity of bacterial communities via Biolog® Ecoplates. *European Journal of Soil Biology* 2011;47:256-263.
- [64] Fountain MT, Brown VK, Gang AC, Symondson WOC, Murray PJ. The effect of the insecticide chlorpyrifos on spider and collembolan communities. *Pedobiologia* 2007;51:147-158.
- [65] Garcia C, Hernandez T, Costa F. Potential use of dehydrogenase activity as an index of microbial activity in degraded soils. *Communications in Soil Science and Plant Analysis* 1997;28:123-134.
- [66] Garg, H, Atri C, Sandhu PS, Kaur B, Renton M, Banga SK, Singh H, Singh C, Barbetti, MJ, Banga, SS. High level of resistance to *Sclerotinia sclerotiorum* in introgression lines derived from wild crucifers and the crop *Brassica* species *B. napus* and *B. juncea*. *Field Crops Research* 2010;117:51-58.
- [67] Garg H, Sivasithamparam K, Banga SS, Barbetti, MJ. Cotyledon assay as a rapid and reliable method of screening for resistance against *Sclerotinia sclerotiorum* in *Brassica napus* genotypes. *Australasian Plant Pathology* 2008;37:106-111.
- [68] Georghiou GP, Taylor CE. Genetic and biological influences in evolution of insecticide resistance. *Journal of Economic Entomology* 1977;70:319-323.

- [69] Ghananand T, Prasad CS, Lok N. Effect of insecticides, bio-pesticides and botanicals on the population of natural enemies in brinjal ecosystem. *Vegetos- An International Journal of Plant Research* 2011;24:40-44.
- [70] Gianelli VR, Bedmar F, Luis J. Persistence and sorption of imazapyr in three Argentinian soils. *Environmental Toxicology and Chemistry* 2013. DOI:10.1002/etc.2400.
- [71] Gianfreda L, Sannino F, Violante A. Pesticide effects on the activity of free, immobilized and soil invertase. *Soil Biology and Biochemistry* 1995;27:1201-1208.
- [72] Giglio A, Giulianini PG, Zetto T, Talarico F. Effects of the pesticide dimethoate on a non-target generalist carabid, *Pterostichus melas italicus* (Dejean, 1828) (Coleoptera:Carabidae). *Italian Journal of Zoology* 2011;78:471-477.
- [73] Gill HK, McSorley R, Branham M. Effect of organic mulches on soil surface insects and arthropods. *Florida Entomologist* 2011;94:226-232.
- [74] Gill HK, McSorley R, Goyal G, Webb SE. Mulch as a potential management strategy for lesser cornstalk borer, *Elasmopalpus lignosellus* (Insecta: Lepidoptera: Pyralidae), in bush bean (*Phaseolus vulgaris*). *Florida Entomologist* 2010;93:183-190.
- [75] Gill HK, McSorley R, Goyal G. Soil Solarization and organic mulches: tools for pest management. In: Dhawan AK, Singh B, Brar-Bhullar M, Arora R (eds.). *Integrated pest management*. Scientific Publishers, Jodhpur, India;2013. p301-324.
- [76] Gill HK, McSorley R, Treadwell DD. Comparative performance of different plastic films for soil solarization and weed suppression. *HortTechnology* 2009;19:769-774.
- [77] Gill HK, McSorley R. Effect of integrating soil solarization and organic mulching on the soil surface insect community. *Florida Entomologist* 2010;93:308-309.
- [78] Gill HK, McSorley R. Impact of different organic mulches on the soil surface arthropod community and weeds. *International Journal of Pest Management* 2011(a); 58:33-40.
- [79] Gill HK, McSorley R. Effect of different inorganic/synthetic mulches on weed suppression during soil solarization. *Proceeding of Florida State Horticultural Society* 2011(b);124:310-313.
- [80] Gill RJ, Ramos-Rodriguez O, Raine NE. Combined pesticide exposure severely affects individual- and colony-level traits in bees. *Nature* 2012;491:105-108.
- [81] Gondhalekar AD, Scherer CW, Saran RK, Scharf ME. Implementation of an indoxacarb susceptibility monitoring program using field-collected German cockroach isolates from the United States. *Journal of Economic Entomology* 2013;106:945-953.
- [82] Gondhalekar AD, Song C, Scharf, ME. Development of strategies for monitoring indoxacarb and gel bait susceptibility in the German cockroach (Blattodea: Blattellidae). *Pest Management Science* 2011;67:262-270.

- [83] Goyal G, Kanta U. Influence of time and method of sowing of kharif maize on the incidence of *Chilo partellus* (Swinhoe). *Journal of Insect Science* 2005a;19:24-29.
- [84] Goyal G, Kanta U. Influence of barrier crops on the incidence of *Chilo partellus* (Swinhoe) in kharif maize. *Journal of Insect Science* 2005b;19:54-58.
- [85] Gonod LV, Martin-Laurent F, Chenu C. 2, 4-D impact on bacterial communities, and the activity and genetic potential of 2, 4-D degrading communities in soil. *FEMS Microbiology Ecology* 2006;58:529-537.
- [86] Grisman KA, Scanlon PF, Fox GA. Reproductive and physiological effects of environmental contaminants in fish-eating birds of the Great Lakes: a review of historical trends. In: *Trends in levels and effects of persistent toxic substances in the Great Lakes*, Springer, Netherlands;1998. p117-145.
- [87] Greenlee AR, Arbuckle TE, Chyou PH. Risk factors for female infertility in an agricultural region. *Epidemiology* 2003;14:429-436.
- [88] Grison R, Grezes-Besset B, Schneider M, Lucante N, Olsen L, Leguay JJ, Toppan A. Field tolerance to fungal pathogens of *Brassica napus* constitutively expressing a chimeric chitinase gene. *Nature Biotechnology* 1996;14:643-646.
- [89] Grover P, Danadevi K, Mahboob M, Rozati R, Banu BS, Rahman MF. Evaluation of genetic damage in workers employed in pesticide production utilizing the Comet assay. *Mutagenesis* 2003;18:201-205.
- [90] Guerrero I, Morales MB, Oñate JJ, Geige, F, Berendse F, Snoo GD, Tschardt T. Response of ground-nesting farmland birds to agricultural intensification across Europe: Landscape and field level management factors. *Biological Conservation* 2012;152:74-80.
- [91] Gupta S, Dikshit AK. Biopesticides: An eco-friendly approach for pest control. *Journal of Biopesticides* 2010;3:186-188.
- [92] Handa SK, Agnihotri NP, Kulshreshtha G. Effect of pesticide on soil fertility. In: *Pesticide residues; significance, management and analysis* 1999;184-198.
- [93] Hargrave BT, Harding GC, Vass WP, Erickson PE, Fowler BR, Scott V. Organochlorine pesticides and polychlorinated biphenyls in the Arctic Ocean food web. *Archives of Environmental Contamination and Toxicology* 1992;22:41-54.
- [94] Hayden KM, Norton MC, Darcey D, Østbye T, Zandi PP, Breitner JCS, Welsh-Bohmer KA. Occupational exposure to pesticides increases the risk of incident AD The Cache County Study. *Neurology* 2010;74:1524-1530.
- [95] Heck JE, Charbotel B, Moore LE, Karami S, Zaridze DG, Matveev V, Janout V, Kollarova H, Foretova L, Bencko V, Szeszenia-Dabrowska N, Lissowska J, Mates D, Ferro G, Chow W-H, Rothman N, Stewart P, Brennan P, Boffetta P. Occupation and renal

cell cancer in Central and Eastern Europe. *Occupational and Environmental Medicine* 2010;67:47-53.

- [96] Henry M, Bèguin M, Requier F, Rollin O, Odoux J-F, Aupinel P, Aptel J, Tchamitchian S, Decourtye A. A common pesticide decreases foraging success and survival in honey bees. *Science* 2012;336:348-350.
- [97] Hoppin JA, Umbach DM, London SJ, Henneberger PK, Kullman GJ, Coble J, Alavanja MCR, Freeman LEB, Sandler DP. Pesticide use and adult-onset asthma among male farmers in the agricultural health study. *European Respiratory Journal* 2009;34:1296-1303.
- [98] Hull LA, Starner R van. Effectiveness of insecticide applications timed to correspond with the development of rosy apple aphid (Homoptera: Aphididae) on apple. *Journal of Economic Entomology* 1983;76:594-59.
- [99] Hunter K. The poisoning of non-target animals. In: Best GA, Ruthven AD (eds.) *Pesticides- developments, impacts, and controls*. The Royal Society of Chemistry, London, UK;1995. p74-86.
- [100] Hussain S, Siddique T, Saleem M, Arshad M, Khalid A. Impact of pesticides on soil microbial diversity, enzymes, and biochemical reactions. *Advances in Agronomy* 2009;102:159-200.
- [101] Ibrahim L, Preuss TG, Ratte HT, Hommen U. A list of fish species that are potentially exposed to pesticides in edge-of-field water bodies in the European Union-a first step towards identifying vulnerable representatives for risk assessment. *Environmental Science and Pollution Research* 2013;20:2679-2687.
- [102] Imfeld G, Vuilleumier S. Measuring the effects of pesticides on bacterial communities in soil: a critical review. *European Journal of Soil Biology* 2012;49:22-30.
- [103] Ingram CW, Coyne MS, Williams DW. Effects of commercial diazinon and imidacloprid on microbial urease activity in soil. *Journal of Environmental Quality* 2005;34:1573-1580.
- [104] IRAC. Resistance management for sustainable agriculture and improved public health;2013. (<http://www.irac-online.org/>).
- [105] Ismail BS, Yapp KF, Omar O. Effects of metsulfuron-methyl on amylase, urease, and protease activities in two soils. *Australian Journal of Soil Research* 1998;36:449-456.
- [106] Jat RS, Meena HN, Singh AL, Surya JN, Misra JB. Weed management in groundnut (*Arachis hypogaea* L.) in India-a review. *Agricultural Reviews* 2011;32:155-171.
- [107] Johnson LA, Welch B, Whitfield SM. Interactive effects of pesticide mixtures, predators, and environmental regimes on the toxicity of two pesticides to red-eyed tree frog larvae. *Environmental Toxicology* 2013;32:2379-2386.

- [108] Julien MH, Griffiths MW. Biological Control of Weeds: a world catalogue of agents and their target weeds. Wallingford, Oxford: CAB International, 4th ed.;1998. p223.
- [109] Kelly DW, Poulin R, Tompkins DM, Townsend C. Synergistic effects of glyphosate formulations and parasite infection on fish malformations and survival. *Journal of Applied Ecology* 2010;47:498-504.
- [110] Kerby JL, Richards-Hrdlicka KL, Storfer A, Skelly DK. An examination of amphibian sensitivity to environmental contaminants: are amphibians poor canaries? *Ecology Letters* 2010;13:60-67.
- [111] Kevan PG. Pollinators as bioindicators of the state of the environments: species, activity and diversity. *Agriculture, Ecosystems and Environment* 1999;74:373-393.
- [112] Kfir R. Prospects for cultural control of the stalk borers, *Chilo partellus* (Swinhoe) and *Busseola fusca* (Fuller), in summer grain crops in South Africa. *Journal of Entomological Society of Southern Africa* 1990;53:41-47.
- [113] Khush GS, Mackill DJ, Sidhu GS. Breeding rice for resistance to bacterial blight. In: *Bacterial Blight of Rice, Manila, Philippines, International Rice Research Institute* 1989. p207-217.
- [114] Kips RH. Environmental aspects. In: Haskell PT (ed.) *Pesticide application: principles and practice*. Clarendon Press, Oxford;1985. p190-201.
- [115] Kizilkaya R, Akça İ, Aşkın T, Yılmaz R, Olekhov V, Samofalovaf I, Mudrykh N. Effect of soil contamination with azadirachtin on dehydrogenase and catalase activity of soil. *Eurasian Journal of Soil Science* 2012;1:98-103.
- [116] Köck-Schulmeyer M, Ginebreda A, González S, Cortina JL, de Alda ML, Barceló D. Analysis of the occurrence and risk assessment of polar pesticides in the Llobregat River Basin (NE Spain). *Chemosphere* 2012;86:8-16.
- [117] Konar SK. Pesticides and aquatic environment. *Indian Journal of Fisheries* 2011;80-85.
- [118] Konradsen F, Van der HW, Cole DC, Hutchinson G, Daisley H, Singh S, Eddleston M. Reducing acute poisoning in developing countries-options for restricting the availability of pesticides. *Toxicology* 2003;192:249-261.
- [119] Kos M, van Loon JJ, Dicke M, Vet LE. Transgenic plants as vital components of integrated pest management. *Trends in Biotechnology* 2009;27:621-627.
- [120] Kumar M, Chimote V, Singh R, Mishra GP, Naik PS, Pandey Sk, Chakrabarti SK. Development of Bt transgenic potatoes for effective control of potato tuber moth by using cry1Ab gene regulated by GBSS promoter. *Crop Protection* 2010;29:121-127.
- [121] Kumar R, Kranthi S, Nitharwal M, Jat SL, Monga D. Influence of pesticides and application methods on pest and predatory arthropods associated with cotton. *Phytoparasitica* 2012;40:417-424.

- [122] Kumari M. Effects of organophosphate pesticide Abate on the ovary of the cat fish, *Heteropneustes fossilis* (Bloch.). *Bangladesh Journal of Zoology* 2012;40:207-212.
- [123] Laabs V, Wehrhan A, Pinto A, Dores E, Amelung W. Pesticide fate in tropical wetlands of Brazil: An aquatic microcosm study under semi-field conditions. *Chemosphere* 2007;67:975-989.
- [124] Larson SJ, Capel PD, Majewski M. Pesticides in surface waters: Distribution, trends, and governing factors (No. 3). CRC Press;2010.
- [125] Laycock I, Lenthall KM, Barratt AT, Cresswell JE. Effects of imidacloprid, a neonicotinoid pesticide, on reproduction in worker bumble bees (*Bombus terrestris*). *Ecotoxicology* 2012;21:1937-1945.
- [126] le Vieux PD, Malan AP. An overview of the vine mealybug (*Planococcus ficus*) in South African vineyards and the use of entomopathogenic nematodes as potential biocontrol agent. *South African Journal of Enology and Viticulture* 2013;34:108-118.
- [127] Lee DH, Lind PM, Jacobs DR, Salihovic S, van Bavel B, Lind L. Polychlorinated biphenyls and organochlorine pesticides in plasma predict development of type 2 diabetes in the elderly. The Prospective Investigation of the Vasculature in Uppsala Seniors (PIVUS) study. *Diabetes Care* 2011a;34:1778-1784.
- [128] Lee SJ, Mehler L, Beckman J, Diebolt-Brown B, Prado J, Lackovic M, Waltz J, Mulay P, Schwartz A, Mitchell Y, Moraga-McHaley S, Gergely R, Calvert GM. Acute pesticide illnesses associated with off-target pesticide drift from agricultural applications: 11 States, 1998-2006. *Environmental Health Perspectives* 2011b;119:1162.
- [129] Lee WJ, Colt JS, Heineman EF, McComb R, Weisenburger DD, Lijinsky W, Ward MH. Agricultural pesticide use and risk of glioma in Nebraska, United States. *Occupational and Environmental Medicine* 2005;62:786-792.
- [130] Lupwayi NZ, Harker KN, Clayton GW, O'Donovan JT, Blackshaw RE. Soil microbial response to herbicides applied to glyphosate-resistant canola. *Agriculture Ecosystem and Environment* 2009;129:171-176.
- [131] Madhuri RJ, Rangaswamy V. Influence of selected insecticides on phosphatase activity in groundnut (*Arachis hypogaeae* L.) soil. *Journal of Environmental Biology* 2002;23:393-397.
- [132] Martinez-Toledo MV, Salermon V, Rodelas B, Pozo C, Gonzalez-Lopez J. Effects of the fungicide captan on some functional groups of soil microflora. *Applied Soil Ecology* 1998;7:245-255.
- [133] Matthews GA. Improved systems of pesticide application. *Philosophical Transactions of the Royal Society B* 1981;295:163-173.
- [134] Mayanglambam T, Vig K, Singh DK. Quinalphos persistence and leaching under field conditions and effects of residues on dehydrogenase and alkaline phosphomono-

- noesterases activities in soil. *Bulletin of Environmental Contamination and Toxicology* 2005;75:1067-1076.
- [135] McClure MS. Resurgence of the scale, *Fiorinia externa* (Homoptera: Diaspididae) on hemlock following insecticide application. *Environmental Entomology* 1977;6:480-484.
- [136] Mcfadyen REC. Successes in biological control of weeds. Proceedings of the X International Symposium on Biological Control of Weeds 3. Montana State University, Bozeman, Montana, USA. Neal RS (ed.);2000. p3-14.
- [137] McSorley R, Gill HK. Introduction to soil solarization. ENY-062, Department of Entomology and Nematology, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL, USA;2010. (<http://edis.ifas.ufl.edu/in856>) (accessed 14 October 2013).
- [138] Megharaj M, Boul HL, Thiele JH. Effects of DDT and its metabolites on soil algae and enzymatic activity. *Biology and Fertility of Soils* 1999;29:130-134.
- [139] Menendez A, Martínez A, Chiochio V, Venedikian N, Ocampo J A, Godeas A. Influence of the insecticide dimethoate on arbuscular mycorrhizal colonization and growth in soybean plants. *International Microbiology* 2010;2:43-45.
- [140] Mesnage R, Clair E, de Vendômois JS, Seralini GE. Two cases of birth defects overlapping Stratton-Parker syndrome after multiple pesticide exposure. *Occupational and Environmental Medicine* 2010;67:359-359.
- [141] Meyers L, Bull J. Fighting change with change: adaptive variation in an uncertain world. *Trends in Ecology and Evolution* 2002;17:551-557.
- [142] Miedaner T. Breeding wheat and rye for resistance to *Fusarium* diseases. *Plant Breeding* 1997;116: 201-220.
- [143] Mineau P, Whiteside M. Pesticide acute toxicity is a better correlate of US grassland bird declines than agricultural intensification. *PloS One* 2013;8:e57457.
- [144] Mironidis GK, Kapantaidaki D, Bentila M, Morou E, Savopoulou-Soultani M, Vontas J. Resurgence of the cotton bollworm *Helicoverpa armigera* in northern Greece associated with insecticide resistance. *Insect Science* 2013;20:505-512.
- [145] Mitra A, Chatterjee C, Mandal FB. Synthetic chemical pesticides and their effects on birds. *Research Journal of Environmental Toxicology* 2011;5:81-96.
- [146] Mostafalou S, Abdollahi M. Concerns of environmental persistence of pesticides and human chronic diseases. *Clinical and Experimental Pharmacology* 2012;S5:e002. (<http://dx.doi.org/10.4172/2161-1459.S5-e002>).
- [147] Munoz-Leoz B, Ruiz-Romera E, Antigüedad I, Garbisu C. Tebuconazole application decreases soil microbial biomass and activity. *Soil Biology and Biochemistry* 2011;43:2176-2183.

- [148] Murage EW, Voroney PR, Kay BD, Deen B, Beyaert RP. Dynamics and turnover of soil organic matter as affected by tillage. *Soil Science Society of American Journal* 2007;71:1363-1370.
- [149] Murray DL, Taylor PL. Claim no easy victories: evaluating the pesticide industry's Global Safe Use campaign. *World Development* 2000;28:1735-1749.
- [150] NAS. Principles of plant and animal pest control, Vol. 3. Insect management and control. National Academy of Sciences, Washington, DC, USA;1969.
- [151] Niethammer KR, White DH, Baskett TS, Sayre MW. Presence and biomagnification of organochlorine chemical residues in oxbow lakes of northeastern Louisiana. *Archives of Environmental Contamination and Toxicology* 1984;13:63-74.
- [152] Niewiadomska A, Klama J. Pesticide side effect on the symbiotic efficiency and nitrogenase activity of Rhizobiaceae bacteria family. *Polish Journal of Microbiology* 2005;54:43-48.
- [153] Niewiadomska A. Effect of carbendazim, imazetapir and thiram on nitrogenase activity, the number of microorganisms in soil and yield of red clover (*Trifolium pretense* L.). *Polish Journal of Environmental Studies* 2004;13:403-410.
- [154] Nowak A, Nowak J, Klodka D, Pryzbulewska K, Telesinski A, Szopa E. Changes in the microflora and biological activity of the soil during the degradation of isoproturon. *Journal of Plant Diseases and Protection* 2004;19:1003-1016.
- [155] Omar SA, Abdel-Sater MA. Microbial populations and enzyme activities in soil, treated with pesticides. *Water, Air, and Soil Pollution* 2001;127:49-63.
- [156] Ono A, Miyazaki R, Sota M, Ohtsubo Y, Nagata Y, Tsuda M. Isolation and characterization of naphthalene-catabolic genes and plasmids from oil contaminated soil by using two cultivation-independent approaches. *Applied Microbiology and Biotechnology* 2007;74:501-510.
- [157] Oppert B, Morgan TD, Kramer KJ. Efficacy of *Bacillus thuringiensis* Cry3Aa protoxin and protease inhibitors against coleopteran storage pests. *Pest Management Science* 2011;67:568-573.
- [158] Pal R, Tah J. Biodegradation and bioremediation of pesticides in Soil: Its objectives, classification of pesticides, factors and recent developments. *World Journal of Science and Technology* 2012;2:36-41.
- [159] Palmer WA, Heard TA, Sheppard AW. A review of Australian classical biological control of weeds programs and research activities over the past 12 years. *Biological Control* 2010;52:271-287.
- [160] Pan-Germany. Pesticide and health hazards. Facts and figures. 2012;1-16 (www.pan-germany.org/download/Vergift_EN-201112-web.pdf) (accessed on 14 October 2013).

- [161] Paoletti MG. Using bioindicators based on biodiversity to assess landscape sustainability. *Agriculture, Ecosystems and Environment* 1999;74:1-18.
- [162] Parshad VR. Rodent control in India. *Integrated Pest Management Reviews* 1999;4:97-126.
- [163] Parsons KC, Mineau P, Renfrew RB. Effects of pesticide use in rice fields on birds. *Waterbirds* 2010;33:193-218.
- [164] Patnaik GK, Kanungo PK, Moorthy BTS, Mahana PK, Adhya TK, Rao VR. Effect of herbicides on nitrogen-fixation (C₂H₂ reduction) associated with rice rhizosphere. *Chemosphere* 1995;30:339-343.
- [165] Pelosi C, Barot S, Capowicz Y, Hedde M, Vandenbulcke F. Pesticides and earthworms. *Agronomy for Sustainable Development* 2013;1-30. DOI:10.1007/s13593-013-0151-z.
- [166] Pereira L, Fernandes MN, Martinez CB. Hematological and biochemical alterations in the fish *Prochilodus lineatus* caused by the herbicide clomazone. *Environmental Toxicology and Pharmacology* 2013;36:1-8.
- [167] Petrelli G, Mantovani A. Environmental risk factors and male fertility and reproduction. *Contraception* 2002;65:297-300.
- [168] Pettis JS, van Engelsdorp D, Johnson J, Dively G. Pesticide exposure in honey bees results in increased levels of the gut pathogen *Nosema*. *Naturwissenschaften* 2012;99:153-158.
- [169] Prasad SM, Zeeshan M, Kumar D. Toxicity of endosulfan on growth, photosynthesis, and nitrogenase activity in two species of Nostoc (*Nostoc muscorum* and *Nostoc caliccola*). *Toxicological and Environmental Chemistry* 2011;93:513-525.
- [170] Pray CE, Huang J, Hu R, Rozelle S. Five years of Bt cotton in China-the benefits continue. *The Plant Journal* 2002;31:423-430.
- [171] Price DRG, Gatehouse JA. RNAi-mediated crop protection against insects. *Trends in Biotechnology* 2008;26:393-400.
- [172] Reader SM, Miller TE. The introduction into bread wheat of a major gene for resistance to powdery mildew from wild emmer wheat. *Euphytica* 1991;53:57-60.
- [173] Reinecke SA, Reinecke AJ. The impact of organophosphate pesticides in orchards on earthworms in the Western Cape, South Africa. *Ecotoxicity and Environmental Safety* 2007;66:244-251.
- [174] Relyea RA, Diecks N. An unforeseen chain of events: lethal effects of pesticides on frogs at sublethal concentrations. *Ecological Applications* 2008;18:1728-1742.
- [175] Relyea RA, Hoverman JT. Interactive effects of predators and a pesticide on aquatic communities. *Oikos* 2008;117:1647-1658.

- [176] Relyea RA. New effects of Roundup on amphibians: predators reduce herbicide mortality; herbicides induce antipredator morphology. *Ecological Applications* 2012;22:634-647.
- [177] Relyea RA. Predator cues and pesticides: a double dose of danger for amphibians. *Ecological Applications* 2003;13:1515-1521.
- [178] Relyea RA. The lethal impact of Roundup on aquatic and terrestrial amphibians. *Ecological Applications* 2005(a);15:1118-1124.
- [179] Relyea RA. The impact of insecticides and herbicides on the biodiversity and productivity of aquatic communities. *Ecological Applications* 2005(b);15:618-627.
- [180] Richter AR, Fuxa JR. Timing, formulations and persistence of a nuclear polyhedrosis virus and microsporidium for control of the velvetbean caterpillar (Lepidoptera: Noctuidae) in soybeans. *Journal of Economic Entomology* 1984;77:1299-1306.
- [181] Richter ED. Acute human pesticide poisonings. *Encyclopedia of Pest Management* 2002;3-6.
- [182] Romero E, Fernández-Bayo J, Díaz JMC, Nogale, R. Enzyme activities and diuron persistence in soil amended with vermicompost derived from spent grape marc and treated with urea. *Applied Soil Ecology* 2010;44:198-204.
- [183] Rosell G, Quero C, Coll J, Guerrero A. Biorational insecticides in pest management. *Journal of Pesticide Science* 2008;33:103-121.
- [184] Rueda-Ayala V, Rasmussen J, Gerhards R. Mechanical weed control. In: *Precision crop protection-the challenge and use of heterogeneity*. Springer Netherlands;2010. p279-294.
- [185] Singh G, Chahil GS, Jyot G, Battu RS, Singh B. Degradation dynamics of emamectin benzoate on cabbage under subtropical conditions of Punjab, India. *Bulletin of Environmental Contamination and Toxicology* 2013. DOI 10.1007/s00128-013-1013-8
- [186] Sannino F, Gianfreda L. Pesticide influence on soil enzymatic activities. *Chemosphere* 2001;45:417-425.
- [187] Santovito A, Cervella P, Delperio M. Chromosomal aberrations in cultured human lymphocytes treated with the fungicide, Thiram. *Drug and Chemical Toxicology* 2012;35:347-351.
- [188] Saravi SSS, Shokrzadeh M. Role of pesticides in human life in the modern age: a review. In: Stoytcheva M (ed.) *Pesticides in the modern world-risks and benefits*. In-Tech;2011. p4-11.
- [189] Satyavani G, Gopi RA, Ayyappan S, Balakrishnamurthy P, Reddy PN. Toxicity effect of expired pesticides to freshwater fish, *Labeo rohita*. *The Journal of Agriculture and Environment* 2011;12:1-9.

- [190] Scharf ME, Meinke LM, Wright RJ, Chandler LD, Siegfried BD. Metabolism of carbar-yl by insecticide-resistant and susceptible western corn rootworm populations (Coleoptera: Chrysomelidae). *Pesticide Biochemistry and Physiology* 1999;63:85-96.
- [191] Scholz NL, Fleishman E, Brown L, Werner I, Johnson ML, Brooks ML, Michaelmore CL, Schlenk D. A Perspective on modern pesticides, pelagic fish declines, and unknown ecological resilience in highly managed ecosystems. *BioScience* 2012;62:428-434.
- [192] Schreck E, Geret F, Gontier L, Treihou M. Neurotoxic effect and metabolic responses induced by a mixture of six pesticides on the earthworm *Aporrectodea caliginosa* nocturna. *Chemosphere* 2008;71:1832-1839.
- [193] Sebiomo A, Ogundero VW, Bankole SA. Effect of four herbicides on microbial population, soil organic matter and dehydrogenase activity. *African Journal of Biotechnology* 2011;10:770-778.
- [194] Shapiro-Ilan DI, Cottrell TE, Jackson MA, Wood BW. Control of key pecan insect pests using biorational pesticides. *Journal of Economic Entomology* 2013;106:257-266.
- [195] Sharma HC, Ortiz R. Host plant resistance to insects: an eco-friendly approach for pest management and environment conservation. *Journal of Environmental Biology* 2002;23:111-135.
- [196] Sharon E, Chet I, Spiegel Y. Trichoderma as a biological control agent. In: *Biological control of plant-parasitic nematodes*. Springer Netherlands;2011. p183-201.
- [197] Shepard M, Carner GR, Turnipseed TG. Colonization and resurgence of insects pests of soybean in response to insecticides and field isolation. *Environmental Entomology* 1977;6:501-506.
- [198] Shim YK, Mlynarek SP, van Wijngaarden, E. Parental exposure to pesticides and childhood brain cancer: US Atlantic coast childhood brain cancer study. *Environmental Health Perspectives* 2009;117:1002-1006.
- [199] Singh B, Mandal K. Environmental impact of pesticides belonging to newer chemistry. In: Dhawan AK, Singh B, Brar-Bhullar M, Arora R (eds.). *Integrated pest management*. Scientific Publishers, Jodhpur, India;2013. p152-190.
- [200] Singh DK, Kumar S. Nitrate reductase, arginine deaminase, urease and dehydrogenase activities in natural soil (ridges with forest) and in cotton soil after acetamiprid treatments. *Chemosphere* 2008;71:412-418.
- [201] Sofo A, Scopa A, Dumontet S, Mazzatura A, Pasquale V. Toxic effects of four sulphonylureas herbicides on soil microbial biomass. *Journal of Environmental Science and Health, Part B* 2012;47:653-659.
- [202] Solaimalai A, Ramesh RT, Baskar M. Pesticides and environment. In: *Environmental contamination and bioreclamation*; 2004. p345-382.

- [203] Soltani H, Agricultural H. Efficacy of spinosad against potato Colorado beetle *Leptinotarsa*. Hamedan Agricultural and Natural Resources Research, Hamedan, Iran; 2011.
- [204] Son HK, Kim SA, Kang JH, Chang YS, Park SK, Lee SK, Jacobs Jr. DR, Lee DH. Strong associations between low-dose organochlorine pesticides and type 2 diabetes in Korea. *Environment International* 2010;36:410-414.
- [205] Song C, Kanthasamy A, Anantharam V, Sun F, Kanthasamy AG. Environmental neurotoxic pesticide increases histone acetylation to promote apoptosis in dopaminergic neuronal cells: relevance to epigenetic mechanisms of neurodegeneration. *Molecular Pharmacology* 2010;77:621-632.
- [206] Souza RC, Cantão ME, Vasconcelos ATR, Nogueira MA, Hungria M. Soil metagenomics reveals differences under conventional and no-tillage with crop rotation or succession. *Applied Soil Ecology* 2013;72:49-61.
- [207] Srinivasulu M, Mohiddin GJ, Madakka M, Rangaswamy V. Effect of pesticides on the population of *Azospirillum* sp. and on ammonification rate in two soils planted to groundnut (*Arachis hypogaea* L.). *Tropical Ecology* 2012a;53:93-104.
- [208] Srinivasulu M, Mohiddin GJ, Subramanyam K, Rangaswamy V. Effect of insecticides alone and in combination with fungicides on nitrification and phosphatase activity in two groundnut (*Arachis hypogaea* L.) soils. *Environmental Geochemistry and Health* 2012b;34:365-374.
- [209] Stuart SN, Chanson JS, Cox NA, Young BE, Rodrigues ASL, Fischman DL, Waller RW. Status and trends of amphibian declines and extinctions worldwide. *Science* 2004;306:1783-1786.
- [210] Sukul P. Enzymes activities and microbial biomass in soil as influenced by metalaxyl residues. *Soil Biology Biochemistry* 2006;38:320-326.
- [211] Sultana V, Ehteshamul-Haque S, Ara J, Athar M. Effect of brown seaweeds and pesticides on root rotting fungi and root-knot nematode infecting tomato roots. *Journal of Applied Botany and Food Quality* 2012;83:50-53.
- [212] Symondson WOC, Hemingway J. Biochemical and molecular techniques. In: Dent DR, Walton MP (eds.) *Methods in ecological and agricultural entomology*. CAB International, Wallingford, UK;1997. p293-350.
- [213] Tabashnik BE, Van Rensburg JBJ, Carrière Y. Field-evolved insect resistance to Bt crops: definition, theory, and data. *Journal of Economic Entomology* 2009;102:2011-2025.
- [214] Tanner CM, Kamel F, Ross GW, Hoppin JA, Goldman SM, Korell M, Marras C, Bhudhikanok GS, Kasten M, Chade AR, Comyns K, Richards MB, Meng C, Priestley B, Fernandez HH, Cambi F, Umbach DM, Blair A, Sandler DP, Langston JW. Rote-

- none, paraquat, and Parkinson's disease. *Environmental Health Perspectives* 2011;119:866-872.
- [215] Tavela ADO, Araújo JV, Braga FR, Silva AR, Carvalho RO, Araujo JM, Carvalho G. Biological control of cyathostomin (Nematoda: Cyathostominae) with nematophagous fungus, *Monacrosporium thaumasium* in tropical southeastern Brazil. *Veterinary Parasitology* 2011;175:92-96.
- [216] Tien CJ, Chen CS. Assessing the toxicity of organophosphorous pesticides to indigenous algae with implication for their ecotoxicological impact to aquatic ecosystems. *Journal of Environmental Science and Health, Part B* 2012;47:901-912.
- [217] Tierney KB, Baldwin DH, Hara TJ, Ross PS, Scholz NL, Kennedy CJ. Olfactory toxicity in fishes. *Aquatic Toxicology* 2010;96:2-26.
- [218] Trajkovska S, Mbaye M, Seye MG, Aaron JJ, Chevreuil M, Blanchoud H. Toxicological study of pesticides in air and precipitations of Paris by means of a bioluminescence method. *Analytical and Bioanalytical Chemistry* 2009;394:1099-1106.
- [219] Trevisan M, Montepiani C, Ragozza L, Bartoletti C, Loannilli E, Del Re AAM. Pesticides in rainfall and air in Italy. *Environmental Pollution* 1993;80:31-39.
- [220] Tu CM. Effects of some pesticides on enzyme activities in an organic soil. *Bulletin of Environmental Contamination and Toxicology* 1981;27:109-114.
- [221] Van Leeuwen T, Vontas J, Tsagkarakou A, Dermauw W, Tirry L. Acaricide resistance mechanisms in the two-spotted spider mite *Tetranychus urticae* and other important Acari: A review. *Insect Biochemistry and Molecular Biology* 2010;40:563-572.
- [222] Vasileiadis VP, Sattin M, Otto S, Veres A, Palinkas Z, Ban R, Pons X, Kudsk P, Weide van der R, Czembor E, Moonen AC, Kiss J. Crop protection in European maize-based cropping systems: Current practices and recommendations for innovative Integrated Pest Management. *Agricultural Systems* 2011;104:533-540.
- [223] Vermeer K, Risebrough RW, Spaans AL, Reynolds LM. Pesticide effects on fishes and birds in rice fields of Surinam, South America. *Environmental Pollution* 1970;7:217-236.
- [224] Vickerman GP. Farm scale evaluation of the long-term effects of different pesticide regimes on the arthropod fauna of winter wheat. In: Greeves MP, Grieg-Smith PW, Smith BD (eds.) *Field methods for the environmental study of the effects of pesticides*. BCPC Monograph No. 40 British Crop Protection Council, Farnham, UK;1988. p127-135.
- [225] Wang LP, Shen J, Ge LQ, Wu JC, Yang GQ, Jahn GC. Insecticide-induced increase in the protein content of male accessory glands and its effect on the fecundity of females in the brown planthopper *Nilaparvata lugens* Stål (Hemiptera: Delphacidae). *Crop Protection* 2010;29:1280-1285.

- [226] Wang MC, Gong M, Zang HB, Hua XM, Yao J, Pang YJ, Yang YH. Effect of methamidophos and urea application on microbial communities in soils as determined by microbial biomass and community level physiological profiles. *Journal of Environmental Science and Health Part B* 2006;41:399-413.
- [227] Ware GW. Effects of pesticides on nontarget organisms. *Residue Reviews* 1980;76:173-201.
- [228] Weber JB, Wilkerson GG, Reinhardt CF. Calculating pesticide sorption coefficients ($K_{sub(d)}$) using selected soil properties. *Chemosphere* 2004;55:157-166.
- [229] Whitehorn PR, O'Connor S, Wackers FL, Goulson D. Neonicotinoid pesticide reduces bumble bee colony growth and queen production. *Science* 2012;336:351-352.
- [230] Williams T. Silent scourge. *Audubon* 1997;99:28-35.
- [231] Winchester PD, Huskins J, Ying J. Agrichemicals in surface water and birth defects in the United States. *Acta Paediatrica* 2009;98:664-669. DOI:0.1111/j.1651-2227.2008.01207.x
- [232] Wu JY, Smart MD, Anelli CM, Sheppard WS. Honey bees (*Apis mellifera*) reared in brood combs containing high levels of pesticide residues exhibit increased susceptibility to *Nosema* (Microsporidia). *Journal of Invertebrate Pathology* 2012;109:326-329.
- [233] Wu KM, Lu YH, Feng HQ, Jiang YY, Zhao JZ. Suppression of cotton bollworm in multiple crops in China in areas with Bt toxin-containing cotton. *Science* 2008;321:1676-1678.
- [234] Xavier R, Rekha K, Bairy KL. Health perspective of pesticide exposure and dietary management. *Malaysian Journal of Nutrition* 2004;10:39-51.
- [235] Xie H, Gao F, Tan W, Wang SG. A short-term study on the interaction of bacteria, fungi and endosulfan in soil microcosm. *Science of the Total Environment* 2011;412:375-379.
- [236] Xu X, Dailey AB, Talbott EO, Ilacqua VA, Kearney G, Asal NR. Associations of serum concentrations of organochlorine pesticides with breast cancer and prostate cancer in US adults. *Environmental Health Perspectives* 2010;118:60-66.
- [237] Xuan TD, Shinkichi T, Khanh TD, Chung IM. Biological control of weeds and plant pathogens in paddy rice by exploiting plant allelopathy: an overview. *Crop Protection* 2005;24:197-206.
- [238] Yadav BC, Veluthambi K, Subramaniam K. Host-generated double stranded RNA induces RNAi in plant-parasitic nematodes and protects the host from infection. *Molecular and Biochemical Parasitology*. 2006;148:219-222.
- [239] Yasmin S, D'Souza D. Effects of pesticides on the growth and development of earthworm: a review. *Applied and Environmental Soil Sciences* 2010;1-9. (Article ID 678360).

- [240] Zhang LW, Liu YJ, Yao J, Wang B, Huang B, Li ZZ, Sun JH. Evaluation of *Beauveria bassiana* (Hyphomycetes) isolates as potential agents for control of *Dendroctonus valens*. *Insect Science* 2011;18:209-216.
- [241] Zhou X, Scharf ME, Parimi S, Meinke LJ, Wright RJ, Chandler LD, Siegfried BD. Diagnostic assays based on esterase-mediated resistance mechanisms in western corn rootworms (Coleoptera: Chrysomelidae). *Journal of Economic Entomology* 2002;95:1261-1265.

IntechOpen

IntechOpen