

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,500

Open access books available

118,000

International authors and editors

130M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



The Epidemiological, Morphological, and Clinical Aspects of the Aberrant Right Subclavian Artery (Arteria Lusoria)

Michał Polgaj, Ludomir Stefańczyk and
Mirosław Topol

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/64604>

Abstract

The most common embryologic abnormality of the aortic arch is aberrant right subclavian artery (ARSA), known clinically as arteria lusoria (AL). This vessel travels to the right arm, crossing the middle line of the body and usually passing behind the esophagus. If the artery compresses the esophagus, it may produce a condition called dysphagia lusoria. Another commonly reported symptoms related to compression of adjacent structures by arteria lusoria were dyspnea, retrosternal pain, cough, and weight loss greater than 10 kg over a 6-month period. The chapter includes information describing demographic, clinical, and morphological characteristics of presence of arteria lusoria such as gender distribution, frequency in population, frequency of the most commonly reported symptoms related to compression of adjacent structures, coexistence with the most common vascular anomalies and diagnostic procedures. The presence of arteria lusoria together with the right nonrecurrent inferior laryngeal nerve (NRILN) is especially clinically important; during thyroid surgery, the right laryngeal nerve cannot be found at the lower pole of the thyroid, and it may be injured by the surgeon if it is not identified in the aberrant area or found lateral to the thyroid.

Keywords: aberrant right subclavian artery, arteria lusoria, anatomical variations, vessels, clinical symptoms

1. Introduction

The one of the most common embryologic vascular abnormalities of the aortic arch is an aberrant right subclavian artery (ARSA), known clinically as arteria lusoria (AL) [1].

In approximately 80% of individuals, three arteries arise from the arch of the aorta. From right to left, the brachiocephalic trunk arises (divided into the right common carotid artery and the right subclavian artery), followed by the left common carotid artery, and finally the left subclavian artery (**Figure 1**) [2].




Figure 1. Three-dimensional computed tomography reconstruction of the arteries (CT-64-row MDCT scanner, Light-speed VCT, GE, Waukesha, Wisconsin, USA). Ao—arch of the aorta, BCT—brachiocephalic trunk, LSA—left subclavian artery, LCCA—left common carotid artery, RCCA—right common carotid artery, RSA—right subclavian artery.

When an aberrant right subclavian artery (arteria lusoria) is present, the brachiocephalic trunk is absent and four arteries arise from the arch of the aorta: the right common carotid artery followed by the left common carotid artery, the left subclavian artery, and finally the right subclavian artery, with the most distal left-sided origin (**Figures 2–4**). This vessel, the aberrant right subclavian artery, travels to the right arm, crossing the middle line of the body and usually passing behind the esophagus (**Figure 3**) [3, 4]. If the artery compresses neighboring structures or organs, it may produce symptoms, the most common example being compression of the esophagus by the arteria lusoria, which results in a condition called dysphagia lusoria [4].

Although the first description of an aberrant right subclavian artery was provided in 1735 by Hunauld, the clinical entity was presented later [2]. In 1794, David Bayford, a physician from London, England, described a 33-year-old woman who succumbed to malnutrition after 20 years of progressive dysphagia [3]. At autopsy, Dr. Bayford noted esophageal compression by an abnormal right subclavian artery and suggested the term “dysphagia lusoria” to describe

this syndrome. Hence, it is also known as Bayford–Autenrieth dysphagia [3]. Kommerell described its radiological findings in 1936 [4].




Figure 2. Three-dimensional computed tomography reconstruction of the arteries (CT-64-row MDCT scanner, Light-speed VCT, GE, Waukesha, Wisconsin, USA). Ao—arch of the aorta, AL—arteria lusoria, LSA—left subclavian artery, LCCA—left common carotid artery, RCCA—right common carotid artery.




Figure 3. Computed tomography transverse scan on the level of the arch of the aorta. Ao—arch of the aorta, AL—arteria lusoria, E—esophagus, KD—Kommerell's diverticulum, T—trachea.




Figure 4. Three-dimensional computed tomography reconstruction of the arteries (CT-64-row MDCT scanner, Light-Speed VCT, GE, Waukesha, Wisconsin, USA). Ao—arch of the aorta, AL—arteria lusoria, KD—Kommerell's diverticulum, LSA—left subclavian artery, LCCA—left common carotid artery, RCCA—right common carotid artery.

2. Embryology

Arteria lusoria results from abnormal embryologic development of the aortic arch. In the normal situation, between the 4th and 5th weeks of embryonic life, blood leaves the heart by a common trunk called the “truncus arteriosus”, which divides into two branches termed the ventral aortae. These branches are connected with the paired dorsal aortae by six aortic arches. The carotid system is formed by segments of the first three arches. The right fourth arch, a segment of the right ventral aorta, and a portion of the right dorsal aorta develop into the right subclavian artery. The left fourth arch persists as the adult aortic arch with the anlagen of the seventh dorsal intersegmental artery, and this forms the left subclavian artery. The fifth arches are both resorbed, and the sixth arches form the pulmonary artery and the ductus arteriosus [5, 6].

The aberrant origin of the right subclavian artery is caused by the involution of the right fourth vascular arch and proximal right dorsal aorta and the persistence of the seventh intersegmental artery originating from the proximal descending thoracic aorta, resulting in the arteria lusoria following an abnormal course [5, 6].

3. Epidemiological and demographic characteristics

The frequency of ARSA varies throughout the world. In Europe, depending on the country, it has been found in 0.11% (Great Britain) [7], 0.16% (Greece) [8], 0.3% (France) [9], or 0.36% (the Netherlands) [10] of the population. Studies have also been performed on other continents: Asia, 0.1–0.2% of cases (respectively: China and Japan) [11, 12]; North America, 0.5% of cases (United State) [13]; or Australia and Oceania, 0.8% of cases (New Zealand) [14] (**Table 1**). However, its detection arguably depends primarily on the sensitivity of the diagnostic procedures employed, for example, cadaveric or CT versus chest X-ray examination.

AL (%)	Country	Type of investigation	N/(total)	Researcher
0.1	China	Angiography	3/3000	Nie et al. [11]
0.11	Great Britain	Gastroscopy	1/920	Kelly [7]
0.16	Greece	Angiography	1/622	Natsis et al. [8]
0.2	Japan	Autopsy	1/516	Saito et al. [12]
0.3	France	coronarography	11/3730	Abhaichand et al. [9]
0.36	The Netherlands	Endoesophageal ultrasonography	12/3334	De Luca et al. [10]
0.5	USA	Computed tomography	36/7174	Haesemeyer and Gavant [13]
0.8	New Zealand	Autopsy	19/2291	Cainey [14]

AL—arteria lusoria.

Table 1. Frequency of aberrant right subclavian artery (arteria lusoria) in different populations.

The gender distribution of the aberrant right subclavian artery was found to be similar: 55.3% females versus 44.7% males [15].

4. Clinical characteristics

An aberrant right subclavian artery is usually asymptomatic, while about 8–10% of adult patients develop symptoms [16]. The anomaly does not cause symptoms in most patients and can be discovered incidentally during life or found at autopsy [17, 18].

When present, symptoms usually occur at the two extremes of life [19]. In children, respiratory symptoms are prevalent, mainly dyspnea or chronic coughing. They can also present with repetitive respiratory infections. Infant patients may demonstrate an increased frequency of pulmonary infections, which is thought to be due to absence of tracheal rigidity in combination with dysphagia and aspiration [19, 20]. As noted above, a rare cause of dysphagia observed in adults is compression of the esophagus by an abnormal course of the aberrant right subclavian artery, which is classically termed “dysphagia lusoria” [3].

Although dysphagia may be the most frequent symptom demonstrated by adults, children demonstrate a different respiratory symptomatology attributed to lack of the tracheal rigidity associated with dysphagia and false routes [21]. In addition, the arteria lusoria can also be revealed by the extension of aortic dissection or by peripheral arterial embolism.

A meta-analysis by Polgaj et al. found the most commonly reported symptoms related to compression of adjacent structures by an aberrant right subclavian artery to be dysphagia (71.2%), dyspnea (18.7%), retrosternal pain (17.0%), cough (7.6%), and weight loss greater than 10 kg over a six-month period (5.9%) [15]. Among the less common symptoms, stomachache, back pain, and numbness of the right upper limb were reported. The mean age of the onset of symptoms was 49.9 ± 19.4 years for the whole group (data shown as mean \pm standard deviation). However, the mean age according to gender was 44.9 ± 18.1 years for males and 54.0 ± 19.6 years for females. This difference was statistically significant [15].

Dysphagia also frequently occurs in elderly patients, for which four mechanisms have been proposed: increased rigidity of the trachea leading to easy compression of esophagus, aneurysm formation, presence of Kommerell's diverticulum, elongation of the aorta, the coexistence of an aberrant artery with a truncus bicaroticus, or a close origin of common carotid arteries from the arch of the aorta [22–25].

To the angiographer who uses the right axillary, brachial or radial approach to the ascending thoracic aorta, the arteria lusoria is also a clinically important element. The presence of an ARSA is suspected in cases in which catheterization of the ascending aorta proves difficult. Using the right radial approach, access to the ascending aorta is usually easy [11]. Previous studies indicate that only 60% of such cases were successfully performed by transradial approach in the setting of AL [26]. This variant makes the right transradial route difficult to approach the ascending aorta, as it requires the catheter to curve back to reach the aortic root [26, 27]. However, the repeated entry of the guide wire from the right subclavian artery to the descending aorta rather than the ascending aorta should indicate this possibility. Thus, angiography can prove to be very challenging in the presence of an arteria lusoria [11, 26, 27].

Finally, the inferior right recurrent laryngeal nerve is an asymptomatic variation anomaly, which can be an important obstacle and be seriously damaged during cervicotomy, thyroid, and parathyroid surgery. In such cases, the inferior right recurrent laryngeal nerve is a classic risk and must be eliminated by location and routine dissection of the nerve [28–30]. This is of particular importance when the diagnosis concerns an asymptomatic neural anomaly discovered by dissection or a vascular anomaly whose symptoms are very variable [30].

5. Morphological characteristics

The literature presents two main classifications of the aberrant right subclavian artery. According to Neuhauser's threefold classification, the first type of arteria lusoria crosses the posterior wall of the esophagus, and this is observed in more than 80% of the cases. In the

second type, this artery passes between the trachea and the esophagus (15% of the cases), and in the third type, it crosses the midline of the body ahead of the trachea (in 5% of the cases) [23].

In contrast, the Adachi and Williams classification recognizes four basic morphological types. Type I/G is characterized by an aberrant right subclavian artery arising from the arch of the aorta as the final branch. Type II/CG is similar to the first type, but an additional left vertebral artery arises from the arch of the aorta. In type III/H, three arteries arise from the arch of the aorta: as the first common trunk of the common carotid arteries (truncus bicaroticus), as the left subclavian artery, and as the last aberrant right subclavian artery. In type IV/N, the aberrant left subclavian artery arises from the right-sided arch of the aorta as the final branch [31].

The most common vascular anomalies coexisting with an aberrant right subclavian artery (arteria lusoria) were found to be truncus bicaroticus (19–29%), Kommerell's diverticulum (15–60%), aneurysm (just after the origin of arteria lusoria) (13%), and right-sided aortic arch (9%) [15]. Klinkhamer regards truncus bicaroticus is a precondition for tracheal–esophageal compression and the development of clinical symptoms. Under these circumstances, the truncus bicaroticus holds the trachea from the front, and the aberrant right subclavian artery compresses the esophagus from behind [22].

In 1936, Kommerell published the first radiological findings of the route of the aortic arch as an aortic diverticulum (Kommerell's diverticulum), which was identified as being located at the origin of an aberrant subclavian artery [4] (**Figure 3** and **4**). Kommerell's diverticulum is usually found incidentally on a chest roentgenogram and is often misdiagnosed as a mediastinal tumor [29]. Kommerell's diverticulum is a normal broadening of the proximal origin of the aberrant right subclavian artery from the aortic arch and is most frequently present in patients with a right aortic arch and an aberrant left subclavian artery [32]. However, Kommerell's diverticulum is not analogous with an aneurysm: The primary indication for surgical repair of Kommerell's diverticulum is a diameter larger than 50 mm and the presence of clinical symptoms.

The identification of arteria lusoria should alert the radiologist and surgeon that a nonrecurrent inferior laryngeal nerve (NRILN) is present and that an anticipating surgical technique should be performed to reduce the risk of neural injury. Due to its anatomical position, an NRILN is not only at risk of being damaged during thyroidectomy, but also during such other surgical procedures as neck dissection, parathyroidectomy, and carotid endarterectomy [30, 33].

6. Diagnosis

The diagnosis of arteria lusoria was reported only on anatomical dissection until 1936, when Burckhard Kommerell described the clinical diagnosis of an aberrant right subclavian artery that originated from an aortic diverticulum, later known as Kommerell's diverticulum, in a 65-year-old man who was believed to have stomach cancer [4]. In 1946, Gross was the first to report the surgical treatment of dysphagia lusoria, in a four-month-old infant [34].

The diagnostic modalities available to visualize an arteria lusoria include barium esophagogram, computed tomography (CT), magnetic resonance imaging (MRI), digital subtraction angiography (DSA), endoscopy and endoscopic ultrasound. New advances in CT technology allow even small vascular structures to be visualized in detail. Multidetector computed tomography (MDCTA) is now an established diagnostic test in the evaluation of many vascular diseases [35–39].

Barium contrast examination of the esophagus shows a characteristic, extrinsic, smooth diagonal impression at the level of the third and fourth dorsal vertebra. Lateral or oblique views show the extrinsic impression to be posterior, and in case of arteria lusoria, just above the level of the aortic arch. As dysphagia occurs frequently with ingestion of solid foods, including a barium soaked bread bolus may improve localization of the defect [35].

Digital subtraction angiography gives valuable information regarding AL. It is an invasive procedure and, in contrast to MDCT, has the disadvantage in showing extravascular structures such as the esophagus. It has also been shown that the effective radiation doses in MDCT angiography studies are moderate and even lower than those associated with DSA in a comparable patient group.

CT or MRI (magnetic resonance imaging) angiography has replaced conventional angiography and is the gold standard for the diagnosis. It not only confirms the diagnosis but also helps to exclude aneurysm of the aorta or other associated anomalies and to plan the operation [36–39].

MRI has the advantage of being a noninvasive procedure and the patient is spared the potential risk of intravenous contrast agents. MRI is not as useful as MDCT due to the generation of respiratory and cardiac motion artifacts. Also it is not a preferred method due to its cost and prolonged scan time. Although MR angiography may reveal the presence of a vascular anomaly, the information regarding nonvascular mediastinal structures is insufficient [36–39].

Endoscopy may reveal pulsatile, shelf-like extrinsic compression in the posterior wall of the esophagus, with intact mucosa. Such an area of narrowing is usually located between 20 and 24 cm from the mouth [7].

Endoscopic ultrasound (EUS) can identify an arteria lusoria, as it lies close to the esophagus. EUS is regarded as the most accurate test for the evaluation of the esophageal wall and the surrounding structures, with the incorporation of Doppler technology in modern echoendoscopes allowing particularly accurate examination of adjacent vessels [10].

The combination of MDCT with 3D volume rendering images provides further advantages. These allow not only the depiction of the thoracic vascular anomalies but also more accurate assessment of the diameter, angle, and compressed area of the esophagus and the relationship between the AL with the esophagus and other mediastinal structures. In addition, MDCT is a noninvasive procedure, unlike DSA, and offers easier application and a shorter time requirement than DSA or MRA [36–39].

7. Conclusion

A familiarity with the anatomy of the some types of vascular anomalies is necessary for clinicians involved in many medical areas. Compression of adjacent structures by an arteria lusoria needs to be differentiated from other conditions presenting symptoms such as dysphagia, dyspnea, retrosternal pain, cough, and weight loss. The knowledge presented in this chapter will allow the best healthcare to be provided for patients.

Author details

Michał Polguj^{1*}, Ludomir Stefańczyk² and Mirosław Topol³

*Address all correspondence to: michal.polguj@umed.lodz.pl

1 Department of Angiology, Medical University of Łódź, Łódź, Poland

2 Department of Radiology, Medical University of Łódź, Łódź, Poland

3 Department of Normal and Clinical Anatomy, Medical University of Łódź, Łódź, Poland

References

- [1] Molz G, Burri B. Aberrant subclavian artery (arteria lusoria): sex differences in the prevalence of various forms of the malformation. Evaluation of 1378 observations. *Virchows Arch A Pathol Anat Histol.* 1978, 380(4): 303–15
- [2] Hunauld PM. Examen de quelques parties d'un singe. *Hist Acad Roy Sci.* 1735;2:516–23.
- [3] Bayford D. An account of a singular case of obstructed deglutition. *Mem M Society London.* 1794, 2, 275–86.
- [4] Kommerell B. Verlagerung des Oesophagus durch eine abnorm verlaufende Arteria subclavia dextra (arteria lusoria). *Fortschr Geb Roentgenstrahlen.* 1936, 54: 590–95.
- [5] Kopp R, Wizgall J, Kreuzer E, Meimarakis G, Weidenhagen R, Kiiknl A, Conrad C, Jauch KW, Lauterjun L. Surgical and endovascular treatment of symptomatic aberrant right subclavian artery (arteria lusoria). *Vascular.* 2007, 15: 84–91.
- [6] Inami T, Seino Y, Mizuno K. Unique case of giant Kommerell diverticulum with aberrant left subclavian artery arising from the left aortic arch associated with situs inversus. *Int J Cardiol.* 2013, 163(3): e47–48.
- [7] Kelly MD. Endoscopy and the aberrant right subclavian artery. *Am Surg.* 2007, 73(12): 1259–61.

- [8] Natsis KI, Tsitouridis IA, Didagelos MV, Fillipidis AA, Vlasits KG, Tsikaras PD. Anatomical variations in the branches of the human aortic arch in 633 angiographies: clinical significance and literature review. *Surg Radiol Anat.* 2009, 31(5): 319–23.
- [9] Abhaichand RK, Louvard Y, Gobeil JF, Loubeyre C, Lefèvre T, Morice MC. The problem of arteria lusoria in right transradial coronary angiography and angioplasty. *Catheter Cardiovasc Interv.* 2001, 54(2): 196–01.
- [10] De Luca L, Bergman JJ, Tytgat GN, Fockens P. EUS imaging of the arteria lusoria: case series and review. *Gastrointest Endosc.* 2000, 52(5): 670–73.
- [11] Nie B, Zhou YJ, Li GZ, Shi DM, Wang JL. Clinical study of arterial anatomic variations for transradial coronary procedure in Chinese population. *Chin Med J (Engl)* 2009, 122(18): 2097–02.
- [12] Saito T, Tamatsukuri Y, Hitosugi T, Miyakawa K, Shimizu T, Oi Y, Yoshimoto M, Yamamoto Y, Spaniel-Browski K, Steinke H. Three cases of retroesophageal right subclavian artery. *J Nippon Med Sch.* 2005, 72(6): 375–82.
- [13] Haesemeyer SW, Gavant ML. Imaging of acute traumatic aortic tear in patients with an aberrant right subclavian artery. *AJR Am J Roentgenol.* 1999, 172(1): 117–20.
- [14] Cainey J. Tortuosity of the cervical segment of the internal carotid artery. *J Anat.* 1924, 59(1): 87–96.
- [15] Polguj M, Chrzanowski Ł, Kasprzak J, Stefanczyk L, Topol M, Majos A. The aberrant right subclavian artery (arteria lusoria)—the morphological and clinical aspects of the one of the most important variations: a systematic study of 141 reports. *Sci World J.* 2014;2014:292734.
- [16] Delap TG, Jones SE, Johnson DR. Aneurysm of an aberrant right subclavian artery presenting as dysphagia lusoria. *Ann Otol Rhinol Laryngol.* 2000, 109(2): 231–34.
- [17] Adachi B. *Das Arteriensystem der Japener.* Kenkyusha Press, Tokyo, Japan, 1928.
- [18] Janssen M, Baggen MGA, Veen HF, Smout AJ, Bekkers JA, Jonkman JG, Ouwendijk RJ. Dysphagia lusoria: clinical aspects, manometric findings, diagnosis and therapy. *Am J Gastroenterol.* 2000, 96: 1411–16.
- [19] Puri SK, Ghuman S, Narang P, Sharma A, Singh S. CT and MR angiography in dysphagia lusoria in adults. *Ind J Radiol Imag.* 2005, 15(4): 497–01.
- [20] Van Son JA, Julsrud PR, Hagler DJ, Sim EK, Pairolero PC, Puga FJ, Schaff HV, Danielson GK. Surgical treatment of vascular rings: the Mayo Clinic experience. *Mayo Clin Proc.* 1993, 68: 1056–63.
- [21] McNally PR, Rak KM. Dysphagia lusoria caused by persistent right aortic arch with aberrant left subclavian artery and diverticulum of kommerell. *Dig Dis Sci.* 1992, 37: 144–49.

- [22] Klinkhamer AC. Aberrant right subclavian artery: clinical and roentgenologic aspects. *Am J Roentgenol Radium Ther Nucl Med.* 1966, 97: 438–46.
- [23] Levitt B, Richter JE. Dysphagia lusoria: a comprehensive review. *Dis Esophagus.* 2007, 20: 455–60.
- [24] Van Son JA, Mierzwa M, Mohr FM. Resection of atherosclerotic aneurysm at origin of aberrant right subclavian artery. *Eur J Cardiovasc Surg.* 1999, 16: 576–79.
- [25] Ulger Z, Ozyurek AR, Levent E, Gurses D, Parlar A. Arteria lusoria as cause of dysphagia. *Acta Cardiol.* 2004, 59(4): 445–47.
- [26] Valsecchi O, Vassileva A, Musumeci G, Rossini R, Tespili M, Guagliumi G, Mihalcsik L, Gavazzi A, Ferrazzi P. Failure of transradial approach during coronary interventions: anatomic considerations. *Catheter Cardiovasc Interv.* 2006, 67: 870–78.
- [27] Huang IL, Hwang HR, Li SC, Chen CK, Liu CP, Wu MT. Dissection of arteria lusoria by transradial coronary catheterization: a rare complication evaluated by multidetector CT. *J Chin Med Assoc.* 2009, 72(7):379–81.
- [28] Nakatani T, Tanaka S, Mizukami S, Okamoto K, Shiraishi Y, Nakamura T. Retroesophageal right subclavian artery originating from the aortic arch distal and dorsal to the left subclavian artery. *Ann Anat.* 1996, 178(3): 269–71.
- [29] Epstein DA, Debord JR. Abnormalities associated with aberrant right subclavian arteries-a case report. *Vasc Endovascular Surg.* 2002, 36(4): 297–03.
- [30] Avisse C, Marcus C, Delattre JF, Marcus C, Cailliez-Tomasi JP, Palot JP, Ladam-Marcus V, Menanteau B, Flament JB. Right nonrecurrent inferior laryngeal nerve and arteria lusoria: the diagnostic and therapeutic implications of an anatomic anomaly. Review of 17 cases. *Surg Radiol Anat.* 1998, 20(3):227–32.
- [31] Williams GD, Aff HM, Schmeckebier M, Edmonds HW, Grand EG. Variations in the arrangement of the branches arising from the aortic arch in the American whites and Negroes. *Anat Rec.* 1932, 54: 247–51.
- [32] Hartyanszky IL, Lozsadi K, Marcsek P, Huttl T, Sapi E, Kovacs AB. Congenital vascular rings: surgical management of 111 cases. *Eur J Cardiothorac Surg.* 1989, 3(3): 250–54.
- [33] Devèze A, Sebag F, Hubbard J, Jaunay M, Maweja S, Henry JF. Identification of patients with a non-recurrent inferior laryngeal nerve by duplex ultrasound of the brachiocephalic artery. *Surg Radiol Anat.* 2003, 25(3–4):263–9
- [34] Gross RE. Surgical treatment for dysphagia lusoria. *Ann Surg.* 1946, 124:532–34.
- [35] Branscom JJ, Austin JH. Aberrant right subclavian artery. Findings seen on plain chest roentgenograms. *Am J Roentgenol Radium Ther Nucl Med.* 1973, 119(3): 539–42.

- [36] Alper F, Akgun M, Kantarci M, Eroglu A, Ceyhan E, Onbas O, Duran C, Okur A. Demonstration of vascular abnormalities compressing esophagus by MDCT: Special focus on dysphagia lusoria. *Eur J Radiol.* 2006, 59(1):82–7.
- [37] Godlewski J, Widawski T, Michalak M, Kmiec Z Aneurysm of the aberrant right subclavian artery – a case report *Pol J Radiol*, 2010, 75(4): 47–50
- [38] Chen X, Qu YJ, Peng ZY, Lu JG, Ma XJ. Diagnosis of congenital aortic arch anomalies in chinese children by multi-detector computed tomography angiography. *J Huazhong Univ Sci Technolog Med Sci.* 2013, 33(3): 447–51.
- [39] Yang M, Mo XM, Jin JY, Wu M, Liu B, Liu ZY, Gao XC, Tang WW, Teng GJ. Diagnostic value of 64 multislice CT in typing of congenital aortic anomaly in neonates and infants. 2010, 90(31): 2167–71.