

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,400

Open access books available

117,000

International authors and editors

130M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Introductory Chapter: Hormonal Regulation in Plant Development and Stress Tolerance

Mohamed A. El-Esawi

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.69806>

1. Introduction

Phytohormones are regulatory compounds produced in low concentrations and serve as chemical messengers to regulate various plant physiological and developmental processes. They also play essential roles in signal transduction pathways during stress response and regulate internal and external stimuli [1]. Phytohormones comprise five main groups, namely auxins (IAAs), cytokinins (CKs), abscisic acid (ABA), gibberellins (GAs) and ethylene (ET). Salicylates (SAs), jasmonates (JAs), brassinosteroids (BRs), strigolactones (SLs), polyamines, and some peptides represent new families of phytohormones.

Hormone actions form a signaling network and regulate various systems in plants. The interacting actions among hormone signal transduction cascades are called crosstalk [2, 3]. Phytohormones interact by activating a phosphorylation cascade or a common second messenger. Furthermore, several phytohormones interact together, forming a defense network against environmental stresses such as JA, SA, and ABA which play a crucial role in regulating signaling pathways [3]. Understanding the crosstalk between these phytohormones and defense signaling pathways helps reveal new important targets for developing host resistance mechanisms [3, 4]. Here, the current work presents an overview and discusses recent progresses of phytohormone roles and their crosstalk in plant development and stress tolerance.

2. Phytohormones signaling roles

2.1. Abscisic acid

Abscisic acid (ABA), termed stress hormone, plays an important role in plant leaves abscission and abiotic stresses tolerance [3]. ABA also has important roles in various plant developmental and physiological processes such as seed dormancy, embryo morphogenesis, stomatal opening, cell turgor maintenance, and biosynthesis of lipids and storage proteins [3, 5]. ABA regulates protein-encoding genes [6]. ABA enables plants to survive under severe environmental factors [7] and water-deficit conditions [8]. ABA is also important for root growth and architectural modifications under nitrogen deficiency [9] and drought stress [10]. Furthermore, ABA is involved in the biosynthesis of dehydrins, osmoprotectants and protective proteins [3, 11, 12].

2.2. Auxins

Some pathways for auxin (IAA) biosynthesis in plants have been reported so far including one tryptophan-independent and four tryptophan-dependent pathways [3, 13]. IAA plays an important role in plant growth and development as well as in regulating growth under stress factors [14]. IAA plays essential roles in plant adaptation to salinity [15] and heavy metal stresses [16]. Furthermore, auxins induce the transcription of the primary auxin response genes which are identified in various plants such as rice, *Arabidopsis* and soybean [3, 17]. Auxin also regulates crosstalk between biotic and abiotic stresses [18].

2.3. Cytokinins

Cytokinins (CKs) regulate plant growth and development [3, 19]. They are also involved in abiotic stresses [20] such as salinity [21] and drought [19]. They are also important for various crop traits such as productivity and enhanced stress tolerance [3, 22]. CKs also release seeds from dormancy [18] and are considered as abscisic acid antagonists [23]. Decreased CK content promotes apical dominance, which assists in the adaptation to drought stress [3, 20].

2.4. Ethylene

Ethylene (ET) is a gaseous phytohormone regulating plant growth and developmental processes, including flower senescence, fruit ripening, and petal and leaf abscission, as well as regulating stress responses [3, 24, 25]. Ethylene biosynthesis begins from methionine via S-adenosyl-L-methionine and the cyclic amino acid ACC. ACC synthase converts S-adenosyl-L-methionine to ACC, whereas ACC oxidase catalyzes the conversion of ACC to ET. Various abiotic stresses affect endogenous ethylene levels in plant species. Higher ET concentrations promote stress tolerance [26]. Ethylene may combine with other hormones such as jasmonates and salicylic acid and plays crucial roles in regulating plant defense against biotic stress factors [3, 1]. Ethylene and abscisic acid may act together to regulate plant growth and development [3].

2.5. Gibberellins

Gibberellins (GAs) are carboxylic acids that may regulate plant growth and development [27]. They positively regulate leaf expansion, seed germination, stem elongation, flower development and trichome initiation [3, 28]. They also play important role in abiotic stress tolerance [29] such as osmotic stress. GAs may interact with other hormones and regulate various developmental processes [30]. These interactions may involve both negative and positive regulatory roles [3, 30].

2.6. Brassinosteroids

Brassinosteroids (BRs) comprise polyhydroxy steroidal phytohormones which regulate plant growth and developmental processes including root and stem growth, and flower initiation and development [3]. BRs were first isolated from *Brassica napus*. Brassinolide, 24-epibrassinolide, and 28-homobrassinolide are the most bioactive BRs widely used in physiological studies [31]. They are found in flower buds, pollen, fruits, vascular cambium, seeds, leaves, roots, and shoots [32]. BRs also play important roles in abiotic stress responses such as chilling, high temperature, soil salinity, drought, light, flooding, and organic pollutants [3].

2.7. Jasmonates

Jasmonates (JAs) are multifunctional phytohormones derived from the membrane fatty acids metabolism and are widely distributed in several plant species [3]. JAs play crucial roles in growth and developmental processes such as fruiting, flowering, senescence and secondary metabolism [3, 33]. JAs are also involved in biotic and abiotic stress responses such as salinity, drought, irradiation and low temperature [3, 34]. Exogenous concentrations of methyl jasmonate (MeJA) minimize salinity stress symptoms [35]. Additionally, endogenous levels of JA are induced in roots under salinity stress [36]. JA levels also reduce heavy metal stress through inducing the antioxidant machinery [3, 37]. MeJA accumulates phytochelatin, conferring tolerance against Cu and Cd stress [38].

2.8. Salicylic acid

Salicylic acid (SA) is a phenolic compound which regulates the expression of pathogenesis-associated proteins [39]. SA plays an important role in plant growth and development, as well as in biotic and abiotic stress responses [3, 40]. SA has two biosynthesis pathways: the major isochorismate (IC) pathway and the phenylalanine ammonia-lyase (PAL) pathway. Low levels of SA promote the plant antioxidant capacity [3]. However, the high SA levels may result in cell death [41]. SA comprises genes encoding chaperones, antioxidants, heat shock proteins, and secondary metabolite biosynthetic genes such as cinnamyl alcohol dehydrogenase, sinapyl alcohol dehydrogenase and cytochrome P450 [3, 41]. SA may also combine with ABA to regulate drought response [39]. However, the SA mechanism in abiotic stress tolerance remains mainly unknown and still needs more investigations.

2.9. Strigolactones

Strigolactones (SLs) are carotenoid-derived compounds, produced in small quantities in roots, or synthesized in several plant species [3, 42]. SLs play an important role in root architecture and development [43]. They induce nodulation during interaction processes [44] and may be used for inducing the parasitic plants seed germination [45]. SLs are also involved in biotic and abiotic responses [3].

3. Phytohormones crosstalk

Sessile plants should maintain growth plasticity and adaptation ability to severe environmental conditions. Stress-responsive hormones assist in the alteration of cellular dynamics and thus regulating plant growth under stress conditions [3, 46]. The interacting actions among hormone signal transduction cascades are called crosstalk and form a signaling network [2, 3]. In this case, hormones interact by activating a phosphorylation cascade or a common second messenger. Several phytohormones interact together forming a defense network against environmental stresses such as JA, SA, and ABA which play a crucial role in regulating signaling pathways [3]. Understanding the crosstalk between these phytohormones and defense signaling pathways helps reveal new important targets for developing host resistance mechanisms [3, 4].

A complex signaling network regulates stomatal closure. ABA regulates gene expression which mediates root growth maintenance and water uptake. ABA interacts with signaling molecules and other phytohormones such as nitric oxide and JA to induce stomatal closure, as well as to induce genes controlling response to cytokinin, ethylene or auxin [2, 3]. Furthermore, exogenous treatment of ABA down-regulated the key cytokinin biosynthetic pathway gene, termed isopentenyltransferase, but up-regulated genes encoding cytokinin dehydrogenases and oxidases [3, 21]. GA is also included in the hormonal crosstalk in environmental signals [47]. In conclusion, like the potential use of molecular and genetic markers in crop improvement [48–57], phytohormones play crucial roles in development and stress tolerance of crops.

Author details

Mohamed A. El-Esawi

Address all correspondence to: mohamed.elesawi@science.tanta.edu.eg

1 Botany Department, Faculty of Science, Tanta University, Tanta, Egypt

2 The Sainsbury Laboratory, University of Cambridge, Cambridge, United Kingdom

References

- [1] Kazan K. Diverse roles of jasmonates and ethylene in abiotic stress tolerance. *Trends in Plant Science*. 2015;**20**:219-229
- [2] Harrison MA. Cross-talk between phytohormone signaling pathways under both optimal and stressful environmental conditions. In: Khan NA, Nazar R, Iqbal N, Anjum NA, editors. *Phytohormones and Abiotic Stress Tolerance in Plants*. Berlin, Heidelberg: Springer-Verlag; 2012. pp. 49-76
- [3] Wani SH, Kumar V, Shriram V, Sah SK. Phytohormones and their metabolic engineering for abiotic stress tolerance in crop plants. *The Crop Journal*. 2016;**4**:162-176
- [4] Grant MR, Jones JD. Hormone (dis)harmony moulds plant health and disease. *Science*. 2009;**324**:750-752
- [5] Sreenivasulu N, Radchuk V, Alawady A, Borisjuk L, Weier D, Staroske N, Fuchs J, Miersch O, Strickert M, Usadel B, Wobus U, Grimm B, Weber H, Weschke W. De-regulation of abscisic acid contents causes abnormal endosperm development in the barley mutant *seg8*. *The Plant Journal*. 2010;**64**:589-603
- [6] Nemhauser JL, Hong F, Chory J. Different plant hormones regulate similar processes through largely non overlapping transcriptional responses. *Cell*. 2006;**126**:467-475
- [7] Keskin BC, Sarikaya AT, Yuksel B, Memon AR. Abscisic acid regulated gene expression in bread wheat. *Australian Journal of Crop Science*. 2010;**4**:617-625
- [8] Wilkinson S, Kudoyarova GR, Veselov DS, Arkhipova TN, Davies WJ. Plant hormone interactions: Innovative targets for crop breeding and management. *Journal of Experimental Botany*. 2012;**63**:3499-3509
- [9] Zhang S, Hu J, Zhang Y, Xie XJ, Knapp A. Seed priming with brassinolide improves lucerne (*Medicago sativa* L.) seed germination and seedling growth in relation to physiological changes under salinity stress. *Australian Journal of Agricultural Research*. 2007;**58**:811-815
- [10] Giuliani S, Sanguineti MC, Tuberosa R, Bellotti M, Salvi S, Landi P. Root-ABA1 a major constitutive QTL affects maize root architecture and leaf ABA concentration at different water regimes. *Journal of Experimental Botany*. 2005;**56**:3061-3070
- [11] Verslues PE, Agarwal M, Katiyar-Agarwal S, Zhu J, Zhu JK. Methods and concepts in quantifying resistance to drought, salt and freezing, abiotic stresses that affect plant water status. *The Plant Journal*. 2006;**45**:523-539
- [12] Sreenivasulu N, Harshavardhan VT, Govind G, Seiler C, Kohli A. Contrapuntal role of ABA: Does it mediate stress tolerance or plant growth retardation under long-term drought stress? *Gene*. 2012;**506**:265-273

- [13] Mano Y, Nemoto K. The pathway of auxin biosynthesis in plants. *Journal of Experimental Botany*. 2012;**63**:2853-2872
- [14] Kazan K. Auxin and the integration of environmental signals into plant root development. *Annals of Botany*. 2013;**112**:1655-1665
- [15] Fahad S, Hussain S, Bano A, Saud S, Hassan S, Shan D, Khan FA, Khan F, Chen YT, Wu C, Tabassum MA, Chun MX, Afzal M, Jan A, Jan MT, Huang JL. Potential role of phytohormones and plant growth-promoting rhizobacteria in abiotic stresses: Consequences for changing environment. *Environmental Science and Pollution Research*. 2015;**22**:4907-4921
- [16] Egamberdieva D. Alleviation of salt stress by plant growth regulators and IAA producing bacteria in wheat. *Acta Physiologiae Plantarum*. 2009;**31**:861-864
- [17] Javid MG, Sorooshzadeh A, Moradi F, Sanavy SA, Allahdadi I. The role of phytohormones in alleviating salt stress in crop plants. *Australian Journal of Crop Science*. 2011;**5**:726-734
- [18] Fahad S, Hussain S, Matloob A, Khan FA, Khaliq A, Saud S, Hassan S, Shan D, Khan F, Ullah N, Faiq M, Khan MR, Tareen AK, Khan A, Ullah A, Ullah N, Huang JL. Phytohormones and plant responses to salinity stress: A review. *Plant Growth Regulation*. 2015;**75**:391-404
- [19] Kang NY, Cho C, Kim NY, Kim J. Cytokininreceptor-dependent and receptor-independent pathways in the dehydration response of *Arabidopsis thaliana*. *Journal of Plant Physiology*. 2012;**169**:1382-1391
- [20] O'Brien JA, Benkova E. Cytokinin cross-talking during biotic and abiotic stress responses. *Frontiers in Plant Science*. 2013;**4**:451. DOI: 10.3389/fpls.2013.00451
- [21] Nishiyama R, Watanabe Y, Fujita Y, Tien LD, Kojima M, Werner T, Vankova R, Yamaguchi-Shinozaki K, Shinozaki K, Kakimoto T, Sakakibara H, Schmuelling T, Lam-Son PT. Analysis of cytokinin mutants and regulation of cytokinin metabolic genes reveals important regulatory roles of cytokinins in drought, salt and abscisic acid responses, and abscisic acid biosynthesis. *Plant Cell*. 2011;**23**:2169-2183
- [22] Zalabák D, Pospíšilová H, Šmehilová M, Mrízová K, Frébort I, Galuszka P. Genetic engineering of cytokinin metabolism: Prospective way to improve agricultural traits of crop plants. *Biotechnology Advances*. 2013;**31**:97-117
- [23] Pospíšilová J. Participation of phytohormones in the stomatal regulation of gas exchange during water stress. *Biologia Plantarum*. 2003;**46**:491-506
- [24] Gamalero E, Glick BR. Ethylene and abiotic stress tolerance in plants. In: Ahmed P, Prasad MNV, editors. *Environmental Adaptations and Stress Tolerance of Plants in the Era of Climate Change*. New York: Springer; 2012. pp. 395-412
- [25] Groen SC, Whiteman NK. The evolution of ethylene signaling in plant chemical ecology. *Journal of Chemical Ecology*. 2014;**40**:700-716

- [26] Shi Y, Tian S, Hou L, Huang X, Zhang X, Guo H, Yang S. Ethylene signaling negatively regulates freezing tolerance by repressing expression of *CBF* and Type-A *ARR* genes in *Arabidopsis*. *Plant Cell*. 2012;**24**:2578-2595
- [27] Sponsel VM, Hedden P. Gibberellin, biosynthesis and inactivation. In: Davies PJ, editor. *Plant Hormones Biosynthesis, Signal Transduction, Action!* Dordrecht: Springer; 2004. pp. 63-94
- [28] Yamaguchi S. Gibberellin metabolism and its regulation. *Annual Review of Plant Physiology*. 2008;**59**:225-251
- [29] Colebrook EH, Thomas SG, Phillips AL, Hedden P. The role of gibberellin signaling in plant responses to abiotic stress. *Journal of Experimental Biology*. 2014;**217**:67-75
- [30] Munteanu V, Gordeev V, Martea R, Duca M. Effect of gibberellin cross talk with other phytohormones on cellular growth and mitosis to endoreduplication transition. *International Journal of Advanced Research in Biological Sciences*. 2014;**1**(6):136-153
- [31] Vardhini BV, Anuradha S, Rao SS. Brassinosteroids: A great potential to improve crop productivity. *Indian Journal of Plant Physiology*. 2006;**11**:1-12
- [32] Bajguz A, Hayat S. Effects of brassinosteroids on the plant responses to environmental stresses. *Plant Physiology and Biochemistry*. 2009;**47**:1-8
- [33] Fahad S, Nie L, Chen Y, Wu C, Xiong D, Saud S, Hongyan L, Cui K, Huang J. Crop plant hormones and environmental stress. *Sustainable Agriculture Reviews*. 2015;**15**:371-400
- [34] Pauwels L, Inze D, Goossens A. Jasmonate-inducible gene: What does it mean? *Trends in Plant Science*. 2009;**14**:87-91. DOI: 10.1016/j.tplants.2008.11.005
- [35] Yoon J, Hamayun M, Lee SK, Lee IJ. Methyl jasmonate alleviated salinity stress in soybean. *Journal of Crop Science and Biotechnology*. 2009;**12**:63-68
- [36] Wang Y, Mopper S, Hasentein KH. Effects of salinity on endogenous ABA, IAA, JA, and SA in *Iris hexagona*. *Journal of Chemical Ecology*. 2001;**27**:327-342
- [37] Yan Z, Chen J, Li X. Methyl jasmonate as modulator of Cd toxicity in *Capsicum frutescens* var. *fasciculatum* seedlings. *Ecotoxicology and Environmental Safety*. 2013;**98**:203-209
- [38] Maksymiec W, Wojcik M, Krupa Z. Variation in oxidative stress and photochemical activity in *Arabidopsis thaliana* leaves subjected to cadmium and excess copper in the presence or absence of jasmonate and ascorbate. *Chemosphere*. 2007;**66**:421-427
- [39] Miura K, Tada Y. Regulation of water, salinity, and cold stress responses by salicylic acid. *Frontiers in Plant Science*. 2014;**5**:4. DOI: 10.3389/fpls.2014.00004
- [40] Rivas-San VM, Plasencia J. Salicylic acid beyond defense: Its role in plant growth and development. *Journal of Experimental Botany*. 2011;**62**:3321-3338
- [41] Jumali SS, Said IM, Ismail I, Zainal Z. Genes induced by high concentration of salicylic acid in *Mitragyna speciosa*. *Australian Journal of Crop Science*. 2011;**5**:296-303

- [42] Koltai H, Beveridge CA. Strigolactones and the coordinated development of shoot and root. In: Baluska F, editor. Long-Distance Systemic Signaling and Communication in Plants. Berlin: Springer; 2013. pp. 189-204
- [43] Ruyter-Spira C, Kohlen W, Charnikhova T, van Zeijl A, van Bezouwen L, de Ruijter N, Cardoso C, Lopez-Raez JA, Matusova R, Bours R, Verstappen F, Bouwmeester H. Physiological effects of the synthetic strigolactone analog GR24 on root system architecture in *Arabidopsis*: Another belowground role for strigolactones? *Plant Physiology*. 2011;**155**:721-734
- [44] Foo E, Davies NW. Strigolactones promote nodulation in pea. *Planta*. 2011;**234**:1073-1081
- [45] Vurro M, Yoneyama K. Strigolactones-intriguing biologically active compounds: Perspectives for deciphering their biological role and for proposing practical application. *Pest Management Science*. 2012;**68**:664-668
- [46] Kohli A, Sreenivasulu N, Lakshmanan P, Kumar PP. The phytohormone crosstalk paradigm takes center stage in understanding how plants respond to abiotic stresses. *Plant Cell Reports*. 2013;**32**:945-957
- [47] Depuydt S, Hardtke CS. Hormone signalling crosstalk in plant growth regulation. *Current Biology*. 2011;**21**:R365-R373
- [48] Consentino L, Lambert S, Martino C, Jourdan N, Bouchet PE, Witczak J, Castello P, El-Esawi M, Corbineau F, d'Harlingue A, Ahmad M. Blue-light dependent reactive oxygen species formation by *Arabidopsis* cryptochrome may define a novel evolutionarily conserved signaling mechanism. *New Phytologist*. 2015;**206**:1450-1462
- [49] El-Esawi MA. Genetic diversity and evolution of *Brassica* genetic resources: From morphology to novel genomic technologies—A review. *Plant Genetic Resources*. C. 2016:1-12. DOI:10.1017/S1479262116000058
- [50] El-Esawi MA. Micropropagation technology and its applications for crop improvement. In: Anis M, Ahmad N, editors. *Plant Tissue Culture: Propagation, Conservation and Crop Improvement*. Springer Singapore; 2016b. pp. 523-545
- [51] El-Esawi MA. Nonzygotic embryogenesis for plant development. In: Anis M, Ahmad N, editors. *Plant Tissue Culture: Propagation, Conservation and Crop Improvement*. Springer Singapore; 2016. pp. 583-598
- [52] El-Esawi MA. Somatic hybridization and microspore culture in *Brassica* improvement. In: Anis M, Ahmad N, editors. *Plant Tissue Culture: Propagation, Conservation and Crop Improvement*. Springer Singapore; 2016. pp. 599-609
- [53] El-Esawi MA, Sammour R. Karyological and phylogenetic studies in the genus *Lactuca* L. (Asteraceae). *Cytologia*. 2014;**79**:269-275
- [54] El-Esawi M, Glascoe A, Engle D, Ritz T, Link J, Ahmad M. Cellular metabolites modulate *in vivo* signaling of *Arabidopsis* cryptochrome-1. *Plant Signaling & Behavior*. 2015;**10**:e1063758

- [55] El-Esawi MA, Germaine K, Bourke P, Malone R. Genetic diversity and population structure of *Brassica oleracea* germplasm in Ireland using SSR markers. *Comptes Rendus Biologies*. 2016;**339**:133-140
- [56] El-Esawi MA, Germaine K, Bourke P, Malone R. AFLP analysis of genetic diversity and phylogenetic relationships of *Brassica oleracea* in Ireland. *Comptes Rendus Biologies*. 2016;**339**:163-170
- [57] Jourdan N, Martino C, El-Esawi M, Witzak J, Bouchet PE, d'Harlingue A, Ahmad M. Blue-light dependent ROS formation by *Arabidopsis* cryptochrome-2 may contribute towards its signaling role. *Plant Signaling & Behavior*. 2015;**10**:e1042647

