

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,500

Open access books available

118,000

International authors and editors

130M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Introductory Chapter: Overview on Echinococcosis

Tonay Inceboz

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.70583>

1. Overview on echinococcosis

Echinococcosis is a zoonotic helminth disease. First, description goes back to 1684 when Francesco Redi has first described the scolex. Then, in 1700s, Philip Jacob Hartmann has defined in an adult form. Since intercontinental travelling is very common, it has been an important health problem in the last decades. Although it has been known for many years, it keeps its peculiarity [1–3].

Echinococcus belongs to Cestoda (class) and the Taeniidae (family). There are four most common species: *Echinococcus granulosus*, *Echinococcus multilocularis*, *Echinococcus vogeli* and *Echinococcus oligarthrus*. The most common one is *E. granulosus*, causing cystic echinococcosis (CE), whereas *E. vogeli* is the rarest species. *E. multilocularis* is also rare; however, it is the most violent species causing alveolar echinococcosis (AE). *E. vogeli* and *E. oligarthrus* cause polycystic echinococcosis (PE). There are other species of *Echinococcus*; *Echinococcus canadensis*, *Echinococcus equinus*, *Echinococcus orteppi* and *Echinococcus shiquicus*. There are also many different genotypes of each species; *E. granulosus* has 10 genotypes (G1–G10), whereas *E. multilocularis* has two genotypes (M1–M2) [4, 5].

Echinococcosis is very widely distributed in all over the world; CE is found in North Pole, Asia, Europe, Africa, Australia and South America. AE is found in Alpine and sub-Arctic or Arctic regions, including Canada, the United States and Central and Northern Europe, China and Central Asia. PE is found in Central and South America [4, 6, 7].

Humans are not definitive hosts; however, two forms of echinococcosis important in humans are CE and AE. Although most people with the disease are asymptomatic, CE causes slowly growing cysts in the liver, lungs and other organs that can be undiagnosed for many years. AE, however, poses a much greater risk than CE, causing parasitic tumors in many organs and can be fatal, if left untreated [4, 6].

The diagnosis of echinococcosis is based on clinical findings, pathology, imaging and serology. There are numerous diagnostic methods for lab diagnosis of echinococcosis. The diagnosis of echinococcosis is based on clinical findings, imaging (radiology, ultrasonography, computed axial tomography, magnetic resonance imaging) [8, 9] and serology tests such as indirect hemagglutination (IHA) [10], enzyme-linked immunosorbent assay (ELISA), IgG-ELISA [11–13] or IgE-ELISA [14], immunoblotting (IB), Western blot 7 kDa and/or 18 kDa for CE and 28 kDa for AE [15] and *E. multilocularis* for diagnosis of Em2plus-ELISA [16], Em18 [17, 18], Em70 and Em90 [19]. On the other hand, various techniques such as random amplification of polymorphic DNA (PCR-RAPD), restriction fragment length polymorphism (PCR-RFLP) [20, 21] and multiplex PCR for a quick identification were used to determine genetic variations [22, 23].

Therapy for echinococcosis depends on the size, location and symptoms of the cysts and the overall health of the patient. There are medical and surgical approaches; however, medical approach—basically albendazole—is a neoadjuvant and adjuvant therapy. It reduces the risk of recurrent disease by the inactivation of the protoscolices. Surgery is the principle treatment modality [24, 25].

Author details

Tonay Inceboz

Address all correspondence to: tonay.inceboz@gmail.com

Medical Faculty, Department of Parasitology, Dokuz Eylul University, Turkey

References

- [1] Redi F. Osservazioni intorno agli animal viventi che si trovano negli animali viventi. Firenze: Piero Matini; 1684. p. 244
- [2] David I. Grove. A History of Human Helminthology. Wallingford, Oxon: CAB International, 1990. p. viii + 848. ISBN: 0-85198-689-7. DOI:10.1017/S0007087400027965
- [3] Craig PS, McManus DP, Lightowers MW, Chabalgoity JA, Garcia HH, Gavidia CM, Schantz PM. Prevention and control of cystic echinococcosis. Lancet Infectious Diseases. 2007;7(6):385-394. DOI: 10.1016/S1473-3099(07)70134-2
- [4] World Health Organization. About Echinococcosis. Available from: http://www.who.int/echinococcosis/about_disease/en/
- [5] Alvarez Rojas CA, Romig T, Lightowers MW. *Echinococcus granulosus* sensu lato genotypes infecting humans—Review of current knowledge. International Journal for Parasitology. 2014;44:9-18. DOI: 10.1016/j.ijpara.2013.08.008

- [6] Sachar S, Goyal S, Sangwan S. Uncommon locations and presentations of hydatid cyst. *Annals of Medical and Health Sciences Research*. 2014;**204**:447-452. DOI: 10.4103/2141-9248.133476
- [7] Hotez PJ, Gurwith M. Europe's neglected infections of poverty. *International Journal of Infectious Diseases*. 2011;**15**:e611-e619. DOI: 10.1016/j.ijid.2011.05.006
- [8] Moro P. Clinical Manifestations and Diagnosis of Echinococcosis. UpToDate. Last updated 2017. Available from: <http://www.uptodate.com/contents/clinical-manifestations-and-diagnosis-of-echinococcosis>
- [9] Inceboz T, Goktay Y, Sagol O, Korkmaz M, Uner A. Mesenteric Doppler ultrasonography findings of *Echinococcus multilocularis* infection: An experimental study. *Türkiye Parazitoloji Dergisi*. 2009;**33**(2):151-154 PubMed PMID: 19598092
- [10] Auer H, Stöckl C, Suhendra S, Schneider R. Sensitivity and specificity of new commercial tests for the detection of specific *Echinococcus* antibodies. *Wiener Klinische Wochenschrift*. 2009;**121**:37e41. DOI: 10.1007/s00508-009-1233-4
- [11] Zhang W, McManus DP. Recent advances in the immunology and diagnosis of echinococcosis. *FEMS Immunology and Medical Microbiology*. 2006;**47**:24e41. DOI: 10.1111/j.1574-695X.2006.00060.x
- [12] Carmena D, Benito A, Eraso E. Antigens for the immunodiagnosis of *Echinococcus granulosus* infection: An update. *Acta Tropica*. 2006;**98**:74e86. DOI: 10.1016/j.actatropica.2006.02.002
- [13] Sarkari B, Rezaei Z. Immunodiagnosis of human hydatid disease: Where do we stand? *World Journal of Methodology*. 2015;**26**;5(4):185-195. DOI: 10.5662/wjm.v5.i4.185
- [14] Lawn SD, Bligh J, Craig PS, Chiodini PL. Human cystic echinococcosis: Evaluation of post-treatment serologic follow-up by IgG subclass antibody detection. *The American Journal of Tropical Medicine and Hygiene*. 2004;**70**:329e335 PMID: 15031526
- [15] Liance M, Janin V, Bresson-Hadni S, Vuitton AD, Houin R, Piarroux R. Immunodiagnosis of echinococcus infections: Confirmatory testing and species differentiation by a new commercial western blot. *Journal of Clinical Microbiology*. 2000;**38**(10):3718-3721 PMID: 11015390
- [16] Gottstein B, Jacquier P, Bresson-Hadni S, Eckert J. Improved primary immunodiagnosis of alveolar echinococcosis in humans by an enzyme-linked Immunosorbent assay using the Em2Plus antigen. *Journal of Clinical Microbiology*. 1993;**31**(2):373-376 PMID: 8432825
- [17] Bart JM, Piarroux M, Sako Y, Grenouillet Bresson-Hadni S, Piarroux R, Ito A. Comparison of several commercial serologic kits and Em18 serology for detection of human alveolar echinococcosis. *Diagnostic Microbiology and Infectious Disease*. 2007;**59**:93-95. DOI: 10.1016/j.diagmicrobio.2007.03.018

- [18] Ito A, Xiao N, Liance M, Sato MO, Sako Y, Mamuti W, Ishikawa Y, Nakao M, Yamasaki H, Nakaya K, Bardonnnet K, Bresson-Hadni S, Vuitton DA. Evaluation of an enzyme-linked immunosorbent assay (ELISA) with affinity-purified Em18 and an ELISA with recombinant Em18 for differential diagnosis of alveolar echinococcosis: Results of a blind test. *Journal of Clinical Microbiology*. 2002;**40**:4161e4165 PMID: 12409391
- [19] Korkmaz M, Inceboz T, Celebi F, Babaoglu A, Uner A. Use of two sensitive and specific immunoblot markers, Em70 and Em90, for diagnosis of alveolar echinococcosis. *Journal of Clinical Microbiology*. 2004;**42**:3350e3352. DOI: 10.1128/JCM.42.7.3350-3352.2004
- [20] Bowles J, McManus DP. Rapid discrimination of echinococcus species and strains using a polymerase chain reaction-based RFLP method. *Molecular and Biochemical Parasitology*. 1993;**57**:231e239 PMID: 8094539
- [21] Xiao N, Nakao M, Qiu J, Budke CM, Giraudoux P, Craig PS, Ito A. Dual infection of animal hosts with different *Echinococcus* species in the eastern Qinghai-Tibet plateau region of China. *The American Journal of Tropical Medicine and Hygiene*. 2006;**75**:292e294 PMID: 16896135
- [22] Boubaker G, Macchiaroli N, Prada L, Cucher MA, Rosenzvit MC, Ziadinov I, Deplazes P, Saarma U, Babba H, Gottstein B, Spiliotis M. A multiplex PCR for the simultaneous detection and genotyping of the *Echinococcus granulosus* complex. *PLoS Neglected Tropical Diseases*. 2013;**7**:e2017. DOI: 10.1371/journal.pntd.0002017
- [23] Can H, Inceboz T, Caner A, Atalay Şahar E, Karakavuk M, Döşkaya M, Çelebi F, Değirmenci Döşkaya A, Gülçe İz S, Gürüz Y, Korkmaz M. Detection of *Echinococcus granulosus* and *Echinococcus multilocularis* in cyst samples using a novel single tube multiplex real-time polymerase chain reaction. *Mikrobiyoloji Bülteni*. 2016;**50**(2):266-277 Turkish PubMed PMID: 27175499
- [24] McManus DP, Gray DJ, Zhang W, Yang Y. Diagnosis, treatment, and management of echinococcosis. *British Medical Journal*. 2012;**344**:e3866. PMID: 22689886. DOI: 10.1136/bmj.e3866
- [25] Kern P. Clinical features and treatment of alveolar echinococcosis. *Current Opinion in Infectious Diseases*. 2010;**23**:505-512. DOI: 10.1097/QCO.0b013e32833d7516