

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,700

Open access books available

121,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# Physiological Features of Red Currant Adaptation to Drought and High Air Temperatures

*Panfilova Olga Vitalevna, Golyaeva Olga Dmitrievna,  
Knyazev Sergey Dmitrievich and Kalinina Olga Vitalevna*

## Abstract

An important requirement for varieties is adaptation to growing conditions. The main indicators of water regime, photosynthesis, and productivity of representatives of different species of *Ribesia* (Berl.) Jancz. subgenus to drought and high temperatures have been studied. Quantitative and qualitative changes of the photosynthetic apparatus are the response to drought and high temperatures. The ratio of chlorophylls to carotenoids is considered to be one of the indicators of adaptability. The total water potential in red currant leaves depends on shoot growth, leaf age, berry formation, variety, and weather conditions, and it is not the main indication of drought resistance. The ratio of bound and free water and water-holding capacity of the leaves is considered to be a determining sign of resistance to hyperthermia. Red currant genotypes do not possess high indicators of heat resistance. The prospects of using physiological rapid diagnostic methods in breeding for adaptability to destructive factors of the growing season are shown. The representatives of *Ribes petraeum* Wulf. (“Hollandische Rote”) and *R. multiflorum* Kit. (1426-21-80) have high levels of drought resistance, making them highly potential for wider growing in (semi)arid agroecological condition.

**Keywords:** *Ribesia* (Berl.) Jancz. subgenus, genotypes, drought resistance, heat hardiness, water regime

## 1. Introduction


The ecological factors of the environment play an important role in the resistance of fruit plants, their productivity, and crop quality [1, 2]. Global climate change is now threatening. Air and soil summer droughts reduce the quality and quantity of horticultural products. Weather anomalies lead to an imbalance of the protective mechanisms of fruit and berry crops and adversely affect the physiological processes occurring in plants [3, 4]. Recently, the task to obtain new high-yielding, precocious genotypes with an amplitude of adaptation to different conditions, with a high content of nutrients and biologically active substances in the fruit, is faced by scientists from different countries [5]. The search of express methods of plant diagnostics to destructive factors of the environment is a priority direction. It is known that the temperature increase causes morphological and anatomical, physiological, and biochemical changes, which affect the growth and development of plants and can lead to large economic losses

[6, 7]. The study of morphological features, structure of photosynthetic apparatus, and water exchange of plants in connection with the area of growth is the main condition for solving fundamental and applied problems in the biology of the culture. The use of physiological and biochemical rapid methods of diagnostics of plant resistance to adverse weather and climatic factors can significantly optimize the long breeding process, minimize crop losses, and obtain genotypes resistant to the destructive effects of climatic anomalies [8–10]. Plant organisms have different mechanisms of adaptation to stressors [9, 11]. Xeromorphic structure of a plant leaf, changes in a pigment complex, and water balance are important diagnostic signs of drought resistance and heat resistance [12–14]. Issues of adaptation of berry crops to drought and high temperatures are poorly studied. Red currant is one of the valuable berry crops due to the high content of vitamins, microelements, sugars, and organic acids. It is valued as a source of healthy nutrition [15–18]. Vitamin and the healing properties of the berries of this culture are also preserved in processed products [19]. Introduction is considered to be an important link in the distribution and production of new red currant genotypes. The success of the introduction is determined by the nature of the interaction of hereditary biological characteristics of plants with specific environmental conditions [18]. Red currants belong to the *Ribes* L. genus and *Ribesia* (Berl.) Jancz. subgenus. As a culture, it was developed on the basis of four species, i.e., *Ribes vulgare* Lam., *Ribes petraeum* Wulf., *Ribes multiflorum* Kit., and *Ribes rubrum* L., and their hybrids [20]. The world assortment of the *Ribes* L. genus includes more than 200 varieties; however, the genetic resources of the *Ribesia* (Berti.) Jancz. subgenus are poorly studied, since there are a number of wild species that exceed the existing varieties by a number of economic and biological characteristics [21].


## 2. Anatomical, morphological, and physiological changes of a red currant leaf due to drought and high temperature

The study of the assimilation apparatus in red currants gives a complete picture of the plasticity of the genotype in relation to the ecological factors of the environment [11, 22–24]. The investigation of the anatomical and morphological features of red currant leaf structure showed that some morphological features, such as shape and venation, depend on the biological characteristics of the variety and the anatomical structure is more influenced by the growing conditions. During the periods with insufficient water supply, the leaf surface area decreased in all studied currant samples compared to optimal conditions. A positive relationship between leaf area and hydrothermal coefficient ( $r = +0.99$ ) and a negative relationship between leaf area and ambient temperature ( $r = -0.97$ ) were identified [10]. Rezanova [14], Patzukova [25], Tokhtar [18], and Panfilova [2] studied the anatomical structure of currant leaves and described the structure of the stomatal apparatus and the conduction system of *Ribes rubrum* L. and *Ribes American* L. According to anatomical studies, the red currant leaf had a mesomorphic structure characteristic of the *Ribesia* (Berl.) Jancz. subgenus. Spongy parenchyma prevailed over palisade parenchyma, epidermal cells were large enough, and on the upper side of the leaf, they were larger than on the lower one (**Figure 1**) [10]. It is shown that stomata are formed on the lower side of the leaf rarely and unevenly, at the level of the epidermis. There are varietal species differences in the size and number of stomata per  $1 \text{ mm}^2$  of leaf area, in the length of the closing cells, and in their shape. The size of stomata and the degree of their openness depend on the temperature and humidity of the air. During the drought, the degree of stomata opening decreases sharply.

The index of leaf mesostructure is labile in red currant genotypes of different ecological and genetic origins, including *Ribes petraeum* Wulf. (“Hollandische Rote”



**Figure 1.**  
 Anatomical structure of a red currant leaf (cross sectional view,  $\times 40$ ), *R. vulgare* Lam. Bar represents 10  $\mu\text{m}$ .



**Figure 2.**  
 The average size ( $\pm$ S.E.) of the cells of adaxial epidermis of a leaf in red currant varieties and selected genotypes of the Russian Research Institute of Fruit Crop Breeding, Orel [10].

and 1518-37-14), *Ribes vulgare* Lam. (“Jonkheer Van Tets,” “Niva” and “Rosa”), and *Ribes multiflorum* Kit. (“Dana,” 1426-21-80 and 1432-29-98), depending on the genotypes and phase of plant development.

High temperature and drought had different effects on cell size of adaxial epidermis and leaf mesostructure. In drought periods (2012, 2013) at the stages of active growth of shoots and formation of berries, the main cells of the adaxial epidermis in “Hollandische Rote” (*Ribes petraeum* Wulf) and 1426-21-80 (*Ribes multiflorum* Kit.) under the action of high temperatures (up to  $+31.2 \dots +28.6^\circ\text{C}$  in May and  $+32.2 \dots +31.5^\circ\text{C}$  in July) were somewhat stretched due to the decrease in turgor of the cells. The remaining samples showed cell compression in the tangent direction (parallel to the stem surface) (**Figure 2**).

Increased temperature and drought lead to the growth of parenchymal cells and increase of the overall thickness of the leaf (**Table 1**). The growth of mesophyll cells occurs mainly due to the increase in the volume of air-bearing cavities of the spongy parenchyma, which contributes to the improvement of gas exchange between the leaf and the environment (**Figure 3**).


In some previous studies [2, 26, 27], it was confirmed that significantly greater growth of spongy parenchyma cells and leaf blade thickness in dry periods were found in most samples of *Ribesia* (Berl.) Jancz. (with the exception of “Niva”). The largest increase in the thickness of the leaf was noted in “Hollandische Rote” and 1426-21-80. The authors consider these changes as a manifestation of high

Variety name/code (B)	Year (A)								
	2011			2012			2013		
	PP	SP	TLT	PP	SP	TLT	PP	SP	TLT
1426-21-80	10.29	13.62	26.94	12.00	20.00	37.30	9.75	15.80	29.91
“Hollandische Rote”	9.56	15.24	27.70	11.25	18.45	35.35	9.55	16.95	31.40
“Niva”	9.28	12.65	24.63	10.15	15.35	29.60	10.35	15.30	29.10
“Dana”	8.10	12.00	22.70	10.05	16.30	29.92	9.10	14.85	27.70
1432-29-98	7.90	12.35	23.25	11.65	17.55	33.65	10.15	15.10	29.10
1518-37-14	7.70	14.68	25.38	9.15	18.15	30.80	8.55	16.90	29.15
“Jonkheer Van Tets”	7.30	10.10	20.33	10.50	15.90	30.05	9.00	12.30	24.80
“Rosa”	6.80	11.23	20.53	7.95	13.85	25.60	8.10	13.80	25.10

Notes: PP—palisade parenchyma,  $\mu\text{m}$ ; SP—sponge parenchyma,  $\mu\text{m}$ ; TLT—total leaf thickness,  $\mu\text{m}$ .  $LSD_{05}$  for palisade parenchyma  $A = 1.57$ ,  $B = 2.56$ , and  $AB = 4.42$ ;  $LSD_{05}$  for sponge parenchyma  $A = 1.92$ ,  $B = 3.14$ , and  $AB = 5.43$ ;  $LSD_{05}$  for total leaf thickness  $A = 1.39$ ,  $B = 2.27$ , and  $AB = 3.93$ .

**Table 1.**

Leaf mesostructural parameters in red currant varieties from the Russian Research Institute of Fruit Crop Breeding, Orel, in 2011–2013 vegetation period [10].

**Figure 3.**

The leaf mesophyll cells in a red currant exposed to drought ((cross sectional view,  $\times 40$ ), *R. multiflorum* Kit.). Bar represents  $10 \mu\text{m}$ .

adaptability of the anatomical structure of the leaf to stressors (high temperature) during the growing season (**Table 1**).

Knowledge of the anatomical structure of the leaves helps to fully reveal the biological characteristics of the variety and species as well as the flow of the most important physiological processes [28]. In the process of evolution, many currant species have acquired a high potential of photosynthetic productivity, and, for the most part, in the real environment, this potential is not fully used [29, 30].

Drought is one of the reasons for the decrease in the intensity of photosynthesis, respiration, and changes in the hormonal status of plants. Violation of hormonal metabolism significantly increases the process of natural fall of the ovaries of red currants, as well as the appearance of necrotic points on the leaves and a decrease in the number of laid generative buds; leads to a decrease in yield, during both current and subsequent years; and also reduces the stability of plants in winter [31–33].

The process of photosynthesis largely depends on the quantitative and qualitative composition of pigments. Changes in the pigment complex can be considered as important mechanisms of culture adaptation to environmental conditions. The impact of high temperatures and low soil moisture leads to a decrease in the concentration of chlorophyll [22, 34–36]. Studies of different red currant species have confirmed the impact of air and soil drought on the work of the assimilation apparatus.

In the dry period (in the Orel region it was 2012), there was a decrease in the content of chlorophyll *a* and chlorophylls *a+b* in the leaves of all red currant genotypes in comparison with the favorable weather conditions (2011) (Figure 4).

A sharp (maximum) decrease in the number of pigments was observed in all derivatives of *R. vulgare* Lam. in 2012; the minimum decrease was observed in the representatives of *R. petraeum* Wulf. and *R. multiflorum* Kit. A slight decrease in drought in 2013 contributed to a slight increase in pigment content.

Drought at high temperatures stimulates the formation of carotenoids in the cell, which is consistent with the literature data on the protective function of this group of pigments under stress [5, 37, 38]. In the representatives of *Ribesia* (Berl.) Jancz. subgenus, the content of carotenoids in drought conditions increased by 2–3 times compared to a favorable period. A positive correlation in red currants was found between the carotenoid content and temperature ( $r = +0.77$ ) [2, 10]. An indirect indicator of drought resistance and high temperature is the ratio of the sum of chlorophylls to carotenoids [23, 39, 40]. In stressful conditions, the high value of this indicator was observed in the variety “Hollandische Rote” (the value of the coefficient 5.14) and the selected form 1426-21-80 (the value of the coefficient 5.51), which may indicate the stability of these samples to drought and high summer temperatures. Hydrothermal regime affects productivity by affecting the functional state of plants, which was confirmed by high correlations between the amount of chlorophyll *a* and yield ( $r = +0.85$ ) and the sum of chlorophylls and yield ( $r = +0.78$ ) [10].




Figure 4. Content of chlorophyll *a* and sum of chlorophylls *a+b* in red currant varieties and selected genotypes of the Russian Research Institute of Fruit Crop Breeding, Orel [10].

### 3. Features of water regime in red currants (drought resistance, heat hardiness)

Changes in the hydration of tissues and their water balance under the influence of adverse environmental conditions are reflected in the level, direction, and relationship of physiological processes that determine the formation of the crop and its quality. The ability to retain and economically consume water in arid conditions is a protective and adaptive reaction of resistant plants, which is due to a number of internal factors [33, 43].

It is established that the increase of water-holding capacity of leaves in extreme conditions of water supply allows stable red currant plants to regulate water metabolism [33, 41]. High water-holding capacity against the background of low water potential (61–67%) was observed in the varieties “Jonkheer Van Tets,” “Hollandische Weisse,” and “Englische Grosse Weisse.” It is noted that the water potential of red currant leaves is not the main indicator of plant resistance to drought, because it depends on a number of factors (weather and climatic conditions, genotype, leaf age, shoot growth, and berry formation). The maximum reduction of water potential of leaf tissues in representatives of the subgenus *Ribesia* (Berl.) Jancz. occurs during the formation of berries, as most of the water is used to form the ovary [10, 42].

The resistance of red currants to hyperthermia is determined by the high content of bound water and water-holding capacity of leaves, low water deficit, and intensity of transpiration [43–45]. High values of the coefficient of bound water to free water and minimal water losses were observed in the variety “Hollandische Rote” and selected form 1426-21-80. The genotypes of these samples of *R. petraeum* Wulf. and *R. multiflorum* Kit. species exhibit higher adaptability to high temperatures against the background of soil moisture deficit (**Table 2**) [10]. In terms of transpiration intensity, red currant samples exhibit different mechanisms of adaptation to drought: either reduce the level of transpiration intensity (plants save water) or the level of transpiration intensity remains high (drought level for these genotypes is not critical) [43].

The relationship between physiological characteristics was revealed in representatives of different red currant species, which was confirmed by the coefficient of pair correlation between the amount of chlorophylls and water loss ( $r = -1.00$ ), the amount of chlorophylls, and the amount of free water ( $p = +0.98$ ). The content of pigments, fractional composition of water, and water-holding capacity of the leaf were associated with the development of water deficit, which depended on the air temperature ( $r = +0.84$ ). The relationship between water regime indicators and meteorological features of the growing season was described using multiple regression coefficients (**Table 3**) [10].

In the study of plant resistance to abiotic environmental factors, their ability to withstand high air temperatures is important. Abnormally high temperatures, regardless of the dehydrating effect of dried soil and air, lead to disruption of water metabolism in plants and damage to membranes and proteins of the cell [46–50]. In this regard, an important feature of the variety is heat resistance [51]. Heat resistance largely depends on the duration of high temperatures and their absolute value. Temperature limits are specific to an individual genotype. In most cases, fruit plants begin to suffer when the temperature rises to 35–40°C. At these and higher temperatures, normal physiological functions are inhibited, and at temperatures around 50°C protoplasm, coagulation and cell death may occur. The exceeding the optimum temperature level leads to partial or global denaturation of proteins. In heat-resistant genotypes in the lipid complex, the saturated fatty acids predominate, and their appearance is a consequence of adaptation to this damaging factor. The mechanism of increasing the heat resistance is interconnected with the genetic apparatus of the cell and is aimed at stabilizing the membrane lipids in the direction of reducing their poly saturation. Under the action of elevated temperatures in the cells, the synthesis of stress proteins is induced [52]. Heat resistance is also associated with a certain

Variety name/code (B)	Year (A)							
	2011		2012		2013		Average	
	BW/FW	WL, %	BW/FW	WL, %	BW/FW	WL, %	BW/FW	WL, %
“Hollandische Rote”	1.33	35.82	2.09	20.36	2.03	21.11	1.82	25.76
1426-21-80	1.31	36.16	2.41	20.62	2.09	19.37	1.94	25.38
1432-29-98	1.12	30.17	1.42	25.01	1.20	26.98	1.25	27.39
“Jonkheer Van Tets”	1.01	38.10	1.31	21.85	1.04	27.40	1.12	29.12
1518-37-14	0.96	40.91	1.06	27.62	1.08	26.66	1.03	31.73
“Niva”	0.82	42.56	1.71	24.84	1.50	25.58	1.34	30.99
“Dana”	0.70	40.97	1.76	25.51	1.66	25.44	1.37	30.64
“Rosa”	0.63	39.75	0.98	26.82	0.87	27.85	0.83	31.47
LSD <sub>05</sub>	0.08	2.06	0.11	3.03	0.14	1.86		

Notes: BW/FW—ratio of bound and free water; WL—water loss in 24 hours; % of water content, LSD<sub>05</sub> for BW/FW—A, 2.18; B, 4.59; and AB, 7.95; and LSD<sub>05</sub> for WL—A, 0.13; B, 0.21; and AB, 0.37.

**Table 2.**  
 The main indicators of the water regime in red currant varieties from the Russian Research Institute of Fruit Crop Breeding, Orel, in 2011–2013 vegetation period [10].

Indicator	Regression ratio
Hydration	0.90×F; r = 0.22
Free water fraction	0.84×F; r = 0.33
Water loss	−0.88×F; r = 0.30

Notes: F—the values of hydrothermal coefficient; and r—the standard error of the experiment.

**Table 3.**  
 Correlation between physiological indicators and meteorological conditions in studied red currant varieties from the Russian Research Institute of Fruit Crop Breeding, Orel [10].

stage of plant development: young actively growing tissues are less stable than old ones. For berry plants, high temperatures are especially dangerous during flowering, because they cause sterility of flowers and the fall of ovaries. Plant organs differ in their heat resistance: shoots and buds are most stable, and the root system is less stable. The cambial tissue is most stable among tissues.

Russian and foreign researchers have achieved certain results in the study of water metabolism and physiological and biochemical parameters of resistance of fruit plants and developed methods for determining heat resistance. Water loss and the degree of water content recovery are informative indicators of heat resistance assessment [34, 53–64].

The effect of temperature “shock” +50°C on the red currant genotypes showed that this culture does not have high heat resistance. Leaves at a young age (less heat-resistant) are most severely damaged by high temperatures; the older the leaf, the higher its heat resistance [65]. The degree of water recovery in red currant leaves increases by the time of berry ripening. The leaves of red currant genotypes recover water better during berry ripening and are less susceptible to damage by temperature +50°C; they are more heat-resistant than the leaves during the active growth of shoots. One hundred percent recovery of water loss does not occur after exposure to temperature stress (+50°C) on the leaf. The most heat-resistant varieties are considered “Niva” (*R. vulgare* Lam.) and “Hollandische Rote” (*R. petraeum* Wulf.) [66].



Field assessment of red currant plants was carried out in Adygea after a long and gradually increasing heat in 1998. “Jonkheer Van Tets,” “Natali,” “Nenagliadnaya,” “Boulogne Blanche,” and *Ribes biebersteinii* Berl. showed resistance to heat (damage up to 2.5 points). The red currant samples of “Englische Grosse Weisse,” “Hollandische Weisse,” “Boulogne Rouge,” “Versailles Rouge,” “White Viksne,” “Hollandische Weisse,” and “London Market” dropped all the leaves, but the kidneys were still alive. Necrotic spots of different degrees appeared on the leaves of “Jonkheer Van Tets,” “Nenagliadnaya,” and “Boulogne Blanche” [33].

#### 4. Conclusion

Summarizing the above, it can be approved that the problem of drought resistance and heat resistance of varieties is currently particularly acute. The increase of variety stability now gets the great relevance, since the numerous experiments prove that only the use of the adapted varieties and the transition to adaptive, biological, and low-cost technologies of cultivation of fruit-growing production are possible [67]. Many issues related to the peculiarities of the water regime and physiological and biochemical processes remain unclear. In our work we have shown that berry crops, including red currants, have complex and little-known mechanisms of adaptation to drought and abnormally high temperatures. The prospects of using physiological methods in assessing the resistance of *Ribesia* (Berl.) Jancz samples to adverse weather factors have been proved. The obtained results were fully confirmed by field observations of stability of red currant varieties and selected forms under conditions of high temperature and moisture deficit. This makes it possible to use the considered methods as a rapid diagnostics and significantly intensify the selection for adaptability. As shown by the complex physiological assessment, the derivatives of different species of the *Ribesia* (Berl.) Jancz subgenus have a different ecological plasticity, which is explained by the difference of their genetic and ecological-geographical origins. The derivatives of different species of the *Ribes petraeum* Wulf. (“Hollandische Rote”) and *R. multiflorum* Kit. (1426-21-80) showed the greatest resistance to dry conditions of the vegetation period in the Central Chernozem Region that gives reason to use them in introduction. Low drought resistance was observed in the varieties and selected forms of *Ribes vulgare* Lam.

#### Author details

Panfilova Olga Vitalevna\*, Golyaeva Olga Dmitrievna, Knyazev Sergey Dmitrievich and Kalinina Olga Vitalevna  
Russian Research Institute of Fruit Crop Breeding, Orel, Russia

\*Address all correspondence to: [olga\\_280682@mail.ru](mailto:olga_280682@mail.ru)

#### IntechOpen

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

## References

- [1] Gudkovsky VA. Resistance of fruit and berry plants to stress factors and ways to improve it. In: Collection of Proceedings: Ways to Improve the Sustainability of Gardening; Michurinsk. 1998. pp. 17-29
- [2] Panfilova OV. Assessment of red currant adaptability to abiotic factors of the north-west of the Central-Chernozem region [thesis]. Orel: Orel State Agrarian University; 2014
- [3] Khaustovich IP. Improving the sustainability of horticulture in response to climate change in the Central-Chernozem Region [thesis]. Voronezh: Voronezh State Agrarian University named after K.D. Glinka; 2006
- [4] Meriño-Gergichevich C, Reyes-Díaz M, Guerrero J, Ondrasek G. Physiological and nutritional responses in two highbush blueberry cultivars exposed to deficiency and excess of boron. *Journal of Soil Science and Plant Nutrition*. 2017;17(2):307-318
- [5] Krivoruchko VP. Ecological and biological bases of increase of productivity of apple orchards of Northern Kyrgyzstan [thesis]. Bishkek: National Science Academy of the Kirghiz Republic; 1998
- [6] Beck EH, Fettig S, Knake C, Hartig K, Bhattarai T. Specific and unspecific responses of plants to cold and drought stress. *Journal of Biosciences*. 2007;32:501-510. DOI: 10.1007/s12038-007-0049-5
- [7] Hasanuzzaman M, Nahar K, Fujita M. Extreme temperature responses, oxidative stress and antioxidant defense in plants. In: Vahadati K, Leslie C, editors. *Abiotic Stress-Plant Response and Applications in Agriculture*. Rijeka, Croatia: Intech; 2013. DOI: 10.5772/54833
- [8] Wise RR, Olson AJ, Schrader SM, Sharkey TD. Electron transport is the functional limitation of photosynthesis in field-grown Pima cotton plants at high temperature. *Plant, Cell & Environment*. 2004;27(6):717-724. DOI: 10.1111/j.1365-3040.2004.01171.x
- [9] Neeru K, Kalpna B, Kadambot HMS, Harsh N. Food crops face rising temperatures: An overview of responses, adaptive mechanisms, and approaches to improve heat tolerance. *Cogent Food & Agriculture*. 2016;2(1):271-313. DOI: 10.1080/23311932.2015.1134380
- [10] Panfilova OV, Golyaeva OD. Physiological features of red currant varieties and selected seedling adaptation to drought and high temperature. *Agrobiology*. 2017;52(5):1056-1064. DOI: 10.15389/agrobiology.2017.5.1056rus
- [11] Davlatov SH, Ashurov AA, Baykova EV. Morphological and anatomical leaf structure of wild species of *Berberis* genus from Tajikistan. *Tomsk State University Journal*;323:348-350
- [12] Skhalyakho TV, Nenko NI, Kisileva GK. Assessment of drought resistance degree of grape varieties in the conditions of the Anapa-Taman zone by physiological, biochemical, anatomical and morphological parameters. *Fruit Growing and Viticulture of South Russia*. 2012;17:69-78
- [13] Rezanova TA, Sorokopudov VN. Some features of anatomical and morphological structure of *Ribes americanum* mill. In: Proc. Sci. Conf. *Biological Active Compounds of Natural Origin: Phytotherapy, Pharmaceutical Marketing, Pharmaceutical Technology, Botany*; Belgorod. 2008. pp. 133-135
- [14] Rezanova TA. Morphological, anatomical and ecological features of

- Ribes americanum* Mill. at introduction in the South of the Central Russian upland [thesis]. Saratov: Saratov State University named after N.G. Chernyshevsky; 2010
- [15] Määttä K, Kamal-Eldin A, Törrönen R. Phenolic compounds in berries of black, red, green, and white currants (*Ribes* sp.). *Antioxidants & Redox Signaling*. 2001;**3**:981-993. DOI: 10.1089/152308601317203521
- [16] Pantelidis GE, Vasilakakis M, Manganaris GA, Diamantidis G. Antioxidant capacity, phenol, anthocyanin and ascorbic acid contents in raspberries, blackberries, red currants, gooseberries and cornelian cherries. *Food Chemistry*. 2007;**102**:777-783. DOI: 10.1016/j.foodchem.2007.06.021
- [17] Skrede G, Martinsen BK, Wold AB, Birkeland SE, Aaby K. Variation in quality parameters between and within 14 Nordic tree fruit and berry species. *Acta Agriculturae Scandinavica Section B Soil and Plant Science*. 2011;**62**:193-208. DOI: 10.1080/09064710.2011.598543
- [18] Tokhtar LA. Biological features of red currants of *Ribes* (Berl.) Jancz. subgenus at introduction in the conditions of Belgorod region [thesis]. Belgorod: Belgorod State National Research University; 2011
- [19] Makarkina MA. Apple and red currant breeding for improvement of fruit chemical composition [thesis]. Bryansk: Bryansk State Agricultural Academy; 2010
- [20] Fedorovsky VD. *Ribes spicatum* Robson. (Taxonomy, Geography, Variability, Introduction); Kiev, 2001
- [21] Golyaeva OD. Red currant breeding results at the Russian Research Institute of Fruit Crop Breeding. In: Collection of Proceedings: The Current State of Currant and Gooseberry Crops; Michurinsk. 2007. pp. 32-35
- [22] Bacelar EA. Ecophysiological Responses of Olive (*Olea europaea* L.) to Restricted Water Availability: Limitations, Damages and Drought Resistance Mechanisms. Vila Real: Universidade De Trás-Os-Montes E Alto Douro; 2006
- [23] Kiseleva NS. Assessment of adaptive capacity of different pear genotypes according to morphoanatomic and physiological state of leaves. *Agricultural Biology*. 2009;**3**:34-38
- [24] Krokmal II, Pugacheva AY. Leaf anatomical features of species of *Hemerocallis* L. genus under the conditions of steppe zone of the Ukraine. *Journal Bulletin of Problems Biology and Medicine*. 2011;**1**(2):72-76
- [25] Patzukova NG. Biological features of *Ribes Alpinum* L. at introduction in Belgorod region [thesis]. Saratov: Saratov State University named after N.G. Chernyshevsky; 2010
- [26] Panfilova OV, Golyaeva OD. Currant adaptation to drought and anomalously high temperatures (overview). *Contemporary Horticulture*. 2015;**2**(14):88-98
- [27] Golyaeva OD, Panfilova OV. Adaptive red currant varieties of VNIISPK breeding. *Pomiculture and Small Fruits Culture in Russia*. 2014;**38**(1):96-100
- [28] Ondrasek G, Clode PL, Kilburn MR, Guagliardo P, Romić D, Rengel Z. Zinc and cadmium mapping in the apical shoot and hypocotyl tissues of radish by high-resolution secondary ion mass spectrometry (NanoSIMS) after short-term exposure to metal contamination. *International Journal of Environmental Research and Public Health*. 2019;**16**:373
- [29] Zhidekhuna TV. Photosynthetic potential of productivity of black

- currant varieties of VNIIS breeding. In: Proc. Sci. Conf.; 22-24 April 2013. Voronezh: Voronezh State Agrarian University; 2013. pp. 65-70
- [30] Zhidekhuna TV. Photosynthetic active radiation and its use by different black currant varieties. In: Proc. Sci. Conf. Contemporary Cultivars and Technologies for Intensive Orchards; 15-18 July 2013; Orel. Orel: VNIISPK; 2013. pp. 93-95
- [31] Shmadlak J. Flower development in pome and stone breeds. In: Kudryavtzeva RP, editor. Physiology of Fruit Plants; Moscow. 1983. pp. 265-276
- [32] Borovkov VV. Biochemical aspects of ripening and falling of black currant fruit [thesis]. Moscow: Moscow Agricultural Academy named after K.A. Tsvetkov; 1997
- [33] Semenova LG, Bzhetzeva NR. Features of Productivity of Black and Red Currants in Adygea; Maykop; 2003. 143 p
- [34] Kushnirenko MD, Pecherskaya SN. Physiology of Water Exchange and Drought Resistance of Plants; Kishinev, 1991.
- [35] Ananina VM, Zholkevich VN. Questions of plant resistance physiology. Agricultural Biology. 2003;1:120-124
- [36] Belous OG. Diagnostics of resistance of tea plants to stress factors. Vestnik of the Russian Agricultural Science. 2008;6:41-43
- [37] Kiseleva NS. Influence of environmental stress factors on the morpho-anatomical structure of pear leaf blade. In: Proc. Sci. Conf. Adaptive Potential and Quality of Products of Fruit Cultivars and Cultivar-Rootstock Combinations; 24-7 July 2012; Orel. Orel: VNIISPK; 2012. pp. 115-122
- [38] Khvostov DS, Kirtbaya EK. Adaptation potential of strawberries in the south of Russia. Horticulture and Viticulture. 1999;4:16-17
- [39] Nenko NI, Ilyina IA, Skhalyakh TV. Adaptation resistance of grapes in summer. In: Proc. Sci. Conf. High-Precision Technologies of Production, Storage and Processing of Grapes. Krasnodar: SKZNIISViV; 2010. pp. 50-59
- [40] Golyaeva OD, Panfilova OV. Adaptive red currant cultivars of VNIISPK breeding. Pomiculture and Small Fruits Culture in Russia. 2014;38(1):96-100
- [41] Bzhetzeva NR. Study of currant gene pool for using in production and breeding in Adygea [thesis]. Krasnodar: Russian Research Institute of Rice; 2002
- [42] Golyeva OD, Petrov AV. Drought resistance of red currant varieties. In: Proc. Sci. Conf. Breeding and Variety Propagation of Orchard Crops; 19-22 June 2007; Orel. Orel: VNIISPK; 2007. pp. 64-74
- [43] Panfilova OV, Golyaeva OD. Effect of drought resistance on the water regime of red currants. Pomiculture and Small Fruits Culture in Russia. 2012;31(2):119-126
- [44] Machado S, Paulsen GM. Combined effects of drought and high temperature on water relations of wheat and sorghum. Plant and Soil. 2001;233(2):179-187. DOI: 10.1023/A:1010346601643
- [45] Mironova LN, Denisova SG, Zeynetdinova GS, Reut AA. To the assessment of the adaptive potential of decorative perennials. In: Vestnik of Voronezh State University, Series: Geography; Geoecology. Vol. 1. 2011. pp. 157-159
- [46] Saratova NA. Some regulatory mechanisms of plant adaptation to drought and high temperatures. In: Physiology of Plant Drought Resistance; Moscow. 1971. pp. 28-29
- [47] Altergot VF, Mordkovich SS, Ignatiev LA. Evaluation principles of drought and heat resistance of plants. In:

Methods of Plant Resistance Evaluation to Unfavorable Conditions of the Environment; Leningrad. 1976. pp. 6-17

[48] Chentzova ES. Prospects of introduction and use of some species and clones of persimmon in the Kuban zone of fruit growing [thesis]. Krasnodar: Kuban State Agrarian University; 2008

[49] Pyanikov VT. Resistance of new cherry cultivars and hybrid seedlings to drought and high temperatures in the south of the central Chernozem zone. In: Proc. Sci. Conf. State and Prospects of Fruit Breeding and Variety Propagation; 12-15 July 2005; Orel. Orel: VNIISPK; 2005. pp. 403-407

[50] Nenko NI, Kiseleva GK, Sergeev YI. Resistance of apple plants to abiotic stress of winter and summer periods in the coenosis of different design. In: Proc. Sci. Conf. Biological Basis for Gardening and Vegetable Growing; 22-25 September 2010; Michurinsk. 2010. pp. 240-244

[51] Savelieva IN. Genetic and selection features and economical value of columnar apple cultivars and forms [thesis]. Ramon: Russian Research Institute of Beet and Sugar named after A.L. Mazlumov; 2012

[52] Fedulov YP, Kotlyarov VV, Dotzenko KA. Plant Resistance to Unfavorable Factors of the Environment. Krasnodar: Kuban State Agrarian University; 2015. 64 p

[53] Kushnirenko MD, Goncharova EA, Bondar EM. Methods of Studying Water Metabolism and Drought Resistance of Fruit Plants; Kishinev; 1970. 79 p

[54] Genkel PA. Physiology of heat and drought resistance of plants. Moscow; 1982. 280 p

[55] Zhuchenko AA. Plant Breeding Adaptive System (Ecological and

Genetic Principles). Vol. 1. Moscow: RUDN University; 2001. 780 p

[56] Leonchenko VG, Evseyeva RP. Development of methods for predicting the resistance of fruit plants to salinization. In book of Ecological Aspects of Intensification of Agricultural Production. Vol. 1. Penza; 2002. pp. 46-48

[57] Leonchenko VG, Evseyeva RP, Zhanova EV. Preliminary selection of promising genotypes of fruit plants for ecological and genetic resistance and biochemical value of fruits. Methodical Recommendations. Michurinsk; 2007. 72 p

[58] Saveliev NI, Savelieva NN, Yushkov AN. Promising Scab Immune Apple Cultivars. Michurinsk; 2009. 128 p

[59] Yushkov AN, Chivilev VV, Borzykh NV, Abyzov VV, Khozhanov A.V. Resistance of fruit and berry plants to dehydration and overheating. In: Potential and Quality of Products of Fruit Cultivars and Cultivar-Rootstock Combinations Conference on scientifically and practical; 24-27 July 2012; Orel: VNISPK; 2012. p. 287-291

[60] Panfilova OV, Bogomolova NI, Goljaeva OD, Ozherel'eva ZE. Valutazione della resistenza e bacche di calore colture nella regione centrale delle terre nere della Russia. Italian Science Review. 2013;9:86

[61] Panfilova OV, Ozherelieva ZE, Golyaeva OD. Comparative assessment of the adaptive potential of red currant genotypes (*Ribes rubrum* L.) to abiotic factors of the vegetation period. Plant Varieties Studying and Protection. 2014;1(22):24-27

[62] Matysik J, Alai Bhalu B, Mohanty P. Molecular mechanisms of quenching of reactive oxygen species by proline under stress in plants. Current Science. 2002;82:525-532

[63] Mittler R. Oxidative stress, antioxidants, and stress tolerance. *Trends in Plant Science*. 2002;7:405-409. DOI: 10.1016/S1360-1385(02)02312-9

[64] Sung DY, Kaplan F, Lee KJ, Guy CL. Acquired tolerance to temperature extremes. *Trends in Plant Science*. 2003;8(3):179-187. DOI: 10.1016/S1360-1385(03)00047-5

[65] Zubov AA, Luk'yanchuk IV. Field evaluation of drought resistance of strawberry plants on wilting and drying of leaves. *Agricultural Biology*. 2004;1:116-118

[66] Panfilova OV, Ozherelieva ZE, Golyaeva OD. Assessment of resistance to extremely high temperatures of vegetation period of red currant genotypes. In: *Proc. Sci. Conf*; 15-18 July 2013; Orel. Orel: VNIISPK; 2013. pp. 171-172

[67] Dragavtzeva IA, Savin IY, Ovechkin SV. *Resource Potential of Land in Krasnodar Region for Fruit Crop Cultivation*; Krasnodar; 2005. 136 p