

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,400

Open access books available

117,000

International authors and editors

130M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# Ion Channels and Their Regulation in Vascular Smooth Muscle

*Arsalan U. Syed, Thanhmai Le, Manuel F. Navedo  
and Madeline Nieves-Cintrón*

## Abstract

Vascular smooth muscle excitability is exquisitely regulated by different ion channels that control membrane potential ( $E_m$ ) and the magnitude of intracellular calcium inside the cell to induce muscle relaxation or contraction, which significantly influences the microcirculation. Among them, various members of the  $K^+$  channel family, voltage-gated  $Ca^{2+}$  channels, and transient receptor potential (TRP) channels are fundamental for control of vascular smooth muscle excitability. These ion channels exist in complex with numerous signaling molecules and binding partners that modulate their function and, in doing so, impact vascular smooth muscle excitability. In this book chapter, we will review our current understanding of some of these ion channels and binding partners in vascular smooth muscle and discuss how their regulation is critical for proper control of (micro)vascular function.

**Keywords:** ion channels, signal transduction, vascular tone, vascular smooth muscle, microcirculation

## 1. Introduction

Vascular smooth muscle cells wrapping around small resistance arteries and arterioles are crucial for vascular reactivity [1]. These cells enable dynamic, moment-to-moment control of vessel diameter and pressure-induced contraction (e.g., vascular tone). This control is central to autoregulation of resistance vessels, maintenance of vessel caliber independently of changes in blood pressure, and proper perfusion to meet the metabolic demands of a given tissue.

To regulate arterial diameter, vascular smooth muscle receives and integrates many inputs, including changes in intraluminal pressure, vasoconstrictor and vasodilatory signals from endothelial cells lining the inner arterial wall, and nerve terminals innervating the vessels [2]. These inputs regulate vascular smooth muscle excitability, at least in part, by modulating the activity of a number of ion channels to control membrane potential ( $E_m$ ) and the magnitude of intracellular  $Ca^{2+}$  concentration ( $[Ca^{2+}]_i$ ) [1]. Among the many ion channels, transient receptor potential (TRP) channels, voltage-gated ( $K_V$ ),  $Ca^{2+}$ -activated ( $BK_{Ca}$ ) and inward rectifier ( $K_{ir}$ )  $K^+$  channels, and voltage-gated  $Ca^{2+}$  channels (VGCC) are fundamental in transducing mechanical force, establishing  $E_m$ , and regulating  $[Ca^{2+}]_i$  [1]. Mechanisms for the regulation of vascular smooth muscle ion channels, including those mentioned above, involve agonist-independent and agonist-dependent activation of  $G_q$  and/or  $G_s$  protein-coupled receptors ( $G_x$ PCR) [3]. The optimal

activation and regulation of these mechanisms are often dependent on the formation of macromolecular complexes driven by scaffold proteins to target and compartmentalize proteins/signaling events to specific regions and substrates [4].


In this chapter, we provide a brief overview of our current understanding of the role of different subunits of TRP channels, several  $K^+$  channel subtypes, as well as key VGCCs in control of vascular smooth muscle excitability and (micro)vascular function. We will also discuss their regulation by signaling pathways and macromolecular complexes. Note that many more studies than those cited here can be found in the scientific literature. Comprehensive studies exploring additional aspects of vascular ion channel regulation, vascular smooth muscle excitability, and (micro)vascular function in health and disease can be found in recent reviews [1, 5–9].

## 2. TRP channels

The TRP channel superfamily is composed of 28 members divided into 6 subfamilies based on their molecular and biophysical properties [10]. Functional TRP channels are composed of four subunits, each with six membrane-spanning helices that can exist in a homomeric or heteromeric form. Vascular smooth muscle cells express a number of these channels, including members of the TRPC, TRPV, TRPM, and TRPP subfamilies [7, 9]. The function of TRP channels in vascular smooth muscle ranges from regulating contractility to modulating the proliferative state of the cells. In this section, we focused our discussion on the functional role and regulation of specific TRP channels in vascular smooth muscle excitability.

### 2.1 Functional role of vascular smooth muscle TRP channels

The function of different TRP channels in vascular smooth muscle has been unmasked using conventional and innovative molecular, pharmacological, and genetic approaches. These approaches revealed distinctive roles of specific TRP channel subfamilies in modulating vascular smooth muscle excitability and vascular reactivity. For instance, studies have found that TRPC3 channels are not essential for pressure-induced constriction. Yet, TRPC3 channels play a key role in receptor-mediated vasoconstriction of resistance arteries upon activation of various  $G_q$ PCRs, including purinergic receptors, endothelin ( $ET_A$ ) receptors, and angiotensin II ( $AT_1$ ) receptors [7, 11, 12]. The mechanisms for TRPC3 channel activation involve direct coupling of the channel with  $IP_3$  receptors located in the sarcoplasmic reticulum (SR) in a process that does not require SR  $Ca^{2+}$  release [13]. TRPC5 channels, in association with TRPC1 channels, have been shown to contribute to store-operated  $Ca^{2+}$  entry in vascular smooth muscle from arterioles [14], which may modulate cell excitability. Expression of TRPP channels in vascular smooth muscle contributes to stretch-activated cation currents, which causes vascular smooth muscle membrane depolarization. The activity of these channels has been associated with stretch-dependent regulation of vascular tone in cerebral arteries and control of systemic blood pressure [15, 16]. The nonselective cation channel TRPC6 is involved in  $Ca^{2+}$  mobilization leading to vascular smooth muscle contraction [17]. More recently, these channels have been shown to be part of a mechanosensation complex [18]. In this model (see **Figure 1**), stretch is “sensed” by  $AT_1$  receptors. These receptors then induce the activation of TRPC6 channels to bolster  $Ca^{2+}$  release from the SR, which triggers TRPM4 channel activity and vascular smooth muscle contraction. This model reveals an exquisite and finely orchestrated macromolecular complex for control of stretch-induced contraction.



**Figure 1.**

*Regulation of vascular smooth muscle excitability by TRP channels. Diverse TRP channels contribute to vascular smooth muscle relaxation or contraction. Ca<sup>2+</sup> influx via TRPV4 channels is associated with an increase in the frequency of Ca<sup>2+</sup> release through ryanodine receptors (RyR) located in the sarcoplasmic reticulum (e.g., Ca<sup>2+</sup> sparks). The production of Ca<sup>2+</sup> sparks triggers the activity of adjacent BK<sub>Ca</sub> channels at the surface membrane to promote K<sup>+</sup> efflux, membrane potential hyperpolarization, and vascular smooth muscle relaxation [18]. TRPV4 channel activity can be stimulated by activation of AT<sub>1</sub> receptors (AT<sub>1</sub>R) by angiotensin II (angII) via an AKAP5-mediated PKC signaling pathway [20, 25]. This AT<sub>1</sub>R/AKAP5/PKC/TRPV4 axis may serve as a negative feedback mechanism to offset stimulation of L-type Ca<sup>2+</sup> channel Ca<sub>v</sub>1.2 (LTCCs) leading to contraction in response to angII. AT<sub>1</sub>R, independent of angII, has also been implicated in a mechanosensitive pathway that activates TRPC6 channels to boost cytosolic Ca<sup>2+</sup> via stimulation of IP<sub>3</sub> receptors (IP<sub>3</sub>R). This increase in cytosolic Ca<sup>2+</sup> promotes the activity of TRPM4 channels leading to vascular smooth muscle membrane depolarization and subsequent contraction in response to stretch [17]. The orange wavelike lines highlight stretch-sensing proteins. The model was generated by taking in consideration studies cited and described above. Na<sup>+</sup> = sodium; Ca<sup>2+</sup> = calcium; AKAP5 = A kinase anchoring protein 5; PKC = protein kinase C; DAG = diacylglycerol; IP<sub>3</sub> = inositol trisphosphate.*

Contrary to the contractile influences of the TRP channels described above, activation of TRPV4 channels is associated with vascular smooth muscle relaxation (**Figure 1**) [19]. The underlying mechanism involves the formation of a macromolecular complex between TRPV4 and BK<sub>Ca</sub> channels in the plasma membrane and ryanodine receptors (RyR) in the SR. This complex does not depend on direct protein-protein interactions between TRPV4 and BK<sub>Ca</sub> channels and RyR but requires their close physical proximity. The close association between these proteins facilitates Ca<sup>2+</sup> influx via TRPV4 channels that stimulates RyR activity resulting in the generation of Ca<sup>2+</sup> sparks. The ensuing Ca<sup>2+</sup> sparks activate BK<sub>Ca</sub> channels leading to vascular smooth muscle membrane potential hyperpolarization and relaxation [20]. Because of the high Ca<sup>2+</sup> permeability of TRPV4 channels in vascular smooth muscle [21], they could potentially activate adjacent BK<sub>Ca</sub> channels directly to regulate cell excitability and vascular reactivity. Examination of this exciting possibility is an excellent opportunity for further research. Moreover, additional work is needed to comprehensively define the role of TRP channels in vascular smooth muscle in the microcirculation during physiological and pathological conditions.

## 2.2 Regulation of TRP channels in vascular smooth muscle

It has been well documented that agonist-dependent activation of G<sub>q</sub>PCRs/protein kinase C (PKC) and G<sub>s</sub>PCRs/protein kinase A (PKA) signaling regulates TRP channels [10]. Protein kinase G (PKG) modulation of TRP channels has also been reported. However, how these signaling pathways control TRP channel activity in vascular smooth muscle, as well as the underlying consequences in vascular

reactivity are not well understood. A limited number of studies have shown that PKG inhibits TRPC3 channels and that this may contribute to nitric oxide (NO)-mediated relaxation, but the mechanisms require further examination [22]. PKA signaling stimulates the activity of several members of the TRPV channel subfamily while inhibiting TRPC5 and TRPC6 channels in vascular smooth muscle [7]. PKA-mediated inhibition of TRPC6 channels in response to agonist stimulation was shown to be dependent on phosphorylation of the TRPC6 subunit at threonine 69, which resulted in a reduction in angiotensin II (angII)-induced vasoconstriction [23], thus revealing a comprehensive mechanism for the regulation of TRP channels and the underlying effects in vascular reactivity.


PKC activity inhibits TRPC3 channels and activates TRPM4 and TRPV4 channels in smooth muscle [7, 9, 21, 24]. TRPM4 activation by PKC proceeds, at least in part, by stimulating channel trafficking and membrane translocation via a mechanism requiring PKC $\delta$  [25]. This PKC $\delta$ -dependent anterograde TRPM4 trafficking is functionally relevant as it promotes vascular smooth muscle contraction. PKC-dependent regulation of TRPV4 channels has been suggested to be critical for counteracting the vasoconstriction stimulated by angII [21]. Intriguingly, TRPV4 channels are found in complex with the scaffold A kinase anchoring protein 5 (AKAP5 = human AKAP79 and murine AKAP150) in vascular smooth muscle (**Figure 1**) [21, 26]. This scaffold protein provides a platform for targeting and compartmentalization of signaling molecules (e.g., PKA, PKC, protein phosphatase 2B = PP2B) to specific substrates (e.g., ion channels) [4]. With a suggested distance between them of ~200 nm [26], optimal AKAP5-anchored PKC modulation of TRPV4 activity is highly dependent on the distance between the targeted kinase and the ion channel. This tight regulation of TRPV4 activity by AKAP5 may be essential for vascular smooth muscle excitability given the high Ca<sup>2+</sup> permeability of these channels, as mentioned previously. Mechanisms regulating TRPP channels by PKA and/or PKC in vascular smooth muscle are currently unclear. The studies discussed above reveal the complex functional role of TRP channels in vascular smooth muscle and how their regulation alters vascular function and highlights unique opportunities for further research. For instance, it will be important to define the role of the AKAP5/PKC/PKA complex and its association with other TRP channels in regulating vascular smooth muscle excitability and vascular reactivity. It also remains to be determined whether dynamic trafficking of other TRP channels and the involvement of the AKAP5/PKC/PKA complex in this process also play a role in fine-tuning vascular reactivity.

### 3. K<sup>+</sup> channels

The activity of K<sup>+</sup> channels determines vascular smooth muscle membrane potential and is therefore key regulators of vascular tone [1]. By setting and controlling the membrane potential, these channels influence the levels of [Ca<sup>2+</sup>]<sub>i</sub> and therefore, vascular smooth muscle contraction. Intriguingly, a subset of K<sup>+</sup> channels has also been linked to regulation of vascular smooth muscle proliferation (see recent review on this topic in [27, 28]). Vascular smooth muscle cells express a wide variety of isoforms from several classes of K<sup>+</sup> channels, including K<sub>v</sub>, BK<sub>Ca</sub>, and K<sub>ir</sub> channels (see **Figure 2**). In the following sections, we will describe the expression, function, and regulation of these K<sup>+</sup> channels and their control of vascular function.

#### 3.1 K<sub>v</sub> channels

A number of K<sub>v</sub> channels isoforms are expressed in vascular smooth muscle, including members of the K<sub>v</sub>1 (K<sub>v</sub>1.1, K<sub>v</sub>1.2, K<sub>v</sub>1.3, K<sub>v</sub>1.5, K<sub>v</sub>1.6), K<sub>v</sub>2 (K<sub>v</sub>2.1), and



**Figure 2.**

Regulation of vascular smooth muscle excitability by  $K^+$  channels. Vascular smooth muscle cells express a number of  $K^+$  channels that, upon activation, provide negative feedback regulation of membrane potential and vascular tone.  $K^+$  channel activation results in efflux of  $K^+$  ions that hyperpolarize the membrane potential ( $\Delta E_m$ ; denoted by dotted black lines) leading to a reduction in the open probability of LTCC  $Ca_{v1.2}$  and TTCC  $Ca_{v3.1}$  that results in a decrease in global  $[Ca^{2+}]_i$  leading to vascular smooth muscle relaxation [32, 95]. As mentioned in **Figure 1**, activation of ryanodine receptors (RyR)  $K^+$  channels can be distinctively regulated by vasoactive agents acting through the  $G_s/AC/PAK$ ,  $NO/sGC/PKG$ , and  $G_q/PLC/PKC$  axes to either increase or decrease their activity. An example is the regulation of  $K_{v1.X}$  channels by  $\beta$  adrenergic signaling (e.g., through  $\beta_1$  adrenergic receptors) [44–46]. Here,  $K_{v1.X}$  channel regulation by  $\beta_1$  adrenergic signaling requires PKA targeting to the channel by postsynaptic density protein 95 (PSD95). Whether PSD95 as well as further scaffolding of proteins by AKAP5 is required for PKA regulation of  $K_{v1.X}$  and other  $K^+$  channels (e.g.,  $K_{v2.X}$ ,  $K_{v7.X}$ ,  $K_{ir2.X}$ ) as well as RyR is unclear (denoted by dotted red lines in the cartoon). The model was generated by taking in consideration studies cited and described above.  $K^+$  = potassium;  $Ca^{2+}$  = calcium; AKAP5 = A kinase anchoring protein 5;  $BK_{Ca}$  = large-conductance  $Ca^{2+}$  activated potassium channels; AC = adenylyl cyclase; PKA = protein kinase A; NO = nitric oxide; sGC = soluble guanylyl cyclase; PKG = protein kinase G; PLC = phospholipase C; PKC = protein kinase C.

$K_V 7$  ( $K_{V7.1-5}$ ), as well as several silent  $K_V$  subunits (e.g.,  $K_{V9.3}$ ) [1].  $K_V$  channels are formed by a homo- or heterotetrameric assembly of pore-forming  $\alpha$  subunits and regulatory  $\beta$  subunits. Key examples with functional relevance in vascular smooth muscle excitability from multiple vascular beds are  $K_V$  channels assembled by  $K_{V1.2}/K_{V1.5}$  subunits,  $K_{V2.1}$  subunits with  $K_{V9.3}$ , and  $K_{V1}/K_{V7}$  subunits. These subunit compositions confer functional and pharmacological diversity essential for fine-tuning vascular smooth muscle excitability and vascular reactivity [29–31]. The mechanisms for this involve activation of  $K_V$  channels by a depolarizing stimulus (e.g., stretch-induced depolarization) and their regulation by vasoactive agents [1, 32, 33]. The resultant  $K^+$  efflux hyperpolarizes the membrane potential of vascular smooth muscle leading to a decrease in the open probability of voltage-gated L-type  $Ca^{2+}$  channels  $Ca_{v1.2}$  (LTCCs), which contributes to decrease  $[Ca^{2+}]_i$  and relaxation (**Figure 2**). Conversely, their inhibition depolarizes the membrane potential, which will increase the open probability of LTCCs and lead to an increase in global  $[Ca^{2+}]_i$  and vascular smooth muscle contraction. This negative feedback regulation, together with other  $K^+$  channels, is essential for fine control of vascular smooth muscle excitability and vascular reactivity. However, in the context of the microcirculation, not much is known regarding the expression, functional composition, physiological role, and regulation of  $K_V$  channels in arteriolar vascular smooth muscle. Consideration of these issues is important as recent studies have implicated that impairment in  $K_V$  channel expression and/or function in the development of channelopathies are associated with small vessel

diseases [34]. Moreover, additional pathologies such as hypertension, metabolic disorders, and diabetic hyperglycemia impair (micro)vascular function, at least in part, by altering the expression/function of  $K_V$  channels, but mechanisms remain not fully understood (see review in [5]).

Many vasoactive agents modulate vascular function by acting on  $K_V$  channels expressed in vascular smooth muscle. For instance, agents that trigger activation of  $G_q$ PCRs, such as angII, phenylephrine, and endothelin 1, are known to stimulate vasoconstriction, at least in part, by decreasing the expression and/or function of  $K_V$  channels, particularly those of the  $K_{V1}$ ,  $K_{V2}$ , and  $K_{V7}$  subfamilies (**Figure 2**) [1, 35–37]. The effects of these vasoconstrictors are related to PKC-mediated phosphorylation and/or changes in surface expression of  $K_V$  subunits [36, 38–40], perhaps via engagement of different PKC isoforms [41]. This may contribute to selective control of  $K_V$  channel activity in response to different stimuli. In addition, increases in  $[Ca^{2+}]_i$  have also been associated with inhibition of  $K_V$  channels [42, 43]. Considering that activation of  $G_q$ PCRs also increases  $[Ca^{2+}]_i$ , these data suggest that both PKC activity and elevated  $[Ca^{2+}]_i$  could synergize to exacerbate  $K_V$  channel inhibition, which will result in membrane potential depolarization, activation of LTCCs, and vascular smooth muscle contraction.

In stark contrast to  $G_q$  signaling, activation of NO/PKG and  $G_s$ /PKA signaling is typically associated with stimulation of vascular  $K_V$  channels, including the members of the  $K_{V1}$  and  $K_{V7}$  subfamilies [1]. The functional consequence of this regulation is vascular smooth muscle relaxation. Phosphorylation of  $K_V$  channels by PKA is opposed by protein phosphatases such as PP2B, which will dephosphorylate the different subunits [44]. Whether  $G_s$ PCR/PKA/PP2B regulation of  $K_V$  channels requires scaffolding proteins that could target all components of the signaling complex within close proximity to the channels is not well understood. A recent series of studies have demonstrated that the scaffold protein postsynaptic density 95 (PSD95), which was thought to be a neuronal-specific protein, is expressed in vascular smooth muscle (**Figure 2**) [45, 46]. Intriguingly, PSD95 was found to be necessary for basal- and isoproterenol-induced PKA-mediated activation of  $K_{V1.X}$  channels that resulted in vascular smooth muscle relaxation [45–47]. This was due to the formation of a distinctive PSD95-mediated signaling complex involving the  $\beta_1$ -adrenergic receptor ( $\beta_1$  AR)-, PKA-, and  $K_{V1.2}$ -containing channels [45, 47]. Since PSD95 is associated with AKAP5 in neurons [48], the argument was made that the PSD95-AKAP5 complex may be essential for PKA targeting and regulation of  $K_{V1.2}$  function and that this will have an impact on vascular smooth muscle excitability and vascular reactivity [47]. However, additional studies have found that  $K^+$  currents produced by  $K_{V1.X}$  and  $K_{V2.X}$  subunits and  $BK_{Ca}$  channels are of similar magnitude in wild-type and AKAP5-depleted ( $AKAP5^{-/-}$ ) vascular smooth muscle cells [5, 49, 50]. These results suggest that, at least basally, AKAP5 is not necessary, and PSD95 may be sufficient for PKA-dependent regulation of  $K^+$  channels in these cells.  $\beta$ -Adrenergic stimulation has also been found to regulate  $K_{V7}$  channels leading to vasorelaxation [51], but whether a scaffold protein is mediating these effects is unknown and thus requires further examination. In addition, it is also unclear how  $K_{V2}$ -containing channels, which contribute about 70% of the  $K_V$  current in mouse cerebral and mesenteric vascular smooth muscle [49], are regulated by PKA signaling, presenting another area of further research.

### 3.2 $BK_{Ca}$ channels

$BK_{Ca}$  channels are abundantly expressed in vascular smooth muscle cells [52]. The pore-forming  $\alpha$  subunit ( $BK_{Ca}$   $\alpha 1.1$ ) assembles into tetramers to form a functional channel, but unlike TRP and  $K_V$  subunits, it contains seven transmembrane

domains and a heavily regulated long carboxyl terminal. Further regulation of BK<sub>Ca</sub> channel function is conferred by the association of the  $\alpha$  subunit with accessory  $\beta 1$  and LRRC26  $\gamma$  subunits, which increases the Ca<sup>2+</sup> sensitivity of the channel [52]. BK<sub>Ca</sub> channels are sensitive to voltage and Ca<sup>2+</sup>. The Ca<sup>2+</sup> source to activate BK<sub>Ca</sub> channels comes from the release of Ca<sup>2+</sup> from the SR via RyRs (**Figure 2**) [20]. This complex does not depend on direct protein-protein interactions between BK<sub>Ca</sub> channels and RyR but requires that both of these ion channels are in close physical proximity to each other. As mentioned above, data also have suggested that TRPV4 is responsible for triggering RyR activity [19], and thus, these channels have been proposed to form part of the same signaling complex with BK<sub>Ca</sub> and RyRs in vascular smooth muscle. Intriguingly, LTCCs have been found to have an indirect or “loose” coupling with RyRs that could enable their activation [53], perhaps by modulating the SR Ca<sup>2+</sup> concentration [54]. Additional components of the same or similar signaling complexes (see T-type Ca<sup>2+</sup> channels section) with divergent roles have been identified [1, 7, 13, 55], but an integrated model remains to be defined. Regardless, the activation of these channels should result in K<sup>+</sup> efflux that hyperpolarizes the plasma membrane of vascular smooth muscle cells leading to relaxation. The physiological influence of BK<sub>Ca</sub> channels in control of membrane potential and vascular tone, however, seems to be species-, stimulus-, and vessel-dependent, even when compared between different orders of the same vascular tree. Indeed, studies have reported a clear involvement of BK<sub>Ca</sub> channels in tonic negative feedback regulation of membrane potential and pressure-induced constriction, even in human resistance arteries, whereas others have failed to establish a relationship (exemplary studies in [56–62]). The differences may be associated with disparities in the BK<sub>Ca</sub> channel's Ca<sup>2+</sup> sensitivity and/or subunit expression levels and composition in the different vascular beds. A recent and broader discussion on this topic can be found in [1]. Thus, given the distinctive role of BK<sub>Ca</sub> channels in the vasculature, further research on how these channels are regulated to control vascular tone in different vascular beds during physiological and pathological conditions is warranted.

Generally, vasoactive agents that act via activation of the G<sub>s</sub>/adenylyl cyclase (AC)/PKA and NO/soluble guanylyl cyclase (sGC)/PKG axes potentiate BK<sub>Ca</sub> channel activity, whereas those acting through the G<sub>q</sub>/phospholipase C (PLC)/PKC axis inhibit the channel [1, 52]. Vasoactive agents may also regulate BK<sub>Ca</sub> channel activity indirectly by modulating the function of RyR in the SR or other ion channels (e.g., L-type Ca<sub>v</sub>1.2 and T-type Ca<sub>v</sub>3.2 channels) in the plasma membrane that are involved with direct activation of RyR or SR Ca<sup>2+</sup> refilling [54, 63, 64]. Although there is some evidence that scaffold proteins such as AKAP5 may help target signaling molecules to BK<sub>Ca</sub> channels [65], whether this indeed occurs in native vascular smooth muscle cells, as well as its functional relevance, is unclear. Considering the role of BK<sub>Ca</sub> channels in negative feedback regulation of vascular tone, the physiological implications of vasoactive agents acting through distinct G<sub>x</sub>PCRs on BK<sub>Ca</sub> channels will be either vascular smooth muscle relaxation or contraction.

### 3.3 K<sub>ir</sub> channels

As their name implies, K<sub>ir</sub> channels produce an inward current. This current is observed at a potential negative to the K<sup>+</sup> equilibrium potential that helps stabilize the resting E<sub>m</sub> [1, 66]. They also produce a small outward current at depolarizing potentials that serves as an electrical amplifier to magnify hyperpolarization. Inward rectification occurs due to voltage-dependent blockade of the channel by polyamines and Mg<sup>2+</sup>. K<sub>ir</sub> channels are regulated by lipids (e.g., phosphatidylinositol 4,5 biphosphate (PIP<sub>2</sub>) and cholesterol) [1, 66]. Intriguingly, a recent study found that cholesterol, but not PIP<sub>2</sub>, regulates K<sub>ir</sub> channel activity in the cerebral

vascular smooth muscle [67], suggesting that these channels may be distinctively modulated by lipids depending on their tissue distribution. A functional  $K_{ir}$  channel is formed when four pore-forming  $\alpha$  subunits, each containing two membrane-spanning domains, come together. Two main  $\alpha$  subunits (e.g.,  $K_{ir2.1}$  and  $K_{ir2.2}$ ) have been identified in vascular smooth muscle from multiple species [68–71]. Intriguingly, the expression of these subunits in a specific vascular bed may be species-dependent. Accordingly, although  $K_{ir}$  subunit expression and channel activity have been extensively reported in murine cerebral vascular smooth muscle [67–69, 71], minimal, if any,  $K_{ir}$  subunit expression and channel activity were found in the human cerebral vascular smooth muscle [72]. The functional implication of the activation of these channels in vascular smooth muscle is relaxation.  $K_{ir}$  channel activity can be modulated by vasoactive agents with those acting through the  $G_q$ /PLC/PKC axis, inducing channel inhibition, and those acting on the  $G_s$ /AC/PKA pathway, promoting channel activity [1]. The physiological relevance of these regulatory mechanisms on  $K_{ir}$  channels and their control of vascular function are less well understood and therefore are in need of further evaluation.

#### 4. Voltage-gated $Ca^{2+}$ channels

Vascular smooth muscle cells express several subtypes of VGCCs [9]. These channels have been shown to be important for vascular smooth muscle contraction, and some subtypes have been implicated in relaxation mechanisms. In this section, we will focus on the role of two key subtypes of VGCCs, namely, LTCCs and T-type  $Ca^{2+}$  channels (TTCCs), in regulation of vascular smooth muscle excitability.


##### 4.1 L-type $Ca^{2+}$ channel $Ca_v1.2$

The L-type  $Ca^{2+}$  channel  $Ca_v1.2$  (i.e., LTCCs) is essential for vascular smooth muscle contraction and vascular reactivity. Therefore, they play a key role in controlling blood flow and blood pressure [33, 73]. LTCCs are comprised of a pore-forming  $\alpha_{1c}$  subunit and auxiliary  $\beta$ ,  $\alpha_2\delta$ , and  $\gamma$  subunits that modulate channel function and trafficking [74]. The  $\alpha_{1c}$  subunit contains four homologous domains (I, II, III, IV). Each domain comprises of six membrane-spanning segments (S1–S6) with intracellular amino- and carboxyl termini, which contain many regions relevant for channel regulation and control of cell excitability. In vascular smooth muscle, expression of the  $\alpha_{1c}$  subunit is critical for pressure-induced constriction as evidenced by an absence of myogenic response after LTCC blockade and depletion of the  $Ca_v1.2$ - $\alpha_{1c}$  subunit in mice (e.g., SMAKO mouse) [33, 73, 75]. The auxiliary subunits  $\alpha_2$  and  $\delta$  are the product of the same gene that gets proteolytically cleaved after translation but remains connected by disulfide bonds, which give rise to the mature subunit. The  $\alpha_2\delta$  subunit has been linked to regulation of  $\alpha_{1c}$  subunit surface expression that controls  $Ca^{2+}$  influx in vascular smooth muscle and the level of myogenic constriction [76]. The  $\beta$  subunit, which remains cytoplasmic, also contributes to the  $\alpha_{1c}$  subunit surface expression and channel regulation and therefore can modulate vascular smooth muscle excitability in health and disease [77, 78]. Unlike the other subunits, the expression, regulation, and function of the  $\gamma$  subunit in vascular smooth muscle are unclear and likely the subject of further research.

LTCCs in vascular smooth muscle are distinctively regulated by the  $G_s$ /AC/PKA, NO/sGC/PKG, and  $G_q$ /PLC/PKC axes [9, 79]. Accordingly, the NO/sGC/PKG signaling axis has been shown to inhibit vascular LTCCs [80]. This has been associated with a reduction in  $[Ca^{2+}]_i$  that may be part of the vasodilatory mechanism underlying the activation of this pathway [79]. Receptor-mediated signaling via the

$G_q$ /PLC/PKC axis typically results in potentiation of LTCC activity [9, 79, 81]. The functional effects of this  $G_q$ /PLC/PKC-mediated activation of LTCCs are vascular smooth muscle contraction and an increase in vascular tone. Intriguingly, activation of the  $G_s$ /AC/PKA axis has been shown to inhibit, activate, or produce no effect on vascular LTCC activity (**Figure 3**) [79]. Irrespectively of this however, PKA signaling has been generally linked with vasodilation, thus raising questions about the functional relevance, if any, of this kinase in the regulation of vascular LTCCs. Intriguingly, recent studies revealed that elevations in extracellular D-glucose (HG) potentiate LTCC activity via a  $G_s$ /AC/PKA pathway in vascular smooth muscle [82–85]. This HG-induced PKA-dependent activation of LTCCs resulted in increased global  $[Ca^{2+}]_i$  and vasoconstriction, thus providing the first example of a PKA-dependent pathway underlying vascular smooth muscle contraction. Future studies should further examine the *in vivo* relevance of this pathway.

LTCC regulation by  $G_s$ /AC/PKA and  $G_q$ /PLC/PKC axes in vascular smooth muscle is mediated by AKAP5 (**Figure 3**) [85, 86]. The involvement of the scaffold in this regulation was initially speculated from total internal reflection fluorescence (TIRF) microscopy experiments that optically recorded the activity of single or clusters of LTCCs [87, 88]. From these experiments, it was clear that the activity and location of functional LTCCs were heterogeneous throughout the surface membrane of vascular smooth muscle cells [81, 87, 89]. Whereas some LTCCs showed stochastic activity with low  $Ca^{2+}$  flux and duration of events, others had persistent activity characterized by increased  $Ca^{2+}$  flux and events with prolonged open time that were produced by the opening of two or more channels [81, 87, 89–91]. The



**Figure 3.**

*Regulation of vascular smooth muscle excitability by voltage-gated  $Ca^{2+}$  channels.  $Ca^{2+}$  influx via L-type  $Ca^{2+}$  channel  $Ca_{v1.2}$  is essential for vascular smooth muscle contraction [33]. Their activity is regulated by  $G_s$ /PKA and  $G_q$ /PKC signaling pathways upon activation of a specific GPCR by a given stimulus (e.g., angII or HG). The association of  $Ca_{v1.2}$  with these signaling pathways is orchestrated by AKAP5 [86]. PKA (and perhaps PKC) augments L-type  $Ca^{2+}$  channel  $Ca_{v1.2}$  activity by increasing  $Ca_{v1.2}$  phosphorylation at serine 1928 ( $pS^{1928}$ ) [82, 85]. The phosphatase PP2B suppresses enhancement of L-type  $Ca^{2+}$  channel  $Ca_{v1.2}$  activity presumably by preventing/opposing channel phosphorylation in vascular smooth muscle cells (gray line). Two TTCC subtypes are expressed in vascular smooth muscle, with  $Ca_{v3.1}$  contributing to contractile mechanisms and  $Ca_{v3.2}$  forming a complex with RyR in the sarcoplasmic reticulum and  $BK_{Ca}$  channels in the surface membrane to foster relaxation [95, 100]. Although PKA has been shown to inhibit  $Ca_{v3.2}$ , whether this requires AKAP5 function is unclear (dotted red lines with perpendicular line at the end). It is also unclear whether AKAP5-anchored PKC and PKA regulate  $Ca_{v3.1}$  channel activity (dotted light and dark red lines with star near  $Ca_{v3.1}$ ). Finally, whether the GPCRs activated by angII and HG are targeted to specific complexes by AKAP5 is unclear (red dotted line with ?? symbols). The model was generated by taking in consideration studies cited and described above. GPCR = G-protein coupled receptors; angiotensin II = angII; high glucose = HG; AKAP5 = A kinase anchoring protein 5; protein kinase A = PKA; protein kinase C = PKC; protein phosphatase 2B; PP2B; phosphorylation at  $Ca_{v1.2}$  serine 1928 =  $pS^{1928}$ ;  $K^+$  = potassium;  $Ca^{2+}$  = calcium; ryanodine receptors = RyR.*

stochastic and persistent activity of LTCCs was modulated by membrane potential [92]. However, the occurrence of LTCCs with persistent activity is limited to specific regions of the surface membrane and has been demonstrated to be highly dependent on PKC activity and AKAP5 expression [81, 86]. The activity of phosphatases, such as PP2B, that are targeted to the channel by AKAP5, counteracts anchored kinase activity and restricts persistent LTCC activity (**Figure 3**) [89]. Accordingly, in vascular smooth muscle in which PKC is inhibited or cells from mice with genetically depleted PKC or AKAP5, the frequency of persistent LTCC activity is minimal [86, 87, 93]. In addition, PP2B inhibition stimulates persistent LTCC events in cells from wild type but not AKAP5<sup>-/-</sup> mice, suggesting that removing this “brake” facilitates kinase-mediated potentiation of channel activity [86, 89]. These results suggest an important role for AKAP5-anchored PKC and PP2B activity in modulating basal persistent LTCC activity. The physiological significance of these findings is underscored by data indicating that persistent LTCC events account for 50% of the total dihydropyridine-sensitive (e.g., LTCCs) Ca<sup>2+</sup> influx at physiological membrane potentials [92], which is critical for vascular smooth muscle contractility in health and disease [82, 84, 86, 93].

#### 4.2 T-type Ca<sup>2+</sup> channels

T-type Ca<sup>2+</sup> channels are formed by pore-forming  $\alpha_1$  subunits with similar topology as that of the LTCC  $\alpha_{1c}$  subunit, but with no known auxiliary subunits that modulate channel function [74]. TTCCs are activated at more hyperpolarized potentials and show similar conductance with Ca<sup>2+</sup> or Ba<sup>2+</sup> as charge carriers. Vascular smooth muscle cells express several TTCC  $\alpha_1$  subunits, including Ca<sub>v</sub>3.1 ( $\alpha_{1G}$ ) and Ca<sub>v</sub>3.2 ( $\alpha_{1H}$ ) [9, 94–98]. Intriguingly, Ca<sub>v</sub>3.1, which is found in murine vascular smooth muscle, seems to be replaced by Ca<sub>v</sub>3.3 ( $\alpha_{1I}$ ) in human cells [96], suggesting that expression of TTCC  $\alpha_1$  subunits is species-dependent. TTCCs have been shown to contribute to vascular smooth muscle excitability in several vascular beds from different species [9, 98]. However, rigorous analysis revealed that different Ca<sub>v</sub>3.X subunits may have very divergent physiological responses. For instance, whereas Ca<sub>v</sub>3.1 (Ca<sub>v</sub>3.3) mediates low-pressure-induced constriction, Ca<sub>v</sub>3.2 contributes to the negative feedback regulation of vascular tone by stimulating the RyR/BK<sub>Ca</sub> axis (**Figure 3**) [64, 95, 96]. TTCCs can also be regulated by signaling molecules. Indeed, the NO/PKG and AC/PKA axes both inhibit vascular TTCCs [99, 100], which may have key implications in vascular smooth muscle excitability. Whether these signaling molecules are organized and targeted by scaffold proteins such as AKAPs to areas near TTCCs to fine-tune their function is unclear and therefore the subject of future studies.

### 5. Conclusions

Vascular smooth muscle excitability is exquisitely controlled by a repertoire of ion channels, which in themselves, are regulated by several vasoactive agents. The precise regulation of ion channels in vascular smooth muscle cells is essential for the dynamic adjustment of vascular tone necessary to maintain adequate tissue perfusion and blood pressure. Here, we have provided a brief overview of our current knowledge of key ion channels and their regulation by receptor-mediated signaling pathways that are activated by various vasoactive agents to modulate vascular smooth muscle excitability and therefore vascular tone. We focused on several TRP channels, multiple K<sup>+</sup> channel subtypes, and various classes of VGCCs. We emphasized ion channel regulation by signaling pathways associated with the G<sub>s</sub>/AC/PKA,

NO/sGC/PKG, and G<sub>q</sub>/PLC/PKC axes given their important role in modulating vascular smooth muscle excitability. When possible, we identified key gaps in knowledge, even in areas that have been extensively studied, which are fertile ground for further research. Because of the importance of all the ion channels and signaling pathways discussed above on vascular control, understanding how they are affected during pathological conditions is essential for the development of rational therapies to treat (micro)vascular diseases.

## Acknowledgements

We thank the members of the Navedo Lab for critically reading early versions of the manuscript. This work was supported by NIH grants R01HL098200, R01HL121059 and R01HL149127 (to MFN), T32HL086350 (to AUS), and a UC Davis Academic Federation Innovative Development Award (to MN-C).

## Conflict of interest

None.

## Author details

Arsalan U. Syed, Thanhmai Le, Manuel F. Navedo and Madeline Nieves-Cintrón\*  
Department of Pharmacology, University of California Davis, Davis, CA,  
United States

\*Address all correspondence to: [mcnieves@ucdavis.edu](mailto:mcnieves@ucdavis.edu)

## IntechOpen

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

## References

- [1] Tykocki NR, Boerman EM, Jackson WF. Smooth muscle ion channels and regulation of vascular tone in resistance arteries and arterioles. *Comprehensive Physiology*. 2017;**7**:485-581
- [2] Bagher P, Segal SS. Regulation of blood flow in the microcirculation: Role of conducted vasodilation. *Acta Physiologica*. 2011;**202**:271-284
- [3] Mederos YSM, Storch U, Gudermann T. Mechanosensitive Gq/11 protein-coupled receptors mediate myogenic vasoconstriction. *Microcirculation*. 2016;**23**:621-625
- [4] Langeberg LK, Scott JD. Signalling scaffolds and local organization of cellular behaviour. *Nature Reviews Molecular Cell Biology*. 2015;**16**:232-244
- [5] Nieves-Cintrón M, Syed AU, Nystoriak MA, Navedo MF. Regulation of voltage-gated potassium channels in vascular smooth muscle during hypertension and metabolic disorders. *Microcirculation*. 2018;**25**:e12423
- [6] Amberg GC, Navedo MF. Calcium dynamics in vascular smooth muscle. *Microcirculation*. 2013;**20**:281-289
- [7] Earley S, Brayden JE. Transient receptor potential channels in the vasculature. *Physiological Reviews*. 2015;**95**:645-690
- [8] Narayanan D, Adebisi A, Jaggar JH. Inositol trisphosphate receptors in smooth muscle cells. *American Journal of Physiology. Heart and Circulatory Physiology*. 2012;**302**:H2190-H2210
- [9] Ghosh D, Syed AU, Prada MP, Nystoriak MA, Santana LF, Nieves-Cintrón M, et al. Calcium channels in vascular smooth muscle. *Advances in Pharmacology*. 2017;**78**:49-87
- [10] Nilius B, Szallasi A. Transient receptor potential channels as drug targets: From the science of basic research to the art of medicine. *Pharmacological Reviews*. 2014;**66**:676-814
- [11] Reading SA, Earley S, Waldron BJ, Welsh DG, Brayden JE. TRPC3 mediates pyrimidine receptor-induced depolarization of cerebral arteries. *American Journal of Physiology. Heart and Circulatory Physiology*. 2005;**288**:H2055-H2061
- [12] Xi Q, Adebisi A, Zhao G, Chapman KE, Waters CM, Hassid A, et al. IP3 constricts cerebral arteries via IP3 receptor-mediated TRPC3 channel activation and independently of sarcoplasmic reticulum Ca<sup>2+</sup> release. *Circulation Research*. 2008;**102**:1118-1126
- [13] Adebisi A, Zhao G, Narayanan D, Thomas-Gatewood CM, Bannister JP, Jaggar JH. Isoform-selective physical coupling of TRPC3 channels to IP3 receptors in smooth muscle cells regulates arterial contractility. *Circulation Research*. 2010;**106**:1603-1612
- [14] Xu SZ, Boulay G, Flemming R, Beech DJ. E3-targeted anti-TRPC5 antibody inhibits store-operated calcium entry in freshly isolated pial arterioles. *American Journal of Physiology. Heart and Circulatory Physiology*. 2006;**291**:H2653-H2659
- [15] Bulley S, Fernandez-Pena C, Hasan R, Leo MD, Muralidharan P, Mackay CE, et al. Arterial smooth muscle cell PKD2 (TRPP1) channels regulate systemic blood pressure. *eLife*. 2018;**7**:e42628
- [16] Narayanan D, Bulley S, Leo MD, Burris SK, Gabrick KS, Boop FA, et al. Smooth muscle cell transient

receptor potential polycystin-2 (TRPP2) channels contribute to the myogenic response in cerebral arteries. *The Journal of Physiology*. 2013;**591**:5031-5046

[17] Welsh DG, Morielli AD, Nelson MT, Brayden JE. Transient receptor potential channels regulate myogenic tone of resistance arteries. *Circulation Research*. 2002;**90**:248-250

[18] Gonzales AL, Yang Y, Sullivan MN, Sanders L, Dabertrand F, Hill-Eubanks DC, et al. A PLC gamma1-dependent, force-sensitive signaling network in the myogenic constriction of cerebral arteries. *Science Signaling*. 2014;**7**:ra49

[19] Earley S, Heppner TJ, Nelson MT, Brayden JE. TRPV4 forms a novel Ca<sup>2+</sup> signaling complex with ryanodine receptors and BKCa channels. *Circulation Research*. 2005;**97**:1270-1279

[20] Nelson MT, Cheng H, Rubart M, Santana LF, Bonev AD, Knot HJ, et al. Relaxation of arterial smooth muscle by calcium sparks. *Science*. 1995;**270**:633-637

[21] Mercado J, Baylie R, Navedo MF, Yuan C, Scott JD, Nelson MT, et al. Local control of TRPV4 channels by AKAP150-targeted PKC in arterial smooth muscle. *The Journal of General Physiology*. 2014;**143**:559-575

[22] Chen J, Crossland RF, Noorani MM, Marrelli SP. Inhibition of TRPC1/TRPC3 by PKG contributes to NO-mediated vasorelaxation. *American Journal of Physiology. Heart and Circulatory Physiology*. 2009;**297**:H417-H424

[23] Nishioka K, Nishida M, Ariyoshi M, Jian Z, Saiki S, Hirano M, et al. Cilostazol suppresses angiotensin II-induced vasoconstriction via protein kinase A-mediated phosphorylation of the transient receptor potential

canonical 6 channel. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 2011;**31**:2278-2286

[24] Albert AP, Large WA. Inhibitory regulation of constitutive transient receptor potential-like cation channels in rabbit ear artery myocytes. *The Journal of Physiology*. 2004;**560**:169-180

[25] Crnich R, Amberg GC, Leo MD, Gonzales AL, Tamkun MM, Jaggar JH, et al. Vasoconstriction resulting from dynamic membrane trafficking of TRPM4 in vascular smooth muscle cells. *American Journal of Physiology. Cell Physiology*. 2010;**299**:C682-C694

[26] Tajada S, Moreno CM, O'Dwyer S, Woods S, Sato D, Navedo MF, et al. Distance constraints on activation of TRPV4 channels by AKAP150-bound PKC $\alpha$  in arterial myocytes. *The Journal of General Physiology*. 2017;**149**:639-659

[27] Lopez-Lopez JR, Ciudad P, Perez-Garcia MT. Kv channels and vascular smooth muscle cell proliferation. *Microcirculation*. 2018;**25**:e12427

[28] Perez-Garcia MT, Ciudad P, Lopez-Lopez JR. The secret life of ion channels: Kv1.3 potassium channels and proliferation. *American Journal of Physiology. Cell Physiology*. 2018;**314**:C27-C42

[29] Dwenger MM, Ohanyan V, Navedo MF, Nystoriak MA. Coronary microvascular Kv1 channels as regulatory sensors of intracellular pyridine nucleotide redox potential. *Microcirculation*. 2018;**25**:e12426

[30] Zhong XZ, Abd-Elrahman KS, Liao CH, El-Yazbi AF, Walsh EJ, Walsh MP, et al. Stromatoxin-sensitive, heteromultimeric Kv2.1/Kv9.3 channels contribute to myogenic control of cerebral arterial diameter. *J Physiol*. 2010;**588**:4519-4537

- [31] Jackson WF. Kv channels and the regulation of vascular smooth muscle tone. *Microcirculation*. 2018;**25**:e12421
- [32] Knot HJ, Nelson MT. Regulation of membrane potential and diameter by voltage-dependent K<sup>+</sup> channels in rabbit myogenic cerebral arteries. *The American Journal of Physiology*. 1995;**269**:H348-H355
- [33] Knot HJ, Nelson MT. Regulation of arterial diameter and wall [Ca<sup>2+</sup>] in cerebral arteries of rat by membrane potential and intravascular pressure. *The Journal of Physiology*. 1998;**508**(Pt 1):199-209
- [34] Dabertrand F, Kroigaard C, Bonev AD, Cognat E, Dalsgaard T, Domenga-Denier V, et al. Potassium channelopathy-like defect underlies early-stage cerebrovascular dysfunction in a genetic model of small vessel disease. *Proceedings of the National Academy of Sciences of the United States of America*. 2015;**112**:E796-E805
- [35] Amberg GC, Rossow CF, Navedo MF, Santana LF. NFATc3 regulates Kv2.1 expression in arterial smooth muscle. *The Journal of Biological Chemistry*. 2004;**279**:47326-47334
- [36] Clement-Chomienne O, Walsh MP, Cole WC. Angiotensin II activation of protein kinase C decreases delayed rectifier K<sup>+</sup> current in rabbit vascular myocytes. *The Journal of Physiology*. 1996;**495**(Pt 3):689-700
- [37] Mackie AR, Brueggemann LI, Henderson KK, Shiels AJ, Cribbs LL, Scrogin KE, et al. Vascular KCNQ potassium channels as novel targets for the control of mesenteric artery constriction by vasopressin, based on studies in single cells, pressurized arteries, and in vivo measurements of mesenteric vascular resistance. *The Journal of Pharmacology and Experimental Therapeutics*. 2008;**325**:475-483
- [38] Kidd MW, Bulley S, Jaggar JH. Angiotensin II reduces the surface abundance of KV 1.5 channels in arterial myocytes to stimulate vasoconstriction. *The Journal of Physiology*. 2017;**595**:1607-1618
- [39] Clement-Chomienne O, Walsh MP. Identification of protein kinase C isoenzymes in smooth muscle: Partial purification and characterization of chicken gizzard PKC zeta. *Biochemistry and Cell Biology*. 1996;**74**:51-65
- [40] Ko EA, Park WS, Firth AL, Kim N, Yuan JX, Han J. Pathophysiology of voltage-gated K<sup>+</sup> channels in vascular smooth muscle cells: Modulation by protein kinases. *Progress in Biophysics and Molecular Biology*. 2010;**103**:95-101
- [41] Rainbow RD, Norman RI, Everitt DE, Brignell JL, Davies NW, Standen NB. Endothelin-I and angiotensin II inhibit arterial voltage-gated K<sup>+</sup> channels through different protein kinase C isoenzymes. *Cardiovascular Research*. 2009;**83**:493-500
- [42] Ishikawa T, Hume JR, Keef KD. Modulation of K<sup>+</sup> and Ca<sup>2+</sup> channels by histamine H1-receptor stimulation in rabbit coronary artery cells. *The Journal of Physiology*. 1993;**468**:379-400
- [43] Gelband CH, Ishikawa T, Post JM, Keef KD, Hume JR. Intracellular divalent cations block smooth muscle K<sup>+</sup> channels. *Circulation Research*. 1993;**73**:24-34
- [44] Brignell JL, Perry MD, Nelson CP, Willets JM, Challiss RA, Davies NW. Steady-state modulation of voltage-gated K<sup>+</sup> channels in rat arterial smooth muscle by cyclic AMP-dependent protein kinase and protein phosphatase 2B. *PLoS One*. 2015;**10**:e0121285

- [45] Moore CL, Nelson PL, Parelkar NK, Rusch NJ, Rhee SW. Protein kinase A-phosphorylated KV1 channels in PSD95 signaling complex contribute to the resting membrane potential and diameter of cerebral arteries. *Circulation Research*. 2014;**114**:1258-1267
- [46] Joseph BK, Thakali KM, Pathan AR, Kang E, Rusch NJ, Rhee SW. Postsynaptic density-95 scaffolding of shaker-type K(+) channels in smooth muscle cells regulates the diameter of cerebral arteries. *The Journal of Physiology*. 2011;**589**:5143-5152
- [47] Moore CL, McClenahan SJ, Hanvey HM, Jang DS, Nelson PL, Joseph BK, et al. Beta1-adrenergic receptor-mediated dilation of rat cerebral artery requires shaker-type KV1 channels on PSD95 scaffold. *Journal of Cerebral Blood Flow and Metabolism: Official Journal of the International Society of Cerebral Blood Flow and Metabolism*. 2015;**35**:1537-1546
- [48] Colledge M, Dean RA, Scott GK, Langeberg LK, Haganir RL, Scott JD. Targeting of PKA to glutamate receptors through a MAGUK-AKAP complex. *Neuron*. 2000;**27**:107-119
- [49] Nieves-Cintrón M, Nystoriak MA, Prada MP, Johnson K, Fayer W, Dell'Acqua ML, et al. Selective down regulation of Kv2.1 function contributes to enhanced arterial tone during diabetes. *Journal of Biological Chemistry*. 2015;**290**:7918-7929
- [50] Nystoriak MA, Nieves-Cintrón M, Nygren PJ, Hinke SA, Nichols CB, Chen CY, et al. AKAP150 contributes to enhanced vascular tone by facilitating large-conductance Ca<sup>2+</sup>-activated K<sup>+</sup> channel remodeling in hyperglycemia and diabetes mellitus. *Circulation Research*. 2014;**114**:607-615
- [51] Chadha PS, Zunke F, Zhu HL, Davis AJ, Jepps TA, Olesen SP, et al. Reduced KCNQ4-encoded voltage-dependent potassium channel activity underlies impaired beta-adrenoceptor-mediated relaxation of renal arteries in hypertension. *Hypertension*. 2012;**59**:877-884
- [52] Latorre R, Castillo K, Carrasquel-Ursulaez W, Sepulveda RV, Gonzalez-Nilo F, Gonzalez C, et al. Molecular determinants of BK Channel functional diversity and functioning. *Physiological Reviews*. 2017;**97**:39-87
- [53] Kotlikoff MI. Calcium-induced calcium release in smooth muscle: The case for loose coupling. *Progress in Biophysics and Molecular Biology*. 2003;**83**:171-191
- [54] Takeda Y, Nystoriak MA, Nieves-Cintrón M, Santana LF, Navedo MF. Relationship between Ca<sup>2+</sup> sparklets and sarcoplasmic reticulum Ca<sup>2+</sup> load and release in rat cerebral arterial smooth muscle. *American Journal of Physiology. Heart and Circulatory Physiology*. 2011;**301**:H2285-H2294
- [55] Zhao G, Neeb ZP, Leo MD, Pachua J, Adebisi A, Ouyang K, et al. Type 1 IP3 receptors activate BKCa channels via local molecular coupling in arterial smooth muscle cells. *The Journal of General Physiology*. 2010;**136**:283-291
- [56] Nelson MT, Brayden JE. Regulation of arterial tone by calcium-dependent K<sup>+</sup> channels and ATP-sensitive K<sup>+</sup> channels. *Cardiovascular Drugs and Therapy*. 1993;**7**(Suppl 3):605-610
- [57] Dabertrand F, Nelson MT, Brayden JE. Acidosis dilates brain parenchymal arterioles by conversion of calcium waves to sparks to activate BK channels. *Circulation Research*. 2012;**110**:285-294
- [58] Nieves-Cintrón M, Syed AU, Buonarati OR, Rigor RR, Nystoriak MA, Ghosh D, et al. Impaired BKCa channel

function in native vascular smooth muscle from humans with type 2 diabetes. *Scientific Reports*. 2017;**7**:14058

[59] Khavandi K, Baylie RL, Sugden SA, Ahmed M, Csato V, Eaton P, et al. Pressure-induced oxidative activation of PKG enables vasoregulation by Ca<sup>2+</sup> sparks and BK channels. *Science Signaling*. 2016;**9**:ra100

[60] Amberg GC, Bonev AD, Rossow CF, Nelson MT, Santana LF. Modulation of the molecular composition of large conductance, Ca<sup>2+</sup> activated K<sup>+</sup> channels in vascular smooth muscle during hypertension. *The Journal of Clinical Investigation*. 2003;**112**:717-724

[61] Jackson WF, Blair KL. Characterization and function of Ca(2+)-activated K<sup>+</sup> channels in arteriolar muscle cells. *The American Journal of Physiology*. 1998;**274**:H27-H34

[62] Krishnamoorthy G, Sonkusare SK, Heppner TJ, Nelson MT. Opposing roles of smooth muscle BK channels and ryanodine receptors in the regulation of nerve-evoked constriction of mesenteric resistance arteries. *American Journal of Physiology. Heart and Circulatory Physiology*. 2014;**306**:H981-H988

[63] Jaggar JH, Porter VA, Lederer WJ, Nelson MT. Calcium sparks in smooth muscle. *American Journal of Physiology. Cell Physiology*. 2000;**278**:C235-C256

[64] Fan G, Kassmann M, Hashad AM, Welsh DG, Gollasch M. Differential targeting and signalling of voltage-gated T-type Cav 3.2 and L-type Cav 1.2 channels to ryanodine receptors in mesenteric arteries. *The Journal of Physiology*. 2018;**596**:4863-4877

[65] Liu G, Shi J, Yang L, Cao L, Park SM, Cui J, et al. Assembly of a Ca<sup>2+</sup>-dependent BK channel signaling complex by binding to beta2 adrenergic

receptor. *The EMBO Journal*. 2004;**23**:2196-2205

[66] Longden TA, Nelson MT. Vascular inward rectifier K<sup>+</sup> channels as external K<sup>+</sup> sensors in the control of cerebral blood flow. *Microcirculation*. 2015;**22**:183-196

[67] Sancho M, Fabris S, Hald BO, Brett SE, Sandow SL, Poepping TL, et al. Membrane lipid-KIR2.x channel interactions enable hemodynamic sensing in cerebral arteries. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 2019;**39**:1072-1087

[68] Sancho M, Samson NC, Hald BO, Hashad AM, Marrelli SP, Brett SE, et al. KIR channels tune electrical communication in cerebral arteries. *Journal of Cerebral Blood Flow and Metabolism: Official Journal of the International Society of Cerebral Blood Flow and Metabolism*. 2017;**37**:2171-2184

[69] Quayle JM, McCarron JG, Brayden JE, Nelson MT. Inward rectifier K<sup>+</sup> currents in smooth muscle cells from rat resistance- sized cerebral arteries. *The American Journal of Physiology*. 1993;**265**:C1363-C1370

[70] Tajada S, Ciudad P, Moreno-Dominguez A, Perez-Garcia MT, Lopez-Lopez JR. High blood pressure associates with the remodelling of inward rectifier K<sup>+</sup> channels in mice mesenteric vascular smooth muscle cells. *The Journal of Physiology*. 2012;**590**:6075-6091

[71] Bradley KK, Jaggar JH, Bonev AD, Heppner TJ, Flynn ER, Nelson MT, et al. Kir2.1 encodes the inward rectifier potassium channel in rat arterial smooth muscle cells. *The Journal of Physiology*. 1999;**515**:639-651

[72] Sancho M, Gao Y, Hald BO, Yin H, Boulton M, Steven DA, et al. An assessment of KIR channel function

- in human cerebral arteries. *American Journal of Physiology. Heart and Circulatory Physiology*. 2019;**316**:H794-H800
- [73] Moosmang S, Schulla V, Welling A, Feil R, Feil S, Wegener JW, et al. Dominant role of smooth muscle L-type calcium channel Cav1.2 for blood pressure regulation. *The EMBO Journal*. 2003;**22**:6027-6034
- [74] Catterall WA. Voltage-gated calcium channels. *Cold Spring Harbor Perspectives in Biology*. 2011;**3**:a003947
- [75] Navedo MF, Amberg GC, Westenbroek RE, Sinnegger-Brauns MJ, Catterall WA, Striessnig J, et al. Ca(v)1.3 channels produce persistent calcium sparklets, but Ca(v)1.2 channels are responsible for sparklets in mouse arterial smooth muscle. *Am J Physiol Heart Circ Physiol*. 2007;**293**:H1359-H1370
- [76] Bannister JP, Adebisi A, Zhao G, Narayanan D, Thomas CM, Feng JY, et al. Smooth muscle cell alpha2delta-1 subunits are essential for vasoregulation by CaV1.2 channels. *Circulation Research*. 2009;**105**:948-955
- [77] Kharade SV, Sonkusare SK, Srivastava AK, Thakali KM, Fletcher TW, Rhee SW, et al. The beta3 subunit contributes to vascular calcium channel upregulation and hypertension in angiotensin II-infused C57BL/6 mice. *Hypertension*. 2013;**61**:137-142
- [78] Reimer D, Huber IG, Garcia ML, Haase H, Striessnig J. Beta subunit heterogeneity of L-type Ca(2+) channels in smooth muscle tissues. *FEBS Letters*. 2000;**467**:65-69
- [79] Keef KD, Hume JR, Zhong J. Regulation of cardiac and smooth muscle Ca(2+) channels (Ca(V)1.2a,b) by protein kinases. *Am J Physiol Cell Physiol*. 2001;**281**:C1743-C1756
- [80] Blatter LA, Wier WG. Nitric oxide decreases [Ca<sup>2+</sup>]<sub>i</sub> in vascular smooth muscle by inhibition of the calcium current. *Cell Calcium*. 1994;**15**:122-131
- [81] Navedo MF, Amberg GC. Local regulation of L-type Ca(2)(+) channel sparklets in arterial smooth muscle. *Microcirculation*. 2013;**20**:290-298
- [82] Prada MP, Syed AU, Buonarati OR, Reddy GR, Nystoriak MA, Ghosh D, et al. A Gs-coupled purinergic receptor boosts Ca(2+) influx and vascular contractility during diabetic hyperglycemia. *eLife*. 2019;**8**:e42214
- [83] Syed AU, Reddy GR, Ghosh D, Prada MP, Nystoriak MA, Morotti S, et al. Adenylyl cyclase 5-generated cAMP controls cerebral vascular reactivity during diabetic hyperglycemia. *J Clin Invest*. 2019;**130**:3140-3152
- [84] Navedo MF, Takeda Y, Nieves-Cintrón M, Molkentin JD, Santana LF. Elevated Ca<sup>2+</sup> sparklet activity during acute hyperglycemia and diabetes in cerebral arterial smooth muscle cells. *American Journal of Physiology. Cell Physiology*. 2010;**298**:C211-C220
- [85] Nystoriak MA, Nieves-Cintrón M, Patriarchi T, Buonarati OR, Prada MP, Morotti S, et al. Ser1928 phosphorylation by PKA stimulates the L-type Ca<sup>2+</sup> channel CaV1.2 and vasoconstriction during acute hyperglycemia and diabetes. *Science Signaling*. 2017;**10**:eaaf9647
- [86] Navedo MF, Nieves-Cintrón M, Amberg GC, Yuan C, Votaw VS, Lederer WJ, et al. AKAP150 is required for stuttering persistent Ca<sup>2+</sup> sparklets and angiotensin II-induced hypertension. *Circulation Research*. 2008;**102**:e1-e11
- [87] Navedo MF, Amberg GC, Votaw VS, Santana LF. Constitutively active

L-type Ca<sup>2+</sup> channels. Proceedings of the National Academy of Sciences of the United States of America. 2005;**102**:11112-11117

[88] Santana LF, Navedo MF. Molecular and biophysical mechanisms of Ca<sup>2+</sup> sparklets in smooth muscle. *Journal of Molecular and Cellular Cardiology*. 2009;**47**:436-444

[89] Navedo MF, Amberg GC, Nieves M, Molkentin JD, Santana LF. Mechanisms underlying heterogeneous Ca<sup>2+</sup> sparklet activity in arterial smooth muscle. *The Journal of General Physiology*. 2006;**127**:611-622

[90] Navedo MF, Santana LF. CaV1.2 sparklets in heart and vascular smooth muscle. *Journal of Molecular and Cellular Cardiology*. 2013;**58**:67-76

[91] Navedo MF, Cheng EP, Yuan C, Votaw S, Molkentin JD, Scott JD, et al. Increased coupled gating of L-type Ca<sup>2+</sup> channels during hypertension and Timothy syndrome. *Circulation Research*. 2010;**106**:748-756

[92] Amberg GC, Navedo MF, Nieves-Cintrón M, Molkentin JD, Santana LF. Calcium sparklets regulate local and global calcium in murine arterial smooth muscle. *The Journal of Physiology*. 2007;**579**:187-201

[93] Nieves-Cintrón M, Amberg GC, Navedo MF, Molkentin JD, Santana LF. The control of Ca<sup>2+</sup> influx and NFATc3 signaling in arterial smooth muscle during hypertension. *Proceedings of the National Academy of Sciences of the United States of America*. 2008;**105**:15623-15628

[94] Abd El-Rahman RR, Harraz OF, Brett SE, Anfinogenova Y, Mufti RE, Goldman D, et al. Identification of L- and T-type Ca<sup>2+</sup> channels in rat cerebral arteries: Role in myogenic tone development. *American Journal*

*of Physiology. Heart and Circulatory Physiology*. 2013;**304**:H58-H71

[95] Harraz OF, Abd El-Rahman RR, Bigdely-Shamloo K, Wilson SM, Brett SE, Romero M, et al. Ca(V)<sub>3.2</sub> channels and the induction of negative feedback in cerebral arteries. *Circulation Research*. 2014;**115**:650-661

[96] Harraz OF, Visser F, Brett SE, Goldman D, Zechariah A, Hashad AM, et al. CaV1.2/CaV3.X channels mediate divergent vasomotor responses in human cerebral arteries. *The Journal of General Physiology*. 2015;**145**:405-418

[97] Harraz OF, Welsh DG. T-type Ca<sub>2</sub>(+) channels in cerebral arteries: Approaches, hypotheses, and speculation. *Microcirculation*. 2013;**20**:299-306

[98] Kuo IY, Wolfle SE, Hill CE. T-type calcium channels and vascular function: The new kid on the block? *The Journal of Physiology*. 2011;**589**:783-795

[99] Harraz OF, Brett SE, Welsh DG. Nitric oxide suppresses vascular voltage-gated T-type Ca<sup>2+</sup> channels through cGMP/PKG signaling. *American Journal of Physiology. Heart and Circulatory Physiology*. 2014;**306**:H279-H285

[100] Harraz OF, Welsh DG. Protein kinase a regulation of T-type Ca<sup>2+</sup> channels in rat cerebral arterial smooth muscle. *Journal of Cell Science*. 2013;**126**:2944-2954