

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,500

Open access books available

136,000

International authors and editors

170M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Chapter

Hot Pepper (*Capsicum annuum* L.): An Alternative Food to Reduce Micronutrient Deficiencies in Human

S.R. Krishna Motukuri and Nallamothu Jaswanthi

Abstract

Hot peppers are good source of bioactive compounds particularly phytochemicals, capsaicin, oleoresins, and rich organic micronutrients with potential health benefits. It plays an important role in diets and possesses micronutrients. Micronutrient dietary deficiency remains a massive problem in the world which may cause several chronic health diseases. More than 2 billion individuals are facing micronutrient deficiencies, viz., zinc, iodine, and iron, followed by vitamins. Among various approaches to overcome human nutrition deficiencies, a diet with fruits and vegetables that are rich in micronutrients is considered as the best solution. Hot pepper consumption worldwide is well known for its high nutritional content which in turn reduces human micronutrient deficiencies. Thus sufficient amounts of micronutrients can be provided by incorporating the nutrient-rich pepper in diet that could beneficially help in combating nutrient deficiencies. Thus, in the present chapter, an overview of the micronutrient deficiency is described, and the nutrition status of hot pepper that utilized in eradicating human micronutrient dietary deficiencies is also discussed.

Keywords: *Capsicum*, hot pepper, micronutrients, human dietary deficiency, advanced approaches

1. Introduction

Micronutrients (minerals and vitamins) are obtained from our diet; though required in small amounts, they are necessary for good growth and development [1]. Micronutrients, namely, iron, cobalt, manganese, boron, molybdenum, chlorine, nickel, zinc, copper, iodine, fluorine, and chromium, are essential for human health. Many studies showed that nutrition plays a vital role in providing the substances essential for the formation of the early brain structure and also supporting for its good functioning [2]. Micronutrients play a unique metabolic role as cofactors in the absorption and digestion of carbohydrates, proteins, amino acids, and lipids to release energy [3]. Micronutrients play a very important role in all the metabolic activities like cell signaling, motility, cell division, and cell differentiation and death and, in turn, regulate the tissue development [4].

Micronutrient deficiencies can badly affect the health of individuals, which influence the world's morbidity and mortality [5]. These deficiencies are referred

to as “hidden hunger” which affects about 2 billion people [6] and may lead to high rates of illness, obesity, underdevelopment, and even deaths in children of age below 5 years [7]. Among various vegetables and fruits that are rich in micronutrients, chilies have various essential nutrients, minerals, vitamins (A, C, E), and other important phytochemical compounds [8]. They also pose some human health benefits like preventing diseases such as obesity, heart diseases, and different cancers [9]. An alkaloid, namely, capsaicin, is present in pepper which has antimicrobial, anti-inflammatory, and anticancer effects on the digestive system and is used in relieving pain and also to lose weight [10]. Thus providing diet that is rich in pepper can be supportive in an ongoing quest to ease micronutrient deficiencies.

2. Micronutrient deficiency in humans

Some studies reveal the importance of micronutrients in pregnant women and generally given as supplements, which include vitamins, viz., A, cobalamin, pyridoxine, C, D, and E, iron, zinc, iodine, copper, and selenium. Micronutrient deficiencies may mimic radiation or some chemicals which damage the nucleic acids and lead to cancer [11]. Some studies reveal that these deficiencies are related with the danger of HIV infection progression and deaths [12]. Deficiency of vitamin A may cause low serum level of retinol which affects nearly 15% of pregnant women and which leads to night blindness in 8% of them [4]. The reduced form of vitamin C [ascorbic acid (AA)] is the principal chemical structure that appears as an important micronutrient that involves in various physiological functions. Ascorbic acid plays a vital role in reducing the antioxidants and enhances the production of ROS to prevent cancer [13]. Vitamin D is generally known as fat-soluble compound with antiproliferative effect and also involved in the development of bone and immune system [14]. Some studies reported that vitamin D metabolites help in protection against cancer [15].

Vitamin E deficiency can lead to enhanced peroxidation which leads to symptoms like walking difficulties and severe development of speech, reduced fat absorption syndrome, and lipoprotein abnormalities [16]. Deficiencies of iron may affect the growth and mental development and also decreases the capability to do physical work [3]. Some studies reported that the supplementation of zinc along with other micronutrients can reduce the severity of diarrhea diseases and respiratory infections [17]. These deficiencies occur in individuals who do not consume food that provides micronutrients sufficiently like fortified foods, fruits, and animal products. This is usually due to its excessive cost or may domestically unreachable. Among the various vegetables available, chili (*Capsicum annuum*), which are available to common man at affordable price and provide various vitamins like E, C, B6, B12, and provitamin A and some minerals, possibly will give rise to significantly enhanced nutrition [8].

3. Role of hot pepper in micronutrient deficiencies

To enhance the micronutrient status in humans, nutritionists suggest integrating foods, which are rich in micronutrients, in diet. Among various vegetables, chili fruits are rich in capsaicinoids, carotenoids, tocopherols, provitamin A, ascorbic acid, and several antioxidants. Chilies are also an excellent source of xanthophylls and vitamins B1 (thiamine), B3 (niacin), and P (citric) [18]. Hot pepper contains proteins (1.9 g), sugars (5.3 g), fiber (1.5 g), fat (0.4 g), energy (8.8 g), ascorbic acid (240%), pyridoxine (39%), vitamin A (32%), copper (14%), iron (13%), potassium (7%), and magnesium (6%) [19]. Nutrient constituents and their composition present in chili species in different countries are represented in **Table 1**. Red sweet


| So. no | Constituent | Name of the country | Fruit species | Composition | References |
|--------|---------------|---------------------|----------------------------------|---------------------------|------------|
| 1 | Vitamin C | United States | <i>C. Francisca</i> | 122.0 mg/100 g DW | [20] |
| | | Brazil | <i>C. chinense</i> | 125 mg/100 g DW | [21] |
| | | India | <i>Capsicum chinense</i> Jacquin | 109.36 mg/100 g DW | [22] |
| | | Ethiopia | <i>C. annuum</i> | 84.011–89.011 mg/100 g DW | [23] |
| 2 | Capsaicin | Brazil | <i>C. chinense</i> | 14.0 mg/g DW | [24] |
| | | Central America | <i>C. annuum</i> | 0.042 mg/g DW | [25] |
| | | India | <i>C. frutescens</i> | 4.45 mg/g DW | [26] |
| | | Ethiopia | <i>C. annuum</i> (Marako fana) | 5.5 mg/g DW | [27] |
| 3 | Crude protein | America | <i>C. annuum</i> | 2.9 g/100 g DW | [28] |
| | | India | <i>C. chinense</i> | 17.5 g/100 g DW | [22] |
| | | Ethiopia | <i>C. annum</i> (Marako fana) | 118.09 g/100 g DW | [29] |
| | | Brazil | <i>C. chinense</i> | 0.08–4.5 g/100 g DW | [21] |
| 4 | Carbohydrates | America | <i>C. annuum</i> | 3.0 g/100 g DW | [28] |
| | | India | <i>C. chinense</i> | 78.1 g/100 g DW | [22] |
| | | Brazil | <i>C. chinense</i> | 1.8–10.8 g/100 g DW | [21] |
| | | Ethiopia | <i>C. annum</i> (Marako fana) | 35.3 ± 0.6 g/100 g DW | [29] |

Table 1.
 Nutrient constituents and their composition in chili species in different countries.

pepper consists of twice the provitamin A than in carrot and double the amount of vitamin C of green pepper, and it also acts as antioxidant and anti-inflammatory phytonutrient [30]. Sweet pepper is fibrous in nature and is the sources of dietary fiber which could help in fighting blood cholesterol level and diabetes better from the body and bloodstream, respectively [31].

4. Medicinal uses of hot pepper

Capsaicin is a pungent alkaloid present in hot pepper species, and it is the principal capsaicinoid that accounts for about 71% of the pungent types, followed by dihydrocapsaicin [32]. Structures of capsaicin and dihydrocapsaicin are in **Figure 1**. The percentage of pungency varies among species by varying the capsaicin and dihydrocapsaicin contents which are due to influence of factors like the developmental stages of fruit and the environmental conditions [33]. Capsaicin possesses good medicinal properties like antimicrobial, anticancer, antidiabetic, and analgesic properties. Some studies reveal that fruits having higher amounts

**Figure 1.**

(A) Capsaicin. (B) Dihydrocapsaicin.

of capsaicin show high antioxidant level [32]. Besides the nutritional benefits, it is also responsible for medical therapeutic and pharmacological uses [34]. Capsaicin has an important role in relieving sore throats, fever, and cold symptoms and also enhances the circulation of blood and strengthens the arteries, thus reducing the risk of heart attacks [3].

Capsaicin is also used as an anti-irritant balm for external use and as painkiller in some topical ointments, nasal sprays, and skin patches and also used in the form of cream for short-term relaxation of muscles and pains related to arthritis, back pain, and other stresses [35]. It also has an antidiabetic activity by enhancing the insulin secretion and releasing bound insulin. Capsaicin helps in producing heat within the body and consumes oxygen which simultaneously burns calories in the body and helps in losing weight [30]. Thus, incorporating pepper-rich food in everyday diet can be useful in enduring pursuit of eliminating micronutrient deficiency. Hot pepper fruits were enriched with antioxidants which protect the human body from harmful effects of free radicals, and it develops resistance against different diseases.

5. Conclusion

Advanced technologies are being used to combat human micronutrient deficiency in developing countries, where poor people are mostly at this risk. In tropical countries, rural farmers traditionally cultivate wide range of vegetables; among them chili is grown globally, and India is the most important producer and exporter of hot pepper which consists of wide range of micro- and macronutrients. In this chapter, it has been concluded that the chilies are cost-effective vegetables that are significantly rich in essential micro- and macronutrients and also possess healthy components to support human health conditions. The presence of antioxidants and phytochemicals in chili increases its importance in controlling the diseases. Integrating hot pepper in our regular diet can be supportive to alleviate nutrient deficiency in humans.

Acknowledgements

The authors thank the head of the department, Department of Biotechnology, Koneru Lakshmaiah Education Foundation deemed to be University, Guntur, Andhra Pradesh, India, for providing support in writing this book chapter.

Conflict of interest

The authors declare that there is no conflict of interest on this book chapter.

IntechOpen

IntechOpen

Author details

S.R. Krishna Motukuri* and Nallamothu Jaswanthi
Department of Biotechnology, Koneru Lakshmaiah Education Foundation,
Guntur, Andhra Pradesh, India

*Address all correspondence to: msrkrishna81@gmail.com

IntechOpen

© 2020 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] McMillan DC, Maguire D, Talwar D. Relationship between nutritional status and the systemic inflammatory response: Micronutrients. *Proceedings of the Nutrition Society*. 2019;**78**:56-67. DOI: 10.1017/S0029665118002501
- [2] Prado EL, Dewey KG. Nutrition and brain development in early life. *Nutrition Reviews*. 2014;**72**:267-284. DOI: 10.1111/nure.12102
- [3] Olatunji Tomi L, Anthony J. Afolayan. The suitability of chili pepper (*Capsicum annuum* L.) for alleviating human micronutrient dietary deficiencies: A review. *Food Science & Nutrition*. 2018;**6**:2239-2251. DOI: 10.1002/fsn3.790
- [4] Gernand AD, Schulze KJ, Stewart CP, West KP, Christian P. Micronutrient deficiencies in pregnancy worldwide: Health effects and prevention. *Nature Reviews. Endocrinology*. 2016;**12**:274-289. DOI: 10.1038/nrendo.2016.37
- [5] Shivakoti R, Christian P, Yang WT, Gupte N, Mwelase N, Kanyama C, et al. Prevalence and risk factors of micronutrient deficiencies pre- and post-antiretroviral therapy (ART) among a diverse multicountry cohort of HIV-infected adults. *Clinical Nutrition*. 2016;**35**:183-189. DOI: 10.1016/j.clnu.2015.02.002
- [6] Bailey RL, West KP, Black RE. The epidemiology of global micronutrient deficiencies. *Annals of Nutrition & Metabolism*. 2015;**66**:22-33. DOI: 10.1159/000371618
- [7] Global Nutrition Report. *Global Nutrition Report 2017: Nourishing the SDGs* (Development Initiatives, 2017)
- [8] Materska M, Perucka I. Antioxidant Activity of the Main Phenolic Compounds Isolated from Hot Pepper Fruit (*Capsicum annuum* L.). *Journal of Agricultural and Food Chemistry*. 2005;**53**:1750-1756. DOI: 10.1021/jf035331k
- [9] Alzrog AM, Mohamed AS, Zakaria RB, Alias AKB. Effect of planting media (rice husk and coco peat) on the uptake of cadmium and some micronutrients in chilli (*Capsicum annuum* l.). *Pure and Applied Biology*. 2013;**2**:76. DOI: 10.19045/bspab.2013.23001
- [10] Srinivasan K. Biological activities of red pepper (*Capsicum annuum*) and its pungent principle capsaicin: A review. *Critical Reviews in Food Science and Nutrition*. 2015;**56**:1488-1500. DOI: 10.1080/10408398.2013.772090
- [11] Ames BN. DNA damage from micronutrient deficiencies is likely to be a major cause of cancer. *Mutation Research, Fundamental and Molecular Mechanisms of Mutagenesis*. 2001;**475**:7-20. DOI: 10.1016/S0027-5107(01)00070-7
- [12] Drain PK, Kupka R, Mugusi F, Fawzi WW. Micronutrients in HIV-positive persons receiving highly active antiretroviral therapy. *The American Journal of Clinical Nutrition*. 2007;**85**:333-345. DOI: 10.1093/ajcn/85.2.333
- [13] Heaney ML, Gardner JR, Karasavvas N, Golde DW, Scheinberg DA, Smith EA, et al. Vitamin C antagonizes the cytotoxic effects of antineoplastic drugs. *Cancer Research*. 2008;**68**:8031-8038. DOI: 10.1158/0008-5472.CAN-08-1490
- [14] Gandini S, Raimondi S, Gnagnarella P, Dore JF, Maisonneuve P, Testori A. Vitamin D and skin cancer: A meta-analysis. *European Journal of Cancer*. 2009;**45**:634-641. DOI: 10.1016/j.ejca.2008.10.003

- [15] Deeb KK, Trump DL, Johnson CS. Vitamin D signalling pathways in cancer: Potential for anticancer therapeutics. *Nature Reviews Cancer*. 2007;7:684-700. DOI: 10.1038/nrc2196
- [16] Mamede AC, Tavares SD, Abrantes AM, Trindade J, Maia JM, Botelho MF. The role of vitamins in Cancer: A review. *Nutrition and Cancer*. 2011;63:479-494. DOI: 10.1080/01635581.2011.539315
- [17] Bhutta ZA. Iron and zinc deficiency in children in developing countries. *British Medical Journal*. 2007;334:104-105. DOI: 10.1136/bmj.39094.513924.BE
- [18] Bosland PW, Votava EJ. Peppers: Vegetable and Spice Capsicums. Wallingford, CT: CABI Publishing; 2000. p. 204
- [19] Ebong PE. Quality and Safety of Medicinal Plants in Phytomedicine: Panacea for Primary Health Care Delivery in Nigeria. Nigeria: University of Calabar Press; 2015. pp. 293-310
- [20] Teodoro AF, Alves RD, Ribeiro LB, Reis K, Reifschneider FJ, Fonseca ME, et al. Vitamin C content in Habanero pepper accessions (*Capsicum chinense*). *Horticultura Brasileira*. 2013;31:59-62. DOI: 10.1590/S0102-05362013000100009
- [21] Da Silveira A-CT, da Silva GI, de Melo LAMP, Reifschneider FJB, da Costa Ribeiro CS. Carotenoid and total vitamin C content of peppers from selected Brazilian cultivars. *Journal of Food Composition and Analysis*. 2017;57:73-79. DOI: 10.1016/j.jfca.2016.12.020
- [22] Malakar S, Sarkar S, Kumar N, Jaganmohan R. Studies of biochemical characteristics and identification of active phyto-compounds of king chili (*Capsicum chinense Jacq.*) using GC-MS. *Journal of Pharmacognosy and Phytochemistry*. 2018;7:3100-3104
- [23] Kinf E. Nutritional composition, physicochemical and functional properties of some capsicum varieties grown in Ethiopia [Doctoral dissertation]. Addis Ababa University. 2009. Available from: //localhost:80/xmlui/handle/123456789/8644
- [24] Rêgo ER, Finger FL, Rêgo MM. Types, uses and fruit quality of Brazilian chili peppers. In: Kralis JF, editor. *Spices: Types, Uses and Health Benefits*. Nova Science Publishers; 2011. 70p
- [25] Cázares-Sánchez E, Ramírez-Vallejo P, Castillo-González F, Soto-Hernández RM, Rodríguez-González MT, Chávez-Servia JL. Capsaicinoids and preference of use in different morphotypes of chili peppers (*Capsicum annum L.*) of east-Central Yucatán. *Agrociencia*. 2005;1:627-638
- [26] Tilahun S, Paramaguru P, Rajamani K. Capsaicin and ascorbic acid variability in chilli and paprika cultivars as revealed by HPLC analysis. *Journal of Plant Breeding and Genetics*. 2013;1:85-89
- [27] Aklilu Alemu SH. Diversity, disease reaction, heterosis and combining ability in hot pepper (*Capsicum annum L.*) genotypes for agro-morphological and quality traits in ethiopia [Doctoral dissertation]. Asela University. 2018. Available from: //hdl.handle.net/123456789/2253
- [28] Kumar S, Kumar R, Singh J. Cayenne/American pepper. In: *Handbook of Herbs and Spices*. Vol. 3. Cambridge, UK: Woodhead Publishing; 2006. pp. 299-312. DOI: 10.1533/9781845691717.3.299
- [29] Esayas K, Shimelis A, Ashebir F, Negussie R, Tilahun B, Gulelat D. Proximate composition, mineral content and antinutritional factors of some capsicum (*Capsicum annum*) varieties grown in Ethiopia. *Bulletin of the Chemical Society of Ethiopia*. 2011;25. DOI: 10.4314/bcse.v25i3.68602

[30] Mercy ER, David U. Potential health benefits of conventional nutrients and phytochemicals of Capsicum peppers. *Pharmacy & Pharmacology International Journal*. 2018;**6**:62-69. DOI: 10.15406/ppij.2018.06.00157

[31] Mateljan G. *The World's Healthiest Foods*. USA: The George Mateljan foundation; 2018

[32] Chu CJ, Huang SM, de Petrocellis L, Bisogno T, Ewing SA, Miller JD, et al. N- Oleoyldopamine, a novel endogenous capsaicin- like lipid that produces hyperalgesia. *The Journal of Biological Chemistry*. 2003;**278**:13633-13639. DOI: 10.1074/jbc.M211231200

[33] Estrada B, Bernal MA, Díaz J, Pomar F, Merino F. Capsaicinoids in vegetative organs of *Capsicum annum* L. in relation to fruiting. *Journal of Agricultural and Food Chemistry*. 2002;**50**:1188-1191. DOI: 10.1021/jf011270j

[34] Peña-Alvarez A, Ramírez-Maya E, Alvarado-Suárez LA. Analysis of capsaicin and dihydrocapsaicin in peppers and pepper sauces by solid phase microextraction – Gas chromatography – Mass spectrometry. *Journal of Chromatography. A*. 2009;**1216**:2843-2847. DOI: 10.1016/j.chroma.2008.10.053

[35] Fattori V, Hohmann MS, Rossaneis AC, Pinho-Ribeiro FA, Verri WA. Capsaicin: Current understanding of its mechanisms and therapy of pain and other pre-clinical and clinical uses. *Molecules*. 2016;**21**:844. DOI: 10.3390/molecules21070844